

Schmid, Timo; Tzavidis, Nikos; Münnich, Ralf; Chambers, Ray

Working Paper

Outlier robust small area estimation under spatial correlation

Diskussionsbeiträge, No. 2015/8

Provided in Cooperation with:

Free University Berlin, School of Business & Economics

Suggested Citation: Schmid, Timo; Tzavidis, Nikos; Münnich, Ralf; Chambers, Ray (2015) : Outlier robust small area estimation under spatial correlation, Diskussionsbeiträge, No. 2015/8, Freie Universität Berlin, Fachbereich Wirtschaftswissenschaft, Berlin

This Version is available at:

<https://hdl.handle.net/10419/107683>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

Outlier robust small area estimation under spatial correlation

Timo Schmid
Nikos Tzavidis
Ralf Münnich
Ray Chambers

School of Business & Economics

Discussion Paper

Economics

2015/8

Outlier robust small area estimation under spatial correlation

Timo Schmid^{*}, Nikos Tzavidis^{**}, Ralf Münnich^{***}, and Ray Chambers^{****}

^{*}Institute of Statistics and Econometrics, Freie Universität Berlin, Berlin, Germany

^{**}Southampton Statistical Sciences Research Institute, University of Southampton, Southampton, UK

^{***}Economic and Social Statistics Department, University of Trier, Trier, Germany

^{****}National Institute for Applied Statistics Research Australia, University of Wollongong, Wollongong, Australia

Abstract

Modern systems of official statistics require the estimation and publication of business statistics for disaggregated domains, for example, industry domains and geographical regions. Outlier robust methods have proven to be useful for small area estimation. Recently proposed outlier robust model-based small area methods assume, however, uncorrelated random effects. Spatial dependencies, resulting from similar industry domains or geographic regions, often occur. In this paper we propose outlier robust small area methodology that allows for the presence of spatial correlation in the data. In particular, we present a robust predictive methodology that incorporates the potential spatial impact from other areas (domains) on the small area (domain) of interest. We further propose two parametric bootstrap methods for estimating the mean-squared error. Simulations indicate that the proposed methodology may lead to efficiency gains. The paper concludes with an illustrative application by using business data for estimating average labour costs in Italian provinces.

Keywords: Bias correction, projective and predictive estimators, spatial correlation, business surveys

1 Introduction

An important set of outputs for National Statistical Institutes (NSIs) are enterprise and trade statistics derived from business survey data. This has recently extended to small area/domain estimates for business statistics. During the last two decades there has been substantial growth in the development and application of model-based small area methods. However, small area estimation using business survey data presents a number of challenges. In this paper we focus on small area estimation when there are representative outliers (Chambers, 1986) in the sample data.

Outliers can invalidate the conventional parametric assumptions of Gaussian random effects and unit level errors of model-based methods, which in turn can impact the validity of model-based small area estimators. Sinha and Rao (2009) proposed a robust methodology for estimating the parameters of the nested error regression model that controls for the influence of outliers. An alternative approach that also uses influence functions to control the impact of outliers but is based on a different model was proposed

by Chambers and Tzavidis (2006). Both outlier robust estimators of Sinha and Rao (2009) and Chambers and Tzavidis (2006) are plug-in estimators. This implicitly assumes that all non-sample values follow a well-behaved working model, so that their sum can be predicted using an outlier robust fit of this working model to the outlier contaminated sample data. Chambers et al. (2014) refer to such methods as robust projective. Robust projective methods typically lead to biased estimators with lower variances. This is due to the fact that it is extremely unlikely that the non-sampled values in the target population are drawn from a distribution with the same mean as the sample non-outliers. To correct for this bias, Dongmo-Jiongo et al. (2013) and Chambers et al. (2014) proposed bias-corrected robust small area estimators with either local (area-specific) or global correction terms. Such bias-corrected estimators are referred to by Chambers et al. (2014) as robust predictive. The methodology used for building these robust predictive estimators goes back to the seminal work of Chambers (1986) and Welsh and Ronchetti (1998), in the sense that the local correction is obtained when the robust estimator is defined as the expected value of the small area mean under the outlier robust distribution function proposed by Welsh and Ronchetti (1998), while the global correction is motivated by extending the outlier robust population mean estimator of Chambers (1986) to a mixed effects working model.

In this paper we extend these ideas to outlier robust small area estimation under spatial correlation. Borrowing strength over space in small area estimation has been shown to offer some efficiency gains. One approach considers the use of geographically weighted regression (cf. Chandra et al., 2012 and Salvati et al., 2012), but this usually assumes that geo-referenced data at the individual level is available. This is hardly ever the case in business survey data. Another approach extends the nested error regression model by allowing for spatially correlated random effects that follow a simultaneously autoregressive (SAR) process (cf. Pratesi and Salvati, 2008). However, none of these methods considered estimation in the presence of outliers.

The approach we follow in this paper assumes a nested error regression model with correlated random effects that follow a SAR process. After reviewing the current literature on outlier robust small area estimation in Section 2, we present robust projective and robust predictive small area estimators under spatial correlation in Section 3. Mean Squared Error (MSE) estimation is studied in Section 4 by using two parametric bootstrap schemes that account for the spatial structure. The first scheme is based on a modified version of the parametric bootstrap proposed by Sinha and Rao (2009), whereas the second one is grounded on a modified version of the parametric bootstrap proposed by Dongmo-Jiongo et al. (2013). In Section 5 we empirically evaluate the proposed methodology by using a Monte-Carlo simulation under a range of scenarios. In Section 6 we apply the proposed methodology to business survey data from Italy. We conclude the paper by summarising our main findings and by providing some ideas for further research.

2 Outlier robust small area estimation

In this section we review the current literature on outlier robust small area estimation. We assume that the population of interest U of size N is divided into m non-overlapping small areas of sizes N_i and that

individual level data is available. The variable of interest is denoted by y_j and is assumed to be linearly related to a set of individual level auxiliary variables x_j and a set of area level covariates z_j . The sample s of size n is selected from the population by using a sampling design which is non-informative given x_j and z_j . The population is divided into n sampled and $N - n$ non-sampled units, indexed by s and r respectively. We use the subscript i to indicate the restriction to the specific area i , i. e., s_i (r_i) stands for the set of sampled (non-sampled) units from area i . Furthermore, we assume that all small areas are sampled and that the small area population averages of \bar{x}_i and \bar{z}_i for $i = 1, \dots, m$ are known.

The industry standard approach to model-based small area estimation with unit level data uses the Battese, Harter and Fuller model (cf. Battese et al., 1988). This is a nested error regression model with area-specific random effects:

$$\begin{aligned} \mathbf{y} &= \mathbf{X}\boldsymbol{\beta} + \mathbf{Z}\mathbf{v} + \mathbf{e}, \\ \mathbf{v} &\sim N(0, \mathbf{G}), \\ \boldsymbol{\varepsilon} &\sim N(0, \mathbf{R}). \end{aligned} \tag{1}$$

In (1) $\boldsymbol{\beta}$ is the $k \times 1$ vector of regression parameters and \mathbf{v} is the $(m \times 1)$ vector of area specific random effects. The unit level errors \mathbf{e} are assumed to be independently normally distributed with mean 0 and covariance $\mathbf{R} = \sigma_e^2 \mathbf{I}_N$. The covariance matrices \mathbf{G} and \mathbf{R} depend on a vector of variance parameters $\boldsymbol{\theta} = \{\sigma_v^2, \sigma_e^2\}$. When the area effects are assumed to be independent, the covariance matrix of the random effects simplifies to $\mathbf{G} = \sigma_v^2 \mathbf{I}_m$. Finally, since the two error terms \mathbf{v} and \mathbf{e} are independent, the covariance matrix of \mathbf{y} is given by $\mathbf{V} = \mathbf{R} + \mathbf{Z}\mathbf{G}\mathbf{Z}^T$. Let $\hat{\boldsymbol{\beta}}(\boldsymbol{\theta})$ define the Best Linear Unbiased Estimator (BLUE) of $\boldsymbol{\beta}$ and $\hat{\mathbf{v}}(\boldsymbol{\theta})$ denotes the Best Linear Unbiased Predictor (BLUP) of \mathbf{v} (cf. Henderson, 1950 or Searle, 1971). In practice, $\boldsymbol{\theta}$ is unknown and needs to be estimated by using e.g. maximum likelihood or restricted maximum likelihood estimation. This leads to the empirical best versions of $\hat{\boldsymbol{\beta}}(\hat{\boldsymbol{\theta}})$ and $\hat{\mathbf{v}}(\hat{\boldsymbol{\theta}})$. The EBLUP of the mean in small area i is then given by

$$\hat{y}_i^{EBLUP} = N_i^{-1} \left\{ \sum_{j \in s_i} y_j + \sum_{j \in r_i} (\mathbf{x}_j^T \hat{\boldsymbol{\beta}} + \hat{v}_i) \right\}. \tag{2}$$

The EBLUP (2) is optimal when the model assumptions (1) hold. However, the presence of outliers in the data can invalidate these assumptions. The impact of outliers on estimation can be controlled by replacing $\hat{\boldsymbol{\beta}}$ and $\hat{\mathbf{v}}$ by robust alternatives. One approach for achieving this was proposed by Sinha and Rao (2009). The paper suggested the use of the Huber influence function ψ for deriving robust versions of the maximum likelihood estimating equations leading to robust estimators $\hat{\boldsymbol{\beta}}^\psi$, $\hat{\boldsymbol{\theta}}^\psi$ and $\hat{\mathbf{v}}^\psi$. An outlier robust version of (2) is

$$\hat{y}_i^{REBLUP} = N_i^{-1} \left\{ \sum_{j \in s_i} y_j + \sum_{j \in r_i} (\mathbf{x}_j^T \hat{\boldsymbol{\beta}}^\psi + \hat{v}_i^\psi) \right\}. \tag{3}$$

Chambers et al. (2014) use the term robust projective to refer to estimators such as (3). Robust projective methods can be severely biased. The reason for the bias is not difficult to find. It is extremely

unlikely that the non-sampled values in the target population are drawn from a distribution with the same mean as the sample non-outliers, and yet these methods are built on precisely this assumption. Chambers (1986) recognised this dilemma and coined the concept of a representative outlier, i. e., a sample outlier that is potentially drawn from a group of population outliers, and hence, cannot be unit-weighted in estimation. He noted that representative outliers cannot be treated on the same basis in estimation as other sample data which are more consistent with the working model for the population, since such values can seriously destabilise the survey estimates. Chambers (1986) suggested an addition of an outlier robust bias correction term to a robust projective survey estimator, e. g., one based on outlier-robust estimates of the model parameters. Such bias-corrected estimators are referred to as robust predictive since they attempt to predict the contribution of the population outliers to the population parameter of interest.

Chambers et al. (2014) proposed a robust-predictive version of the REBLUP (3) based on the ideas of Chambers (1986) and Welsh and Ronchetti (1998). This estimator is

$$\hat{y}_i^{CCST} = \hat{y}_i^{REBLUP} + \left(1 - \frac{n_i}{N_i}\right) \frac{1}{n_i} \sum_{j \in s_i} \omega_i^\psi \phi_k \left\{ (y_j - \mathbf{x}_j^T \hat{\beta}^\psi - \hat{v}_i^\psi) / \omega_i^\psi \right\}, \quad (4)$$

where ϕ_k denotes a less restrictive influence function ($|\phi| \geq |\psi|$) with tuning constant $k > 0$ (cf. Tzavidis et al., 2010) and ω_i^ψ is the median absolute deviation of the residuals in area i . Note that setting $k = 0$ leads to the robust projective estimator (3). Hence, the bias-correction term depends only on local information from area i . Although estimator (4) has very good bias properties, it can suffer from high variability due to the fact that the correction term depends on the area-specific sample size which is usually fairly small. To amend this issue, Dongmo-Jiongo et al. (2013) proposed alternative robust predictive estimators that incorporate a global bias correction term which also depends on information from other small areas.

The first estimator of Dongmo-Jiongo et al. (2013) uses the ideas of Chambers (1986). The EBLUP (2) can be reformulated as a weighted linear function of the values in the sample s (cf. Royall, 1976) by

$$\hat{y}_i^{EBLUP} = \frac{1}{N_i} \sum_{j \in s} w_j^{EBLUP} y_j = (\mathbf{w}_{is}^{EBLUP})^T \mathbf{y}_s. \quad (5)$$

Dongmo-Jiongo et al. (2013) used this result and derived the following robust predictive estimator

$$\begin{aligned} \hat{y}_i^{CHAM} = & \hat{y}_i^{REBLUP} + N_i^{-1} \sum_{j \in s_i} \psi_{k_1} \{ (w_j - 1)(y_j - \mathbf{x}_j^T \hat{\beta}^\psi - \hat{v}_i^\psi) \} \\ & + N_i^{-1} \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} \psi_{k_1} \{ w_j (y_j - \mathbf{x}_j^T \hat{\beta}^\psi - \hat{v}_h^\psi) \} + N_i^{-1} \sum_{h=1}^m \psi_{k_2} \{ \varpi_h \hat{v}_h^\psi \} \end{aligned} \quad (6)$$

where

$$\varpi_h = \begin{cases} \sum_{j \in s_i} w_j - N_i, & h = i \\ \sum_{j \in s_h} w_j, & h \neq i \end{cases}$$

and ψ_k is the Huber influence function with tuning constant k . Dongmo-Jiongo et al. (2013) pointed out that the robust predictor (6) converges to the EBLUP (2) for large values of the tuning constant and to the (3) for small values of the tuning constant. A second approach by Dongmo-Jiongo et al. (2013) uses the close relationship between the concept of conditional bias (CB) and the influence function (cf. Beaumont et al., 2013). The conditional bias, originally developed by Muñoz-Pichardo et al. (1995), measures the average influence of a unit j on an estimator. It is defined by

$$B_j(y_j, v_h, \beta, \theta) = E(\hat{y}_i(\theta) - \bar{y}_i | s, y_j, v_h) \quad (7)$$

of a unit j in an area h (cf. Beaumont et al., 2013).

Using the concept of conditional bias, Dongmo-Jiongo et al. (2013) proposed a second robust predictive estimator,

$$\begin{aligned} \hat{y}_i^{CB} &= \hat{y}_i^{REBLUP} + N_i^{-1} \sum_{j \in s_i} \psi_{k_1} \{ (w_j - 1)(y_j - \mathbf{x}_j^T \hat{\beta}^\psi - \hat{v}_i^\psi) \} \\ &\quad + N_i^{-1} \sum_{\substack{h \neq i \\ h=1}}^m \sum_{j \in s_h} \psi_{k_1} \{ w_j (y_j - \mathbf{x}_j^T \hat{\beta}^\psi - \hat{v}_h^\psi) \} + N_i^{-1} \sum_{h=1}^m \varpi_h \hat{v}_h^\psi. \end{aligned} \quad (8)$$

3 Robust projective and predictive estimators with spatially correlated random effects

The robust projective estimators (3) of Sinha and Rao (2009) and the robust predictive estimators (4), (6) or (8) of Chambers et al. (2014) and Dongmo-Jiongo et al. (2013) have all been developed under a nested error regression model that assumes uncorrelated random effects. Recent literature has demonstrated that in some cases borrowing strength over space can improve the efficiency of small area estimators. In this section we define robust projective and robust predictive estimators in the presence of correlated random effects.

One way to model spatial correlation is to allow for correlated random effects. We extend the nested error regression model along the lines of the work by Petrucci et al. (2005) and Pratesi and Salvati (2008) assuming a SAR process. The SAR model is defined by

$$\mathbf{y} = \mathbf{X}\beta + \mathbf{Z}\mathbf{v} + \mathbf{e}, \quad (9)$$

where the covariance matrices of \mathbf{v} and \mathbf{e} are defined as in (1) but with

$$\mathbf{G} = \sigma_v^2 ((\mathbf{I} - \rho \mathbf{W}^T)(\mathbf{I} - \rho \mathbf{W}))^{-1} \quad (10)$$

$$\mathbf{R} = \text{diag}(\sigma_e^2) \quad (11)$$

respectively. The parameter ρ describes the strength of the spatial correlation and \mathbf{W} is a given contiguity matrix that describes neighbourhood structure between the small areas. Under (9) an estimator of the

small area mean is defined by

$$\hat{y}_i^{SEBLUP} = N_i^{-1} \sum_{j \in s} w_j^{sp} y_j = N_i^{-1} (\mathbf{w}_{is}^{sp})^T \mathbf{y}_s. \quad (12)$$

The weights \mathbf{w}_{is}^{sp} are given by

$$(\mathbf{w}_{is}^{sp})^T = (\mathbf{1}_s^T + (N_i - n_i)(\bar{\mathbf{x}}_{ri}^T \mathbf{A} + \bar{\mathbf{z}}_{ri}^T \mathbf{B}(\mathbf{I}_s - \mathbf{X}_s \mathbf{A}))) , \quad (13)$$

where

$$\begin{aligned} \mathbf{A} &= (\mathbf{X}_s^T \mathbf{V}^{-1} \mathbf{X}_s)^{-1} \mathbf{X}_s^T \mathbf{V}^{-1} \\ \mathbf{B} &= (\mathbf{Z}^T \mathbf{R}^{-1} \mathbf{Z} + \mathbf{G}^{-1})^{-1} \mathbf{Z}^T \mathbf{R}^{-1} \end{aligned}$$

and $\mathbf{1}_s$ is a vector of size n with

$$\mathbf{1}_s = \begin{cases} 1, & j \in s_i \\ 0, & j \in s_h \end{cases}.$$

As we saw in Section 2, outliers can severely impact upon the estimation of model (9). In order to control their influence, a robust projective estimator of the small area mean under (9) uses robust ML equations for the estimation of β , θ , \mathbf{v} and ρ leading to

$$\begin{aligned} \hat{y}_i^{SREBLUP} &= N_i^{-1} \left\{ \sum_{j \in s_i} y_j + \sum_{j \in r_i} (\mathbf{x}_j^T \hat{\beta}^{\psi, sp} + \hat{v}_i^{\psi, sp}) \right\} \\ &= N_i^{-1} \left\{ n_i \bar{y}_{si} + (N_i - n_i)(\bar{\mathbf{x}}_{ri}^T \hat{\beta}^{\psi, sp} + \bar{\mathbf{z}}_{ri}^T \hat{v}_i^{\psi, sp}) \right\}, \end{aligned} \quad (14)$$

where the superscript ψ points to the dependence of the estimator on the influence function and the superscript sp indicates that the parameters depend on the spatial correlation parameter ρ (cf. Schmid and Münnich, 2014). The robust projective estimator (14) is an extension of the Sinha and Rao estimator under (9), and hence, can suffer from the same bias problems. Thus, there is need to develop robust predictive estimators in the presence of spatial correlation.

The first robust predictive estimator we propose in this paper uses a local bias correction and is defined by

$$\begin{aligned} \hat{y}_i^{SCCST} &= \hat{y}_i^{SREBLUP} \\ &\quad + (1 - \frac{n_i}{N_i}) \frac{1}{n_i} \sum_{j \in s_i} \omega_i^{\psi, sp} \phi_k \left\{ (y_j - \mathbf{x}_j^T \hat{\beta}^{\psi, sp} - \hat{v}_i^{\psi, sp}) / \omega_i^{\psi, sp} \right\}, \end{aligned} \quad (15)$$

where $\omega_i^{\psi, sp}$ is the median absolute deviation of the residuals under the SAR model (9).

Two additional robust predictive estimators that use global bias correction terms are proposed under model (9). We start by noticing that (12) can be reformulated as follows (the detailed derivation is

available in the Appendix),

$$\begin{aligned}\hat{y}_i^{SEBLUP} &= \hat{y}_i^{SREBLUP} + N^{-1} \sum_{j \in s_i} (w_j^{sp} - 1)(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp}) \\ &\quad + N^{-1} \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} (y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp}) + N^{-1} \sum_{h=1}^m \varpi_h^{sp} \hat{v}_h^{\psi, sp},\end{aligned}\quad (16)$$

where

$$\varpi_h^{sp} = \begin{cases} \sum_{j \in s_i} w_j^{sp} - N_i, & h = i \\ \sum_{j \in s_h} w_j^{sp}, & h \neq i \end{cases}.$$

The influence of outliers on (16) can be controlled by using a a Huber influence function in the last three terms of the estimator leading to the following robust predictive estimator,

$$\begin{aligned}\hat{y}_i^{SCHAM} &= \hat{y}_i^{SREBLUP} + N_i^{-1} \sum_{j \in s_i} \psi_{k_1} \{ (w_j^{sp} - 1)(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp}) \} \\ &\quad + N_i^{-1} \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} \psi_{k_1} \{ w_j^{sp} (y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp}) \} \\ &\quad + N_i^{-1} \sum_{h=1}^m \psi_{k_2} \{ \varpi_h^{sp} \hat{v}_h^{\psi, sp} \}.\end{aligned}\quad (17)$$

The tuning constants used in the influence function are defined utilising the ideas in Dongmo-Jiongo et al. (2013). For small values of the tuning constants k_1 and k_2 the robust predictive estimator converges to the robust projective estimator (14), whereas for large tuning constants it converges to (12). A detailed discussion about the choice of the constants is given in Chambers et al. (2014) or Dongmo-Jiongo et al. (2013). Additional discussion is also provided in Section 5 of the present paper.

The second robust predictive estimator we propose in this paper is based on the concept of conditional bias, however it is developed under the SAR model. The conditional bias in this case is given by

$$B_j(y_j, v_h, \boldsymbol{\beta}, \boldsymbol{\theta}) = \begin{cases} \frac{1}{N_i} (w_j^{sp} - 1)(y_j - \mathbf{x}_j^T \boldsymbol{\beta} - v_i) + N_i^{-1} \varpi_i^{sp} v_i, & j \in s_i \\ \frac{1}{N_i} w_j^{sp} (y_j - \mathbf{x}_j^T \boldsymbol{\beta} - v_h) + N_i^{-1} \varpi_h^{sp} v_h, & j \in s_h \text{ with } h \neq i \\ -\frac{1}{N_i} w_j^{sp} (y_j - \mathbf{x}_j^T \boldsymbol{\beta} - v_i) + N_i^{-1} \varpi_i^{sp} v_i, & j \in r_i \\ \frac{1}{N_i} \varpi_h^{sp} v_h, & j \in r_h \text{ with } h \neq i \end{cases}.$$

This leads to the following robust predictive estimator

$$\begin{aligned}\hat{y}_i^{SCB} &= \hat{y}_i^{SREBLUP} + N_i^{-1} \sum_{j \in s_i} \psi_{k_1} \{ (w_j^{sp} - 1)(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp}) \} \\ &\quad + N_i^{-1} \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} \psi_{k_1} \{ w_j^{sp} (y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp}) \} + N_i^{-1} \sum_{h=1}^m \varpi_h^{sp} \hat{v}_h^{\psi, sp}.\end{aligned}\quad (18)$$

4 Mean squared error estimation

Sinha and Rao (2009) and Dongmo-Jiongo et al. (2013) have already pointed out that the estimation of the mean squared error (MSE) is a challenging problem in the case of robust predictors. In this section we propose two parametric bootstrap schemes for estimating the MSE of the proposed robust predictive small area estimators we presented in Section 3. These bootstrap MSE estimators are extended to incorporate the spatial structure in the data. Both schemes are based on the work in Hall and Maiti (2006), Sinha and Rao (2009) and Dongmo-Jiongo et al. (2013). The difference between these bootstraps is in the mechanism used for generating the bootstrap population. In particular, the first bootstrap scheme generates bootstrap realisations of the random effects \mathbf{v}^* and the error terms \mathbf{e}^* by using the non-robust estimates of the variance components (cf. Dongmo-Jiongo et al., 2013). In contrast, the second bootstrap uses robust estimates of the variance components for generating the bootstrap population (cf. Sinha and Rao, 2009). The comparison between these alternative bootstrap methods is considered for the first time in the present paper.

The steps of the bootstrap are as follows:

1. For given $\hat{\beta}^{\psi, sp}$, $\hat{\theta}$ and $\hat{\rho}$ estimated with the original sample generate area-specific random effects \mathbf{v}^* from $N(0, G(\hat{\theta}, \hat{\rho}))$ and \mathbf{e}^* from $N(0, R(\hat{\theta}))$. Use these to generate the bootstrap population and bootstrap sample,

$$\mathbf{y}^{*(b)} = X\hat{\beta}^{\psi, sp} + Z\mathbf{v}^* + \mathbf{e}^*, \quad (19)$$

and calculate the corresponding bootstrap population parameter for each area i which we denote by $\tau_i^{(b)}$.

For the first bootstrap method, we generate the bootstrap population by using non-robust estimates $\hat{\theta} = (\hat{\sigma}_v, \hat{\sigma}_e)$ and $\hat{\rho} = \hat{p}$. For the second parametric bootstrap approach we utilise robust estimates $\hat{\theta} = (\hat{\sigma}_v^\psi, \hat{\sigma}_e^\psi)$ and $\hat{\rho} = \hat{p}^\psi$. It is important to note that generation of the bootstrap population accounts for spatial correlation between the random effects.

2. Using the bootstrap sample, estimate model (9). Using the estimated model parameters from the bootstrap sample, compute the corresponding small area estimator of the population mean in area i , $\hat{\tau}_i^{(b)}$. $\hat{\tau}$ is a generic notation used to denote an estimator of the small area average.
3. Using B bootstrap samples, the MSE estimator of the corresponding estimator in area i , τ_i , is given by

$$\widehat{\text{MSE}}(\hat{\tau}_i) = \frac{1}{B} \sum_{b=1}^B \left(\hat{\tau}_i^{(b)} - \tau_i \right)^2.$$

5 Empirical evaluations

In this section, we present results from Monte-Carlo simulations that we carried out for assessing the performance of the proposed spatial robust estimators from Section 3. These estimators are compared

against those suggested by Chambers et al. (2014) and Dongmo-Jiongo et al. (2013) in Section 2. We further evaluate the performance of the MSE estimators.

Following Chambers et al. (2014), we generated population data for $m = 40$ small areas with $N_i = 100$ using a nested error regression model as follows

$$y_j = 100 + 5x_j + v_i + e_{ij}. \quad (20)$$

The covariates were generated from a log-normal-distribution with $\mu_x = 1$ and $\sigma_x = 0.5$. The unit level errors e_{ij} and the random effects v_i were generated under a range of scenarios. In particular, following Chambers et al. (2014), we focus on non-symmetric outlier contamination in the area and unit level errors. This is due to the fact that outlier robust projective estimators suffer from bias, especially under asymmetric contamination. The results in the case of symmetric contamination are available from the authors on request. Random effects and the individual errors were generated by using a mixture type contamination mechanism,

$$\begin{aligned} v_i &\sim (1 - \gamma_v) \cdot N(0, \mathbf{G}) + \gamma_v \cdot N(9, 20) \\ e_{ij} &\sim (1 - \gamma_e) \cdot N(0, 6) + \gamma_e \cdot N(10, 150), \end{aligned} \quad (21)$$

where γ_v and γ_e control the percentage of outlier contamination in the population. \mathbf{G} is defined as in (10) by

$$\mathbf{G} = \sigma_v^2 ((\mathbf{I} - \rho \mathbf{W}^T)(\mathbf{I} - \rho \mathbf{W}))^{-1}$$

using $\sigma_v^2 = 3$. We assume that the outlying areas are uncorrelated in (21). The spatial autoregressive parameter ρ is set to 0 for the non-spatial scenarios and to 0.8 for the scenarios with spatial correlation. \mathbf{W} is defined by a 8×5 first-order contiguity matrix based on a rook structure (cf. Bivand et al., 2008). According to the rook structure, neighbours are cells to the horizontal and to the vertical of a cell, but not to the diagonal. \mathbf{W} is row standardized to meet the requirements of the SAR model (cf. Petrucci et al., 2005). This leads to a row stochastic non-symmetric contiguity matrix. In the present of outliers in the data, the contamination levels are set to $\gamma_v = 0.1$ and $\gamma_e = 0.03$. We investigated three spatial and three non-spatial scenarios in total:

1. $(0, 0)$ - No contamination and no spatial correlation; $\gamma_v = \gamma_e = \rho = 0$.
2. $(v, 0)$ - Contamination only at the area level and no spatial correlation: $v_i \sim N(0, 3)$ for areas 1-36, $v_i \sim N(9, 20)$ for areas 37-40, $e_{ij} \sim N(0, 6)$ and $\rho = 0$.
3. (v, e) - Contamination at both levels and no spatial correlation: $v_i \sim N(0, 3)$ for areas 1-36, $v_i \sim N(9, 20)$ for areas 37-40, $e_{ij} \sim 0.97N(0, 6) + 0.03N(10, 150)$ and $\rho = 0$.
4. $(0, 0)_\rho$ - No contamination and spatial correlation; $\gamma_v = \gamma_e = 0$ and $\rho = 0.8$.
5. $(v, 0)_\rho$ - Contamination only at area level and spatial correlation: $v_i \sim N(0, G)$ for areas 1-36, $v_i \sim N(9, 20)$ for areas 37-40, $e_{ij} \sim N(0, 6)$ and $\rho = 0$.

6. $(v, e)_\rho$ - Contamination and spatial correlation: $v_i \sim N(0, G)$ for areas 1-36, $v_i \sim N(9, 20)$ for areas 37-40, $e_{ij} \sim 0.97N(0, 6) + 0.03N(10, 150)$ and $\rho = 0.8$.

Samples of size $n_i = 5$ were selected from the population by simple random sampling without replacement within each area. The population and sample sizes were held fixed for all areas. Each setting was repeated independently 500 times. The choice of small sample sizes is for two reasons. First, we investigate the proposed small area estimators under extreme cases. Second, the sample sizes in our application in Section 6 are approximately of similar size.

Several estimators of the small area population averages are evaluated. These are the standard EBLUP (2), which serves as a benchmark for other estimators, the spatial EBLUP (SEBLUP) (12), the robust EBLUP (REBLUP) (3) and the spatial robust EBLUP (SREBLUP) (14). Furthermore, the robust predictive estimators CCST (4), CHAM (6) and CB (8) are evaluated alongside their proposed spatial extensions SCCST (15), SCHAM (17) and SCB (18). Following Dongmo-Jiongo et al. (2013), the tuning constants for the bias-correction we used are equal to 3.

The following quality measures, over Monte-Carlo simulations R , are used to evaluate the performance of an estimator of the average in area i , $\hat{\tau}_i$,

- The relative bias (RB)

$$\text{RB}(\hat{\tau}_i) = \frac{1}{R} \sum_{r=1}^R \frac{\hat{\tau}_i - \tau_i}{\tau_i}.$$

- The relative root mean squared error (RRMSE) for each area i :

$$\text{RRMSE}(\hat{\tau}_i) = \sqrt{\frac{1}{R} \sum_{r=1}^R \left(\frac{\hat{\tau}_i - \tau_i}{\tau_i} \right)^2}.$$

Table 1 reports the RB and the RRMSE of the small area estimators. It can be observed that all estimators are unbiased in the scenarios without outliers $(0, 0)$ and $(0, 0)_\rho$. Furthermore, the results confirm our expectations regarding the performance of the robust estimators. The robust projective estimators (REBLUP and SREBLUP) suffer from a negative bias in the case of non-symmetric outlier contamination (cf. for instance scenarios $[v, e]$ or $[v, e]_\rho$). Chambers et al. (2014) have already pointed out that this is mainly caused by the implicit assumption of robust projective estimators which assume that the outlier contamination component has zero expectation and that all outliers are observed in the sample. Our simulation scenarios have been specifically constructed to violate the assumptions of robust projective estimators. It is therefore natural that the robust projective estimators suffer from bias. On the other hand, the robust predictive estimators correct this bias. In particular, the robust (CCST and CHAM) and spatial robust predictive estimators (SCCST and SCHAM) have lower relative bias than the projective estimators (REBLUP and SREBLUP), especially in the areas 37-40. Furthermore, the CCST and SCCST estimators reveal slightly smaller relative bias compared to the CHAM and SCHAM estimators. In contrast, the CB and SCB estimators suffer from a relative large bias in the outlying areas 37-40 which is expected due to

Table 1: Model-based simulation results: performance of predictors of small area means

Median values of the RB in %										
Scenarios	$(0, 0)$	$(0, 0)_\rho$	$(v, 0)$	$(v, 0)_\rho$	$(v, 0)$	$(v, 0)_\rho$	(v, e)	$(v, e)_\rho$	(v, e)	$(v, e)_\rho$
Areas	1-40	1-40	1-36	1-36	37-40	37-40	1-36	1-36	37-40	37-40
Estimators										
EBLUP	0.00	0.00	0.09	0.09	-0.53	-0.53	0.12	0.11	-1.14	-1.13
SEBLUP	0.00	0.01	0.06	0.04	-0.44	-0.44	0.07	0.05	-1.02	-1.03
REBLUP	0.00	0.00	0.11	0.11	-0.47	-0.47	-0.08	-0.08	-0.77	-0.77
SREBLUP	0.00	0.00	0.07	0.05	-0.43	-0.45	-0.13	-0.13	-0.74	-0.77
CCST	0.00	-0.00	0.01	0.02	0.00	0.01	-0.11	-0.11	-0.18	-0.19
SCCST	0.00	0.00	0.02	0.02	0.01	0.01	-0.11	-0.11	-0.19	-0.19
CHAM	0.00	0.00	0.04	0.04	-0.03	-0.03	-0.06	-0.06	-0.28	-0.28
SCHAM	0.00	0.00	0.03	0.02	-0.03	-0.04	-0.08	-0.08	-0.27	-0.29
CB	0.00	0.00	0.09	0.09	-0.53	-0.53	0.03	0.03	-1.24	-1.23
SCB	0.00	0.01	0.06	0.04	-0.44	-0.45	-0.01	-0.03	-1.11	-1.12
Median values of the RRMSE in %										
Scenarios	$(0, 0)$	$(0, 0)_\rho$	$(v, 0)$	$(v, 0)_\rho$	$(v, 0)$	$(v, 0)_\rho$	(v, e)	$(v, e)_\rho$	(v, e)	$(v, e)_\rho$
Areas	1-40	1-40	1-36	1-36	37-40	37-40	1-36	1-36	37-40	37-40
Estimators										
EBLUP	0.81	0.82	0.86	0.86	1.01	1.01	1.19	1.18	1.70	1.69
SEBLUP	0.82	0.77	0.85	0.82	0.99	1.03	1.17	1.11	1.65	1.71
REBLUP	0.83	0.84	0.83	0.84	1.01	1.00	0.91	0.91	1.24	1.24
SREBLUP	0.84	0.79	0.84	0.79	0.98	1.04	0.92	0.87	1.21	1.27
CCST	0.92	0.92	0.93	0.93	0.86	0.86	1.12	1.12	1.02	1.02
SCCST	0.92	0.92	0.92	0.92	0.86	0.86	1.12	1.11	1.02	1.02
CHAM	0.86	0.86	0.90	0.91	0.86	0.86	1.02	1.02	1.02	1.02
SCHAM	0.87	0.83	0.90	0.89	0.85	0.85	1.02	1.00	1.02	1.02
CB	0.81	0.82	0.86	0.86	1.01	1.01	0.96	0.96	1.66	1.65
SCB	0.82	0.77	0.85	0.82	0.99	1.03	0.96	0.92	1.60	1.67

the contamination mechanism we use in the simulations. This bias is due to the non-robustified last terms in (8) and (18). Dongmo-Jiongo et al. (2013) present encouraging results for the CB estimator under a different data generation mechanism. Although the relative biases are small, we notice that SEBLUP and the SREBLUP estimators offer some bias reduction when compared respectively to the EBLUP and REBLUP estimators in the case of spatial scenarios.

We now turn to the RRMSE results in Table 1. As expected, in scenario $(0, 0)$ the robust predictive estimators CB and SCB are almost as efficient as the EBLUP, whereas the CHAM and SCHAM and the robust projective methods (REBLUP and SREBLUP) are somewhat less efficient.

In the case of the spatial scenario $(0, 0)_\rho$ the spatial methods have a lower RRMSE compared to the corresponding non-spatial methods, for instance, the RRMSE of SCB is 0.77% compared to 0.82% for the CB estimator. For scenario $(v, e)_\rho$ the bias-corrected predictive spatial methods (SCCST, SCHAM and SCB) offer some gains in RRMSE when compared to the non-spatial predictive estimators (CCST, CHAM and CB) in the areas 1-36. It appears that accounting for the spatial structure in the estimation leads to some efficiency gains.

The CB and SCB estimators are slightly more efficient than the CCST or CHAM based methods in the areas 1-36 at the cost of a higher RRMSE in areas 37-40, where area and individual outliers occur at the same time. The bias part of the MSE is not negligible for the CB and SCB estimators in these areas. The CCST and CHAM spatial and non-spatial methods indicate a more consistent performance with low RRMSEs over all areas in scenarios (v, e) and $(v, e)_\rho$. In contrast to the relative bias, the CHAM and SCHAM estimators perform slightly better in terms of RRMSE compared to the CCST and SCCST estimators. It seems that using information from other areas for the bias-correction in the CHAM methods helps to stabilise the results at the cost of a slightly higher bias compared to the CCST methods. These results illustrate the potentially beneficial effect of combining the spatial information with the bias correction offered by robust predictive estimators leading to more efficient results. However, robustness against outliers appears to be more important than incorporating the spatial structure in the estimation.

We now investigate the performance of the different MSE estimators. The objective of this part is twofold. First, we assess the performance of the proposed bootstrap methods for the spatial versions of the projective and predictive estimators. Second, we compare the performance of the robust and non-robust bootstrap methods in more detail in the case of spatial correlation. For both aims we focus on the scenarios with and without contamination in v and e .

MSE estimation is implemented with the parametric bootstrap methods presented in Section 4 with $B = 100$ bootstrap replicates. We denote by NonRobB and RobB the parametric bootstrap schemes where the v^* and e^* are generated by using non-robust or robust variance estimators respectively. We have also tested two selected scenarios, $(0, 0)_\rho$ and $(v, e)_\rho$, by using $B = 500$ bootstrap replicates and the results did not change considerably. MSE results for some estimators such as for the EBLUP (using the Prasad-Rao estimator and the bootstrap MSE estimator), the CB (using the bootstrap MSE) and the SCB (using the bootstrap MSE) are excluded but are available from the authors on request.

Starting with the first aim, Table 2 presents the results for the MSE estimators and shows the median values of their area-specific RB and RRMSE. We observe that both the robust and the non-robust spatial bootstrap methods work well for the scenarios without contamination. Under contamination, the bootstrap methods for the spatial estimators, SREBLUP, SCCST and SCHAM, lead to similar biases and stability compared to the bootstrap methods for the REBLUP, CCST and CHAM. Hence, the proposed MSE estimators that account for the additional spatial correlation in the data perform similarly to the bootstrap methods proposed by Sinha and Rao (2009) and Dongmo-Jiongo et al. (2013).

Having assessed the overall performance of the spatial versions of the bootstrap schemes, we now compare the proposed spatial robust and non-robust bootstrap methods explicitly. Figure 1 presents a more detailed picture of this comparison. It shows the ratio of the estimated non-robust MSE to the empirical (true) MSE (x-axis) against the ratio of the estimated robust MSE to the empirical (true) MSE (y-axis) for three estimators, the SREBLUP, the SCCST and the SCHAM. The crosses in the plot indicate the outlying areas 37-40. The vertical grey line indicates the sector where the empirical MSE corresponds to the estimated non-robust MSE. Areas left to the vertical grey line suffer from underestimation, whereas areas right to the vertical line indicate overestimation. The same principle applies to the

Table 2: Performance of MSE estimators in model-based simulations

Median values of the RB in %							
		(0, 0)	$(0, 0)_\rho$	(v, e)	$(v, e)_\rho$	(v, e)	$(v, e)_\rho$
Areas		1-40	1-40	1-36	1-36	37-40	37-40
Estimators	MSE						
REBLUP	RobB	-2.97	-2.64	1.28	1.75	-31.04	-30.66
	NonRobB	-1.27	-1.56	39.91	41.02	-5.65	-5.57
SREBLUP	RobB	-2.82	-5.76	-0.90	0.93	-30.25	-34.84
	NonRobB	-3.51	-5.86	34.07	37.48	-6.03	-10.24
CCST	RobB	0.75	1.02	-11.18	-11.32	-10.30	-10.57
	NonRobB	0.38	0.19	22.37	22.74	23.29	23.38
SCCST	RobB	0.98	0.87	-10.29	-9.86	-10.38	-10.41
	NonRobB	-0.66	-0.90	21.83	22.20	23.87	23.43
CHAM	RobB	-1.46	-1.44	-7.47	-7.08	-13.70	-13.63
	NonRobB	-0.60	-0.91	27.30	27.68	17.82	18.07
SCHAM	RobB	-1.73	-4.01	-7.16	-8.69	-14.33	-16.55
	NonRobB	-2.76	-4.61	24.66	24.68	16.25	13.89
Median values of the RRMSE in %							
		(0, 0)	$(0, 0)_\rho$	(v, e)	$(v, e)_\rho$	(v, e)	$(v, e)_\rho$
Areas		1-40	1-40	1-36	1-36	37-40	37-40
Estimators	MSE						
REBLUP	RobB	11.12	9.90	10.70	10.61	32.32	31.98
	NonRobB	9.73	9.26	44.38	45.33	17.12	17.09
SREBLUP	RobB	11.69	12.33	10.80	11.84	31.46	35.92
	NonRobB	10.55	11.55	38.94	42.37	17.63	19.46
CCST	RobB	10.26	9.95	14.46	15.09	14.21	14.54
	NonRobB	9.30	9.28	30.63	31.01	31.26	31.33
SCCST	RobB	10.42	10.27	14.47	14.24	14.03	14.31
	NonRobB	9.26	9.33	30.06	30.55	31.40	31.00
CHAM	RobB	10.63	9.82	12.22	11.99	16.73	16.83
	NonRobB	9.64	9.06	33.55	33.59	25.79	25.96
SCHAM	RobB	10.84	10.72	12.14	13.54	17.10	19.16
	NonRobB	9.74	10.62	31.08	31.24	24.61	23.14

horizontal grey line for the estimated robust MSE (underestimation below the grey line, overestimation above). The diagonal represents the sector where the estimated robust and non-robust MSE lead to the same results. Figure 1 reveals that the robust and the non-robust bootstrap schemes perform similarly in scenario $(0, 0)_\rho$. As a first comment in the scenario $(v, e)_\rho$, we note that the non-robust bootstrap method suffers from overestimation when estimating the MSE of the predictive estimators (SCHAM and SCCST). Second, the non-robust bootstrap method for the robust projective estimator (SREBLUP) overestimates the MSE for most areas, but not for the outlying areas 37-40 which is indicated by the crosses. In contrast, the robust version of the bootstrap MSE for the SREBLUP estimator is more stable, especially in the areas with only individual contamination (cf. areas 1-36), but underestimates the MSE for the outlying areas, because of a negative bias shown in Table 2. Third, the robust bootstrap MSE estimator performs consistently better than the non-robust version when used to estimate the MSE of the predictive estimators (SCCST and SCHAM). These results suggest that using the robust version of




Figure 1: Robust and non-robust bootstrap estimators under spatial correlation.

the bootstrap for estimating the MSE of the predictive estimators appears to have appealing properties regarding the stability and bias in these scenarios.

6 Application: Small area estimation of average labour costs for small and medium-sized enterprises in Italy

In Section 5 we evaluated the performance of different small area estimators under known contamination mechanisms. When using real data, however, the outlier contamination mechanism is unknown. In this section we employ the proposed methodology for estimating average labour costs for provinces in Italy. The focus here is on small and medium-sized enterprises with less than 100 employees as this group of firms is of great interest for economic policy (cf. Acs and Audretsch, 1988 or Sawyerr et al., 2003). In addition, for practical reasons, we only focus on one industry domain NACE 25 which refers to enterprises producing rubber and plastic products.

The business survey data of interest for this application come from Small and Medium Enterprises (SME) survey carried out annually by the Italian National Statistical Institute (ISTAT). In addition to the survey data, small area estimation is facilitated by the use of auxiliary information from the Italian Statistical Business Register (Asia - Archivio Statistico delle Imprese Attive). However, the use of business survey small area data is governed by strict confidentiality rules. To overcome this problem, in the application we use a synthetic dataset (TRItalia), which was created by using real data from the SME survey and the Asia register and preserves the structure of the original data (cf. Kolb et al., 2013). Similar to the Asia register, the TRItalia dataset consists of around 12,000 enterprises in the NACE 25 industry domain. Using the TRItalia dataset we selected a sample of 795 enterprises following a sampling design similar to the one of the SME survey. The design is a single stage stratified random sampling with strata defined by the cross-classification of administrative areas (NUTS 1) and a categorical variable for

Table 3: Summary statistics over provinces.

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
Sample size	3.00	5.00	6.00	7.72	9.00	39.00
LCO (in KEUR)	0.01	9.67	50.68	216.10	176.60	15800.00
EMT	1.00	2.00	6.00	10.82	13.00	96.00

company size leading to 20 strata. In this application we are interested in providing reliable small area estimates of the average labour costs for enterprises in the NACE 25 sector for 103 Italian provinces. Similar to Fabrizi et al. (2013), the outcome of interest is the labour costs (LCO) in thousand Euro (KEUR).

Table 3 presents the distribution of the sample sizes, labour costs and number of employees (EMT) over provinces. The asymmetry in the unconditional distribution of labour costs is evident. It is more appropriate, though, to analyse the conditional distribution of labour costs given a set of auxiliary variables available.

To do so, we use a 2-level random effects model (level 1 = enterprise; level 2 = province) for labour costs with random effects specified at the level of provinces. The auxiliary variables used in this model consist of the number of employees. To control for the stratified design of our sample, we further included the stratification variable defined by the cross classification between administrative areas and a categorical variable for company size. This defines 20 strata. This model provides a starting point which can later be extended. To assess departures from the model assumptions, Figure 2 presents normal probability plots of level 1 and level 2 residuals. The figure indicates that the Gaussian assumptions of the




Figure 2: Normal probability plots of level 1 (left) and level 2 (right) residuals.

model are violated. Furthermore, the Shapiro-Wilk test, which rejects the null hypothesis of normal distributed residuals (p-values: level 1 = $2.2e-16$ and level 2 = $2.2e-16$) confirms this. Using a robust estimation method to control for the influence of outliers may be advisable in this case.

Evidence of moderate spatial correlation in the data is also present. Following Pratesi and Salvati (2008), we compute two diagnostics for the presence of spatial autocorrelation. Moran's I (cf. Moran,

Table 4: Values of the Moran’s I and Geary’s C tests.

	Value	p value
Moran’s I	0.443	$< 1.9e^{-11}$
Geary’s C	0.509	$< 1.3e^{-11}$

Table 5: Distribution of average labour costs (in KEUR) over provinces in Italy.

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
REBLUP	10.39	53.57	85.48	94.19	125.20	311.90
SREBLUP	10.43	54.02	85.05	93.87	125.90	313.10
CHAM	9.81	57.49	100.60	102.40	136.30	323.60
SCHAM	9.71	60.42	98.80	102.00	134.50	329.60
CCST	11.06	54.84	83.00	98.43	124.30	348.20
SCCST	11.06	54.01	86.52	98.20	124.00	350.10

1950) is analogous to the correlation coefficient with values between -1 (strong negative spatial correlation) and 1 (strong positive spatial correlation), whereas 0 indicates no spatial correlation. Geary’s C (cf. Cliff and Ord, 1981) ranges between 0 (strong positive spatial correlation) and 2 (strong negative spatial correlation). Both tests, shown in Table 4, indicate the presence of positive spatial correlation in the data.

Estimates of average labour costs for each province are calculated by using the robust projective estimators REBLUP (3) and SREBLUP (14). We also computed estimates using the robust predictive estimators CCST (4) and CHAM (6), as well as their corresponding spatial extensions SCCST (15) and SCHAM (17). In order to save space, we omit the results for the standard EBLUP (2) and spatial EBLUP (12). We further exclude the CB (8) and SCB (18). These results are available from the authors on request.

Table 5 presents the distribution of the estimated average labour costs in the sector of interest (NACE 25) for provinces in Italy. Our first observation is that there are some differences between the estimates obtained from the robust projective estimators (REBLUP and SREBLUP) and those from the robust predictive estimators (CCST, CHAM, SCCST and SCHAM). In order to further investigate the reason for these differences we performed the following sensitivity analysis. Using the Cook’s distance measure, we removed the most influential outliers from the sample. In total we removed 56 out of 795 observations. Using this reduced dataset, we recomputed the small area estimates. Figure 3 shows the robust projective estimator (REBLUP) (x-axis) against the robust predictive estimator (CHAM) (y-axis) for the reduced and original samples. The corresponding plot for the SREBLUP and SCHAM estimators is presented in Figure 4. We observe that the point estimates of the projective and predictive estimators are closer in the reduced sample. This behaviour can be explained by the fact that the restricted dataset includes a smaller number of outliers, and hence, the robust projective and predictive estimators provide closer results. In contrast, the stronger contamination in the original sample leads to systematically smaller estimates of the REBLUP and SREBLUP compared to the robust predictive estimators (CHAM and SCHAM). The

same picture holds also for the comparison between the projective estimators (REBLUP and SREBLUP) and the predictive estimators (CCST and SCCST). The results from this sensitivity analysis provide evidence that using a predictive estimator is possibly appropriate in this case.




Figure 3: Differences between the projective estimator (REBLUP) and the predictive estimator (CHAM) for the reduced and original samples.




Figure 4: Differences between the projective estimator (SREBLUP) and the predictive estimator (SCHAM) for the reduced and original samples.

MSE estimates, using the two bootstrap schemes we described in Section 4, are presented in Table 6. Based on the findings from the model-based simulations, in the presence of contamination we may expect overestimation of the MSE when bootstrapping from the non-robust estimates of the variance components. Indeed, the results suggest that this MSE estimator provides estimates that are very different to the ones obtained from the bootstrap method that uses the robust variance estimators. Examining further the MSE estimates, we note that the SREBLUP MSE estimates are lower than the corresponding REBLUP estimates. This suggests some efficiency gains obtained by incorporating the spatial information in estimation. The same holds true when comparing the CHAM to the SCHAM and the CCST to the SCCST estimators. Considering the point and MSE estimates, we can conclude that the SCHAM or SCCST estimators appear to be a good choice for producing small area estimates of average labour

costs. The conservative analyst would choose the SCCST estimator, because of the slightly lower bias at the cost of a higher variance compared to the SCHAM estimator. In contrast, the SCHAM estimator offers more stability at the expense of a higher bias in relation to the SCCST estimator.

Table 6: MSE estimation						
MSE estimates using robust parametric bootstrap						
	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
REBLUP	19.12	44.10	51.38	54.74	63.80	105.40
SREBLUP	17.76	41.97	48.22	51.10	58.84	89.46
CHAM	20.38	49.84	56.33	59.75	69.27	113.10
SCHAM	18.36	45.29	54.09	55.65	64.90	97.99
CCST	25.95	117.10	168.90	185.70	243.20	447.90
SCCST	26.51	113.50	171.20	183.10	241.30	398.80
MSE estimates using non-robust parametric bootstrap						
	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
REBLUP	5191	11890	14060	14240	15870	26980
SREBLUP	4843	9927	11110	11780	13650	26000
CHAM	5693	13210	15640	15680	17780	28300
SCHAM	4899	11310	12190	12760	14560	28970
CCST	6748	27670	38160	39780	50800	93640
SCCST	6278	25070	35510	37980	46460	83500

In Table 7 we investigate the estimates obtained from the SCHAM estimator in detail (SCCST estimates are available from the authors upon request). This table shows the mean labour costs per province divided by the average number of employees obtained from the synthetic register. An average employee in a small and medium-sized enterprise earns in Italy around 16,440 EUR per year. This is consistent with official results provided by ISTAT (cf. ISTAT, 2007). A more detailed picture of the spatial distribution of labour costs can be found in Figure 5. The figure shows the labour costs per employee for the 103 Italian provinces. As expected, we detect large regional disparities. The north-south divide in labour costs in Italy is clearly depicted in this figure.

7 Concluding remarks

The paper proposes extensions to the currently available outlier robust projective and robust predictive small area estimators in the presence of spatial correlation. The gains offered by the proposed methodology are twofold. First, the outlier robust nature of the estimators reduces the impact of the misspecification of the model assumptions. Second, the methodology allows for modelling the spatial structure in the data. MSE estimation is performed by using two parametric bootstrap schemes that incorporate the

Table 7: Labour costs per employee per province (in KEUR) using the SCHAM estimator.

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
LCO per employee	5.26	13.94	16.76	16.44	19.13	42.35




Figure 5: Mean labour costs per employee per province (in KEUR) using the SCHAM estimator.

spatial dimension. The results from the model-based simulations show that in the presence of contamination bootstrapping the data should be done as in Sinha and Rao (2009) by using the robust estimates of the variance components.

Currently, estimation relies on the a-priori choice of the tuning constants to be used in the influence functions. One could develop an adaptive (data driven) tuning constant to be used for small area prediction. This provides one avenue for future research. Moreover, the spatial small area estimators depend on the structure of a given contiguity matrix \mathbf{W} . Another line for further work could be to investigate the effect of errors in specifying \mathbf{W} on the estimation. Furthermore, the use of a bootstrap MSE estimator is computationally very demanding. Although the implementation of the proposed methodology is facilitated by the availability of a computationally efficient algorithm using C++ in R, its application in practice is challenging. Developing analytic MSE estimators similar to the one proposed in Chambers et al. (2014) offers an additional avenue for future research.

Acknowledgements

The research was conducted within the BLUE-ETS research project which is funded by the European Commission within the 7th Framework Programme. For more information on the project, we refer to the project page <http://www.blue-ets.eu>. The authors are grateful to ISTAT and the BLUE-ETS project coordinator for kindly providing the business data. In addition, the research has received support from the InGRID (Inclusive Growth Research Infrastructure Diffusion) grant within the 7th Framework Programme. Furthermore, we would like to thank Dongmo-Jiongo et al. (2013) for providing R-code used in the simulations in this paper.

Appendix

In the appendix we show how the SEBLUP in equation (16) can be reformulated.

$$\begin{aligned}
N_i \hat{y}_i^{SEBLUP} &= \sum_{j \in s} w_j^{sp} y_j \\
&= \sum_{h=1}^m \sum_{j \in s_h} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp} + \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} + \hat{v}_h^{\psi, sp} \right) \\
&= \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \hat{v}_h^{\psi, sp} + \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp} \right) + \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} \\
&\quad + \sum_{j \in s_i} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp} \right) + \sum_{j \in s_i} w_j^{sp} \left(\mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} + \hat{v}_i^{\psi, sp} \right) \\
&= \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \hat{v}_h^{\psi, sp} + \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp} \right) + \sum_{j \in s} w_j^{sp} \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} \\
&\quad + \sum_{j \in s_i} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp} \right) + \sum_{j \in s_i} w_j^{sp} \hat{v}_i^{\psi, sp} \\
&= \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \hat{v}_h^{\psi, sp} + \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp} \right) + \sum_{j \in s_i} w_j^{sp} \hat{v}_i^{\psi, sp} \\
&\quad + \sum_{j \in U_i} \left(\mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} + \hat{v}_i^{\psi, sp} \right) + \sum_{j \in s_i} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp} \right) - N_i \hat{v}_i^{\psi, sp} \\
&= \sum_{j \in s_i} y_j + \sum_{j \in r_i} \left(\mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} + \hat{v}_i^{\psi, sp} \right) + \sum_{j \in s_i} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp} \right) \\
&\quad - \sum_{j \in s_i} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp} \right) + \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp} \right) \\
&\quad + \sum_{j \in s_i} w_j^{sp} \hat{v}_i^{\psi, sp} - N_i \hat{v}_i^{\psi, sp} + \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \hat{v}_h^{\psi, sp} \\
&= \sum_{j \in s_i} y_j + \sum_{j \in r_i} \left(\mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} + \hat{v}_i^{\psi, sp} \right) + \sum_{j \in s_i} (w_j^{sp} - 1) \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp} \right) \\
&\quad + \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp} \right) + \left(\sum_{j \in s_i} w_j^{sp} - N_i \right) \hat{v}_i^{\psi, sp} + \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \hat{v}_h^{\psi, sp} \\
&= N_i \hat{y}_i^{SREBLUP} + \sum_{j \in s_i} (w_j^{sp} - 1) \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_i^{\psi, sp} \right) \\
&\quad + \sum_{h=1}^m \sum_{\substack{j \in s_h \\ h \neq i}} w_j^{sp} \left(y_j - \mathbf{x}_j^T \hat{\boldsymbol{\beta}}^{\psi, sp} - \hat{v}_h^{\psi, sp} \right) + \sum_{h=1}^m w_h^{sp} \hat{v}_h^{\psi, sp}
\end{aligned}$$

References

- Acs, Z. and D. Audretsch (1988). Innovation in large and small firms: An empirical analysis. *American Economic Review* 78 (4), 678–690.
- Battese, G. E., R. M. Harter, and W. A. Fuller (1988). An error component model for prediction of

- county crop areas using survey and satellite data. *Journal of the American Statistical Association* 83 (401), 28–36.
- Beaumont, J.-F., D. Haziza, and A. Ruiz-Gazen (2013). A unified approach to robust estimation in finite population sampling. *Biometrika* 100 (3), 555–569.
- Bivand, R., E. Pebesma, and V. Gomez-Rubio (2008). *Applied Spatial Data Analysis with R*. New York: Springer.
- Chambers, R. (1986). Outlier robust finite population estimation. *Journal of the American Statistical Association* 81, 1063–1069.
- Chambers, R., H. Chandra, N. Salvati, and N. Tzavidis (2014). Outlier robust small area estimation. *Journal of the Royal Statistical Society: Series B* 76 (1), 47–69.
- Chambers, R. and N. Tzavidis (2006). M-quantile models for small area estimation. *Biometrika* 93 (2), 255–268.
- Chandra, H., N. Salvati, R. Chambers, and N. Tzavidis (2012). Small area estimation under spatial nonstationarity. *Computational Statistics and Data Analysis* 56, 2875–2888.
- Cliff, A. and J. Ord (1981). *Spatial processes*. London: Pion.
- Dongmo-Jiongo, V., D. Haziza, and P. Duchesne (2013). Controlling the bias of robust small area estimators. *Biometrika* 100, 843–858.
- Fabrizi, E., M. Ferrante, and C. Trivisano (2013). Small area estimation of labor productivity for the Italian manufacturing SME cross-classified by region, industry and size. Ersra conference papers, European Regional Science Association.
- Hall, P. and T. Maiti (2006). On parametric bootstrap methods for small area prediction. *Journal of the Royal Statistical Society: Series B* 68 (2), 221–238.
- Henderson, C. R. (1950). Estimation of genetic parameters. *Annals of Mathematical Statistics* 21 (2), 309–310.
- ISTAT (2007). Conti economici delle imprese: Anno 2003. informazioni n. 8. Downloaded from http://www3.istat.it/dati/catalogo/20070827_01/inf.07_08.Conti_economici_delle_imprese.2003.pdf (7th May 2014).
- Kolb, J.-P., R. Münnich, F. Volk, and T. Zimmermann (2013). *Best practice recommendations on variance estimation and small area estimation in business surveys (final report)*, Chapter 5. TRItalia dataset, pp. 168–188. BLUE-ETS Project, FP7-SSH-2009-A, Deliverable 6.2.
- Moran, P. (1950). Notes on continuous stochastic phenomena. *Biometrika* 37, 17–33.

- Muñoz-Pichardo, J., J. Muñoz-García, J. Moreno-Rebollo, and R. Pino-Mejías (1995). A new approach to influence analysis in linear models. *Sankhya: The Indian Journal of Statistics, Series A (1961-2002)* 57 (3), 393–409.
- Petrucci, A., M. Pratesi, and N. Salvati (2005). Geographic information in small area estimation: Small area models and spatially correlated random area effects. *Statistics in Transition* 7 (3), 609–623.
- Pratesi, M. and N. Salvati (2008). Small area estimation: the eblup estimator based on spatially correlated random area effects. *Statistical Methods & Applications* 17, 113–141.
- Royall, R. (1976). The linear least-squares prediction approach to two-stage sampling. *Journal of the American Statistical Association* 71, 657–664.
- Salvati, N., N. Tzavidis, M. Pratesi, and R. Chambers (2012). Small area estimation via m-quantile geographically weighted regression. *Test* 21 (1), 1–28.
- Sawyer, O., J. McGee, and M. Peterson (2003). Perceived uncertainty and firm performance in small and medium enterprises. *International Small Business Journal* 21 (3), 269–290.
- Schmid, T. and R. Münnich (2014). Spatial robust small area estimation. *Statistical Papers* 55, 653–670.
- Searle, S. R. (1971). *Linear Models*. New York: Wiley.
- Sinha, S. K. and J. N. K. Rao (2009). Robust small area estimation. *The Canadian Journal of Statistics* 37 (3), 381–399.
- Tzavidis, N., S. Marchetti, and R. Chambers (2010). Robust estimation of small area means and quantiles. *Australian and New Zealand Journal of Statistics* 52 (2), 167–186.
- Welsh, A. and E. Ronchetti (1998). Bias-calibrated estimation from sample surveys containing outliers. *Journal of the Royal Statistical Society: Series B* 60, 413–428.

Diskussionsbeiträge - Fachbereich Wirtschaftswissenschaft - Freie Universität Berlin
Discussion Paper - School of Business and Economics - Freie Universität Berlin

2015 erschienen:

- 2015/1 GÖRLITZ, Katja und Christina GRAVERT
The effects of increasing the standards of the high school curriculum on school dropout
Economics
- 2015/2 BÖNKE, Timm und Clive WERDT
Charitable giving and its persistent and transitory reactions to changes in tax incentives: evidence from the German Taxpayer Panel
Economics
- 2015/3 WERDT, Clive
What drives tax refund maximization from inter-temporal loss usage? Evidence from the German Taxpayer Panel
Economics
- 2015/4 FOSSEN, Frank M. und Johannes KÖNIG
Public health insurance and entry into self-employment
Economics
- 2015/5 WERDT, Clive
The elasticity of taxable income for Germany and its sensitivity to the appropriate model
Economics
- 2015/6 NIKODINOSKA, Dragana und Carsten SCHRÖDER
On the Emissions-Inequality Trade-off in Energy Taxation: Evidence on the German Car Fuel Tax
Economics
- 2015/7 GROß, Marcus; Ulrich RENDTEL, Timo SCHMID, Sebastian SCHMON und Nikos TZAVIDIS
Estimating the density of ethnic minorities and aged people in Berlin: Multivariate kernel density estimation applied to sensitive geo-referenced administrative data protected via measurement error
Economics