

A Service of



Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Feizollahi, Mohammad Javad; Feyzollahi, Hadi

# Article Robust quadratic assignment problem with budgeted uncertain flows

**Operations Research Perspectives** 

**Provided in Cooperation with:** Elsevier

*Suggested Citation:* Feizollahi, Mohammad Javad; Feyzollahi, Hadi (2015) : Robust quadratic assignment problem with budgeted uncertain flows, Operations Research Perspectives, ISSN 2214-7160, Elsevier, Amsterdam, Vol. 2, pp. 114-123, https://doi.org/10.1016/j.orp.2015.06.001

This Version is available at: https://hdl.handle.net/10419/178255

# Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.



http://creativecommons.org/licenses/by/4.0/

# Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



# WWW.ECONSTOR.EU

#### Operations Research Perspectives 2 (2015) 114-123

Contents lists available at ScienceDirect

# **Operations Research Perspectives**

journal homepage: www.elsevier.com/locate/orp

# Robust quadratic assignment problem with budgeted uncertain flows

# Mohammad Javad Feizollahi<sup>a,\*</sup>, Hadi Feyzollahi<sup>b</sup>

<sup>a</sup> H. Milton Stewart School of Industrial & Systems Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA
<sup>b</sup> Department of Industrial and Systems Engineering, University at Buffalo, Buffalo, NY 14260, USA

#### HIGHLIGHTS

- A robust approach for quadratic assignment problem (RQAP) with budgeted uncertainty.
- An exact and two heuristic methods to solve RQAP.
- Extensive experiments to show performance of methods and quality of solutions.
- RQAP can be solved significantly faster than minmax regret QAP.
- RQAP has adjustable conservativeness while minmax regret QAP has not.

# ARTICLE INFO

Article history: Available online 9 June 2015

Keywords: Robust optimization Budgeted uncertainty Quadratic assignment problem 2-Opt Tabu search

# ABSTRACT

We consider a generalization of the classical quadratic assignment problem, where material flows between facilities are uncertain, and belong to a budgeted uncertainty set. The objective is to find a robust solution under all possible scenarios in the given uncertainty set. We present an exact quadratic formulation as a robust counterpart and develop an equivalent mixed integer programming model for it. To solve the proposed model for large-scale instances, we also develop two different heuristics based on 2-Opt local search and tabu search algorithms. We discuss performance of these methods and the quality of robust solutions through extensive computational experiments.

> © 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

# 1. Introduction

[1] introduced the standard quadratic assignment problem (QAP). Standard QAP deals with choosing an optimal way to assign n facilities to n locations to minimize the total material handling cost, given all distances between locations and the amount of material flow between each pair of facilities. A more general form of the QAP was proposed by [2]. [3,4] considered multi-dimensional QAP.

QAP is one of the hardest problems in combinatorial optimization [5,6] and even finding a constant-factor approximate solution for the QAP is NP-hard [7]. However, some specific cases of QAP are easy to solve [8,9]. Many exact and heuristic methods have been developed to solve different cases of QAP. Approximated dynamic programming [10], genetic algorithm [11], parallel algorithms [12,13], hybrid algorithms [14], teaching learning based optimization [15], semidefinite programming relaxations [16,17], mixed integer linear programming reformulation [18–22], reformulation linearization technique (RLT) [23–25], formulation reductions [26], and exploiting data structure [27] are some of these techniques.

QAP has numerous applications such as backboard wiring [28], scheduling problems [29], economic problems [30], designing typewriter keyboards [31], facility layout [32–34], assembling printed circuit boards [35] and many other applications. For a detailed discussion about applications and solution methods for QAP see [36,6,37,38].

In deterministic optimization, it is assumed that input data (e.g. flows between facilities and distances between locations in QAP) are precisely known in advance. Although this assumption can be true in some applications, it is not realistic in many others [39]. [40] proposed a design for robust facility layout under the dynamic demand environment. In their approach, the layout of expected flow or expected demand is applied in all the periods. [41] developed a fuzzy model to address uncertainty in QAP. [42] reviewed facility location problems under uncertainty. [43] studied integration of facility layout design and flow assignment problem under demand uncertainty. [44] considered uncertainty in a hospital layout problem and proposed a robust model for this problem.





<sup>\*</sup> Corresponding author. Tel.: +1 404 425 0885.

*E-mail addresses*: feizollahi@gatech.edu (M.J. Feizollahi), hadifeyz@buffalo.edu (H. Feyzollahi).

QAP with uncertain locations was studied in [39]. [45] used a robust deviation (minmax regret) approach to deal with uncertainty in material flows.

In uncertain optimization problems with discrete variables, in addition to robust deviation, we can use budgeted uncertainty which has the same complexity as the original model and adjustable conservativeness [46,47]. In practice, for an uncertain mixed integer programming (MIP) problem with interval data, solving robust counterparts for budgeted uncertainty sets is much easier than finding the minmax regret solution. For example, in redundancy allocation problems, this difference is obvious by comparing the results in [48–51], respectively. In addition, former method can find solutions with different levels of conservative ness, while the latter approach outputs only one conservative solution.

In this paper, we consider a generalization of the QAP where the flows are uncertain for some subset *J* of pairs of facilities. For the flow between each pair of facilities only an interval estimate (uncertainty interval) is available, and the flow can take on any value from the corresponding uncertainty interval. But, for a given protection level  $\Gamma \in [0, |J|]$ , it is assumed that at most  $\lfloor \Gamma \rfloor$  of uncertain flows are allowed to change, and one flow changes by a ratio of at most  $(\Gamma - \lfloor \Gamma \rfloor)$  of its uncertainty interval. We are interested in an assignment which minimizes the maximum cost for any possible realization of flows. In other words, because it is unlikely that all uncertain flows adversely affect the cost of assignment, it is assumed that only a subset of uncertain flows change from their nominal values.

This paper is organized as follows. In Section 2, we present notation and problem statement for deterministic and uncertain QAP, and an efficient MIP equivalent for QAP. In Section 3, we develop a mathematical programming formulation of the problem as well as an equivalent MIP model. Then, two heuristic algorithms are described in Section 4. Experimental results are discussed in Section 5, and conclusions are presented in Section 6.

## 2. Notation and problem statement

In this section, we first present notation and problem statement for classical QAP which is mostly quoted from [45] with some slight adjustments. Then, we present an efficient MIP equivalent for QAP from the literature. Finally, we introduce budgeted uncertainty in flow between facilities and describe some concepts related to the proposed uncertain QAP.

#### 2.1. Classical QAP

In the standard version of QAP, it is assumed that there are *n* facilities that should be assigned to *n* locations, in order to minimize the total material handling cost [1]. Let  $N = \{1, 2, ..., n\}$ . For each pair  $i, j \in N$  of facilities, let  $f_{ij} \ge 0$  be the flow from facility *i* to facility *j*. In addition, for each pair  $k, l \in N$  of locations, let  $d_{kl} \ge 0$  be the travel distance from location *k* to location *l*. An assignment of facilities to locations can be represented by an  $n \times n$  binary matrix *X*, where

 $x_{ik} = \begin{cases} 1 & \text{if facility } i \text{ is assigned to location } k, \\ 0 & \text{otherwise.} \end{cases}$ 

In any feasible assignment  $X \in \{0, 1\}^{n \times n}$ , each location must be assigned exactly to one facility, and similarly each facility must be located exactly in one location. Therefore, the set *P* of all possible assignments is defined by constraints

$$\sum_{k=1}^{n} x_{ik} = 1 \quad \forall i \in N, \tag{1}$$

$$\sum_{i=1}^{n} x_{ik} = 1 \quad \forall k \in N,$$
(2)

$$x_{ik} \in \{0, 1\} \quad i, k \in \mathbb{N}.$$
 (3)

For any  $X \in P$ , let  $\phi_i^X$  denote the location assigned to facility i in assignment X. Let  $d_{ij}^X$  be the distance between facilities i and j in assignment X. Therefore,

$$d_{ij}^{X} := d_{\phi_{i}^{X}\phi_{j}^{X}} = \sum_{k=1}^{n} \sum_{l=1}^{n} d_{kl} x_{ik} x_{jl}.$$
(4)

Given an  $n \times n$  flow matrix  $f = (f_{ij})$  and an assignment  $X = (x_{ik}) \in P$ , let  $\langle f, X \rangle$  denote the corresponding cost of the assignment,

$$\langle f, X \rangle := \sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{k=1}^{n} \sum_{l=1}^{n} f_{ij} d_{kl} x_{ik} x_{jl} = \sum_{i=1}^{n} \sum_{j=1}^{n} f_{ij} d_{ij}^{X}.$$
 (5)

For a given flow matrix *f*, the *classical* QAP is:

**QAP**(
$$f$$
): Minimize { $\langle f, X \rangle | X \in P$ }. (6)

#### 2.2. Xia-Yuan linearization

[2] proposed a more general form of the QAP as follows:

$$\min_{X \in P} \sum_{i=1}^{n} \sum_{j=1}^{n} \sum_{k=1}^{n} \sum_{l=1}^{n} c_{ijkl} x_{ik} x_{jl}$$
(7)

where  $c_{ijkl} \ge 0, i, j, k, l \in N$  are given coefficients. Note that **QAP**(f) is a special case of (7) where  $c_{ijkl} = f_{ij}d_{kl}$ , for all  $i, j, k, l \in N$ . As discussed in [6], different approaches have been developed to linearize the general QAP (7). [22] demonstrated experimentally that [19] linearization is quite effective as a MIP formulation for the general QAP. The Xia-Yuan linearization for general QAP (7) is:

$$\min_{X \in P} \sum_{i=1}^{n} \sum_{k=1}^{n} (w_{ik} + c_{iikk} x_{ik}),$$
(8)

$$w_{ik} \ge \sum_{\substack{j=1\\ i \ne k}}^{n} \sum_{\substack{l=1\\ i \ne k}}^{n} c_{ijkl} x_{jl} - \hat{a}_{ik} (1 - x_{ik}), \quad \forall i, k \in \mathbb{N},$$
(9)

$$w_{ik} \ge l_{ik} x_{ik}, \quad \forall i, k \in N,$$
(10)

where

$$l_{ik} = \min_{\substack{X \in P \\ x_{ik}=1}} \sum_{\substack{j=1 \\ j \neq i}}^{n} \sum_{\substack{l=1 \\ l \neq k}}^{n} c_{ijkl} x_{jl}, \text{ and}$$

$$\hat{a}_{ik} = \max_{\substack{X \in P \\ x_{ik}=1}} \sum_{\substack{j=1 \\ j \neq i}}^{n} \sum_{\substack{l=1 \\ l \neq k}}^{n} c_{ijkl} x_{jl}.$$
(11)

Observe that constants  $l_{ik}$ ,  $\hat{a}_{ik}$  are obtained by means of solving the regular linear assignment problems (11) which can be done in polynomial time [6]. Values  $l_{ik}$  are called Gilmore–Lawler constants [6]. Formulation (8)–(10) has  $n^2$  binary variables,  $n^2$  continuous variables, and  $2n^2 + 2n$  linear constraints.

## 2.3. QAP with budgeted uncertainty

**QAP**(f) is a valid optimization problem as long as the values of flows and distances are known precisely. However, flows between facilities are typically only estimated within most likely intervals. In the remainder of the paper, we deal with uncertain flows.

Suppose that for any  $(i, j) \in N \times N$ , two numbers  $f_{ij}^-, f_{ij}^+$  are given,  $0 \le f_{ij}^- \le f_{ij}^+$ , where  $f_{ij}^-$  and  $f_{ij}^+$  are lower and upper bounds

5)

for  $f_{ij}$ , respectively. Let  $J \subset N \times N$  be the set of pairs (i, j) of facilities with uncertain flow; that is,  $f_{ij}^- < f_{ij}^+$  for all  $(i, j) \in J$ , and  $f_{ij}^- = f_{ij}^+$ for all  $(i, j) \in J^c = (N \times N) \setminus J$ . Assume that the flow  $f_{ij}$  can take any value on  $[f_{ij}^-, f_{ij}^+]$ . To normalize the uncertainty set, suppose that  $f_{ij} = f_{ij}^- + \delta_{ij}\xi_{ij}$ , where  $\delta_{ij} := f_{ij}^+ - f_{ij}^-$  is the interval length and  $\xi_{ij}$  is an uncertain variable in [0, 1]. Let  $f^-, f^+$  and  $f^=$  be the scenarios (flow matrices) where for all  $i, j \in N$ ,  $f_{ij}$  equals to  $f_{ij}^-$ ,  $f_{ij}^+$  and  $0.5 \times (f_{ij}^- + f_{ij}^+)$ , respectively. For a given protection level  $\Gamma \in [0, |J|]$ , the set

$$U_{\Gamma} := \left\{ \xi \in \mathfrak{N}^{|\mathcal{I}|} : \sum_{(i,j) \in J} \xi_{ij} \le \Gamma, \ 0 \le \xi_{ij} \le 1, \ \forall (i,j) \in J \right\}$$
(12)

is called a budgeted uncertainty set. [46] introduced budgeted uncertainty to construct robust counterparts for LP and MIP models with adjustable conservativeness. While  $\Gamma = 0$  gives the most optimistic solution (deterministic problem with  $f = f^-$ ),  $\Gamma = |J|$  generates the most pessimistic one (deterministic problem with  $f = f^+$ ).

For any perturbation vector  $\xi$ , denote the corresponding flow matrix by  $f(\xi)$ . Then, the uncertainty set for flow matrix f is:

$$V_{\Gamma} := \left\{ f \in \mathfrak{R}^{n \times n} : \exists \xi \in U_{\Gamma} \text{ such that } f = f(\xi) \right\}$$
$$= \left\{ f \in \mathfrak{R}^{n \times n} : \sum_{(i,j) \in J} \frac{f_{ij} - f_{ij}^{-}}{\delta_{ij}} \leq \Gamma, \ f_{ij}^{-} \leq f_{ij} \leq f_{ij}^{+}, \right.$$
$$\forall (i,j) \in N \times N \left\} .$$
(13)

For a given  $\Gamma \in [0, |J|]$ , the robust QAP is formulated as follows:

**RQAP**: 
$$\min_{X \in P} \max_{f \in V_T} \langle f, X \rangle.$$
 (14)

For any assignment  $X \in P$ , the value

$$Z(X) := \max_{f \in V_{\Gamma}} \langle f, X \rangle = \max_{\xi \in U_{\Gamma}} \langle f(\xi), X \rangle$$
(1)

is called the *worst-case* or *robust cost* for *X*. A maximizer in (15) is called a *worst-case scenario* for *X*, which is denoted by  $f^X$ . The corresponding *worst-case* perturbation vector is  $\xi^X$  where  $f^X = f(\xi^X)$ . RQAP seeks a feasible solution with the smallest robust cost. Let  $Z^*$  and  $X^*$  be the optimal objective value and solution for RQAP, respectively.

Let RC(X) denotes the difference between Z(X) and  $\langle f^-, X \rangle$ (i.e. the robust and optimistic costs of *X*, respectively) which is called the *robustness cost* of assignment *X*. In other words,

$$RC(X) := Z(X) - \langle f^-, X \rangle.$$
(16)

By substituting  $\langle f, X \rangle$  from (5) into (15), we have

$$Z(X) = \max_{\xi \in U_{\Gamma}} \left\{ \sum_{i=1}^{n} \sum_{j=1}^{n} f_{ij} d_{ij}^{X} \right\}$$
$$= \max_{\xi \in U_{\Gamma}} \left\{ \sum_{i=1}^{n} \sum_{j=1}^{n} (f_{ij}^{-} + \delta_{ij}\xi_{ij}) d_{ij}^{X} \right\}$$
$$= \sum_{i=1}^{n} \sum_{j=1}^{n} f_{ij}^{-} d_{ij}^{X} + \max_{\xi \in U_{\Gamma}} \left\{ \sum_{(i,j) \in J} \delta_{ij} d_{ij}^{X} \xi_{ij} \right\}$$
$$= \langle f^{-}, X \rangle + \max_{\xi \in U_{\Gamma}} \left\{ \sum_{(i,j) \in J} \delta_{ij} d_{ij}^{X} \xi_{ij} \right\}$$
(17)

where the first and last equalities hold by definition. The second equality is true because  $f_{ij} = f_{ii}^- + \delta_{ij}\xi_{ij}$ ,  $\forall i, j$ . The third equality

follows from the facts that for a given X,  $f_{ij}^- d_{ij}^X$  is a constant  $\forall i, j$ , and  $\delta_{ij} = 0$  if  $(i, j) \notin J$ . Eqs. (16) and (17) imply

$$RC(X) = \max_{\xi \in U_{\Gamma}} \left\{ \sum_{(i,j) \in J} \delta_{ij} d_{ij}^{X} \xi_{ij} \right\}.$$
 (18)

# 2.4. Worst-case scenario for a given assignment

In the following proposition, we show how to find a worst-case scenario for a given *X*.

**Proposition 1.** Consider a fixed  $\Gamma \in [0, |J|]$ , and a given  $X \in P$ . Based on the value of  $\Gamma$ , a worst-case scenario for X can be found as follows:

- If  $\Gamma$  is an integer, a scenario with  $\xi_{ij} = 1$  for  $\Gamma$  pairs  $(i, j) \in J$  with the largest values of  $\delta_{ij}d_{ij}^X$ , and  $\xi_{ij} = 0$  for the remaining pairs is a worst-case scenario for X.
- If  $\Gamma$  is not integer valued, a scenario with  $\xi_{ij} = 1$  for  $\lfloor \Gamma \rfloor$  pairs  $(i, j) \in J$  with the largest values of  $\delta_{ij}d_{ij}^X$ ,  $\xi_{ij} = (\Gamma \lfloor \Gamma \rfloor)$  for the next largest value of  $\delta_{ij}d_{ij}^X$ , and  $\xi_{ij} = 0$  for the remaining pairs is a worst-case scenario for X.

**Proof.** The distance matrix has non-negative elements which implies  $d_{ij}^X \ge 0$ , for all  $i, j \in N$ . Therefore,  $\langle f, X \rangle$  is a non-decreasing linear function of  $f \in V_{\Gamma}$ , or equivalently  $\langle f(\xi), X \rangle$  is a non-decreasing linear function of  $\xi \in U_{\Gamma}$ . That is, for any pair of matrices  $\tilde{f}, \hat{f} \in V_{\Gamma}$  if  $\tilde{f}_{ij} \le \hat{f}_{ij}$  for all  $(i, j) \in N$ , then  $\langle \tilde{f}, X \rangle \le \langle \hat{f}, X \rangle$ . Thus, there exists an extreme point  $\xi^X$  of polytope  $U_{\Gamma}$  such that  $\sum_{(i,j) \in J} \xi_{ij}^X = \Gamma$  and  $\xi^X$  is a worst-case scenario for X maximizing  $\langle f(\xi), X \rangle$  over  $\xi \in U_{\Gamma}$ . Note that in all extreme points of  $U_{\Gamma}$  with  $\sum_{(i,j) \in J} \xi_{ij}^X = \Gamma, \lfloor \Gamma \rfloor$  of the  $\xi_{ij}^X$  are equal to 1, up to one  $\xi_{ij}^X$  is  $(\Gamma - \lfloor \Gamma \rfloor)$ , and the remaining are 0.  $\Box$ 

It is worth mentioning that for a given assignment, finding the worst-case scenario requires sorting  $\delta_{ij}d_{ij}^X$  for  $(i, j) \in J$  which can be done in  $O(|J| \log(|J|))$ , i.e polynomial time in size of J and consequently in n. In contrast, obtaining the worst-case scenario in minmax regret QAP requires solving a general QAP which is NP-hard [45].

# 3. Mathematical formulations of robust QAP

In this section, we propose a mixed integer quadratic programming (MIQP) formulation of RQAP and a linear MIP equivalent.

#### 3.1. MIQP formulation of RQAP

An alternative formulation for RQAP (14) is

$$\min_{X \in P} \left\{ \langle f^{-}, X \rangle + \max_{\xi \in U_{\Gamma}} \left\{ \sum_{(i,j) \in J} \xi_{ij} \delta_{ij} d_{ij}^{X} \right\} \right\}$$

$$= \min_{X \in P} \left\{ \langle f^{-}, X \rangle + RC(X) \right\}.$$
(19)

An equivalent MIQP for (19) is proposed in Proposition 2.

Proposition 2. Problem (19) is equivalent to problem (20)-(21).

$$\min_{q\in\mathfrak{M}_+,r\in\mathfrak{M}_+^{|J|},X\in P} \langle f^-,X\rangle + \Gamma q + \sum_{(i,j)\in J} r_{ij}$$
(20)

$$q + r_{ij} \ge \delta_{ij} d_{ij}^X, \quad \forall (i,j) \in J.$$
(21)

**Proof.** By using the same idea as [46], one can substitute the inner maximization problem in (19) with its dual

$$\min_{q \in \mathcal{R}_+, r \in \mathfrak{N}_+^{|J|}} \Gamma q + \sum_{(i,j) \in J} r_{ij}$$
(22)

$$q + r_{ij} \ge \delta_{ij} d^X_{ij}, \quad \forall (i,j) \in J,$$
(23)

which results in problem (20)–(21).  $\Box$ 

Note that by definitions (4) and (5),  $d_{ij}^X$  and  $\langle f^-, X \rangle$  are quadratic functions of *x*. Thus, (20)–(21) is a MIQP problem with a quadratic objective function, |J| quadratic and 2*n* linear constraints,  $n^2$  binary and |J| + 1 continuous variables. To solve RQAP with an off the shelf MIP solver, we need to linearize the objective function and constraints in (20)–(21).

#### 3.2. Linearization of RQAP

Next, we extend [19] linearization to RQAP.

**Proposition 3.** Problem (24)-(27) is a MIP equivalent of problem (20)-(21).

$$\min_{\substack{X \in P, w \in \Re^{N \times N} \\ q \in \Re_{+,r} \in \Re^{|J|}}} \sum_{i=1}^{n} \sum_{k=1}^{n} (w_{ik} + f_{ii}^{-} d_{kk} x_{ik}) + \Gamma q + \sum_{(i,j) \in J} r_{ij}$$
(24)

$$w_{ik} \ge \sum_{\substack{j=1\\j\neq i}}^{n} \sum_{\substack{l=1\\l\neq k}}^{n} f_{j}^{-} d_{kl} x_{jl} - \hat{a}_{ik} (1 - x_{ik}), \quad \forall i, k \in N;$$
(25)

$$w_{ik} \ge l_{ik}x_{ik}, \quad \forall i, k \in N,$$

$$q + r_{ij} \ge \delta_{ij} \left( \sum_{l=1}^{n} d_{kl}x_{jl} + d_{k}^{\max}(x_{ik} - 1) \right),$$

$$\forall (i, j) \in J, \quad \forall k \in N$$

$$(26)$$

where,

$$l_{ik} = \min_{\substack{X \in P \\ x_{ik}=1}} \sum_{\substack{j=1 \\ j \neq i}}^{n} \sum_{\substack{l=1 \\ l \neq k}}^{n} f_{ij}^{-} d_{kl} x_{jl}, \quad \forall i, k \in N,$$

$$\hat{a}_{ik} = \max_{\substack{X \in P \\ x_{ik}=1}} \sum_{\substack{j=1 \\ j \neq i}}^{n} \sum_{\substack{l=1 \\ l \neq k}}^{n} f_{ij}^{-} d_{kl} x_{jl}, \quad \forall i, k \in N,$$

$$d_{k}^{\max} = \max_{l \in N} \{d_{kl}\}, \quad \forall k \in N.$$
(28)

**Proof.** Note that equivalence of problems (20)–(21) and (24)–(27) can be easily checked by using the same insights in [19]. First part of the objective function in (24) together with constraints (25) and (26) correspond to the minimization of  $\langle f^-, X \rangle$  in (20). Moreover, in any  $X \in P$ , for all  $(i, j) \in J$  and  $k \in N$ 

$$\sum_{l=1}^{n} d_{kl} x_{jl} + d_{k}^{\max}(x_{ik} - 1) = \begin{cases} \sum_{l=1}^{n} d_{kl} x_{jl} = d_{ij}^{X}, & \text{if } x_{ik} = 1, \\ \sum_{l=1}^{n} d_{kl} x_{jl} - d_{k}^{\max} \le 0, & \text{otherwise.} \end{cases}$$

Therefore, (27) is a valid linearization of (21).

Formulation (24)–(27) has  $2n^2 + n(2 + |J|)$  linear constraints and  $n^2$  binary and  $n^2 + |J| + 1$  continuous variables.

## 4. Heuristic algorithms

Due to difficulty of the regular QAP, only relatively small instances can be solved by exact methods [6]. To solve larger problems, a number of heuristics have been developed in the literature [36,6]. Since the robust QAP is at least as hard as regular QAP, heuristics are needed even more. In this section, we extend the 2-Opt local search [52] and tabu search [53,54] for classical QAP to solve RQAP.

# 4.1. 2-Opt local search

[52] used a local search based on 2-Opt neighborhood to solve the general QAP(7). In this method, a neighbor of a current solution  $X^1$  is obtained by swapping locations of any two facilities. Suppose that  $X^2$  is obtained from  $X^1$  by swapping locations of facilities  $a, b \in N, a \neq b$ . Then,

$$\phi_{i}^{X^{2}} = \begin{cases} \phi_{i}^{X^{1}}, & \forall i \neq a \text{ and } i \neq b, \\ \phi_{b}^{X^{1}}, & i = a, \\ \phi_{a}^{X^{1}}, & i = b. \end{cases}$$

The difference between the objective values of  $X^2$  and  $X^1$  is

$$\begin{split} \Delta_{ab} &= \sum_{i=1}^{n} \sum_{j=1}^{n} c_{ij\phi_{i}^{X^{2}}\phi_{j}^{X^{2}}} - \sum_{i=1}^{n} \sum_{j=1}^{n} c_{ij\phi_{i}^{X^{1}}\phi_{j}^{X^{1}}} \\ &= (c_{ab\phi_{b}^{X^{1}}\phi_{a}^{X^{1}}} - c_{ab\phi_{a}^{X^{1}}\phi_{b}^{X^{1}}}) + (c_{ba\phi_{a}^{X^{1}}\phi_{b}^{X^{1}}} - c_{ba\phi_{b}^{X^{1}}\phi_{a}^{X^{1}}}) \\ &+ (c_{aa\phi_{b}^{X^{1}}\phi_{b}^{X^{1}}} - c_{aa\phi_{a}^{X^{1}}\phi_{a}^{X^{1}}}) + (c_{bb\phi_{a}^{X^{1}}\phi_{a}^{X^{1}}} - c_{bb\phi_{b}^{X^{1}}\phi_{b}^{X^{1}}}) \\ &+ \sum_{i=1, i \neq a, b}^{n} [(c_{ai\phi_{b}^{X^{1}}\phi_{a}^{X^{1}}} - c_{ai\phi_{a}^{X^{1}}\phi_{b}^{X^{1}}}) + (c_{ia\phi_{i}^{X^{1}}\phi_{b}^{X^{1}}} - c_{ia\phi_{i}^{X^{1}}\phi_{a}^{X^{1}}})] \\ &+ \sum_{i=1, i \neq a, b}^{n} [(c_{ib\phi_{i}^{X^{1}}\phi_{a}^{X^{1}}} - c_{ib\phi_{i}^{X^{1}}\phi_{b}^{X^{1}}}) + (c_{bi\phi_{a}^{X^{1}}\phi_{b}^{X^{1}}} - c_{bi\phi_{b}^{X^{1}}\phi_{a}^{X^{1}}})]. \end{split}$$

In RQAP (19), the objective function consists of two parts  $\langle f^-, X \rangle$  and RC(X). Next, we extend the 2-Opt procedure to heuristically solve RQAP. For this purpose, we need to change the way of computing  $\Delta_{ab}$  values. For all  $(i, j) \in J$ , let  $g_{ij}^{\chi} := \delta_{ij} d_{ij}^{\chi}$ . For each pair of facilities *a* and *b*, let  $\Delta_{ab}$  be as follows:

$$\Delta_{ab} = \Delta_{ab}^1 + \Delta_{ab}^2, \tag{29}$$

where,

$$\begin{split} \Delta_{ab}^{1} &= \langle f^{-}, X^{2} \rangle - \langle f^{-}, X^{1} \rangle \\ &= \sum_{i=1}^{n} \sum_{j=1}^{n} f_{ij}^{-} d_{\phi_{i}^{X^{2}} \phi_{j}^{X^{2}}} - \sum_{i=1}^{n} \sum_{j=1}^{n} f_{ij}^{-} d_{\phi_{i}^{X^{1}} \phi_{j}^{X^{1}}} \\ &= (f_{ab}^{-} d_{\phi_{b}^{X^{1}} \phi_{a}^{X^{1}}} - f_{ab}^{-} d_{\phi_{a}^{X^{1}} \phi_{b}^{X^{1}}}) + (f_{ba}^{-} d_{\phi_{a}^{X^{1}} \phi_{b}^{X^{1}}} - f_{ba}^{-} d_{\phi_{b}^{X^{1}} \phi_{a}^{X^{1}}}) \\ &+ (f_{aa}^{-} d_{\phi_{b}^{X^{1}} \phi_{b}^{X^{1}}} - f_{aa}^{-} d_{\phi_{a}^{X^{1}} \phi_{a}^{X^{1}}}) + (f_{bb}^{-} d_{\phi_{a}^{X^{1}} \phi_{a}^{X^{1}}} - f_{bb}^{-} d_{\phi_{b}^{X^{1}} \phi_{b}^{X^{1}}}) \\ &+ \sum_{i=1, i \neq a, b}^{n} [(f_{ai}^{-} d_{\phi_{b}^{X^{1}} \phi_{i}^{X^{1}}} - f_{ai}^{-} d_{\phi_{a}^{X^{1}} \phi_{a}^{X^{1}}})] \\ &+ (f_{ia}^{-} d_{\phi_{i}^{X^{1}} \phi_{b}^{X^{1}}} - f_{ia}^{-} d_{\phi_{i}^{X^{1}} \phi_{a}^{X^{1}}})] \\ &+ \sum_{i=1, i \neq a, b}^{n} [(f_{ib}^{-} d_{\phi_{i}^{X^{1}} \phi_{a}^{X^{1}}} - f_{ib}^{-} d_{\phi_{i}^{X^{1}} \phi_{b}^{X^{1}}})] \\ &+ (f_{bi}^{-} d_{\phi_{a}^{X^{1}} \phi_{i}^{X^{1}}} - f_{bi}^{-} d_{\phi_{b}^{X^{1}} \phi_{i}^{X^{1}}})], \end{split}$$

$$(30)$$

and

$$\Delta_{ab}^{2} = RC(X^{2}) - RC(X^{1})$$

$$= \max_{\xi \in U_{\Gamma}} \left\{ \sum_{(i,j) \in J} \xi_{ij} \delta_{ij} d_{ij}^{\chi^{2}} \right\} - \max_{\xi \in U_{\Gamma}} \left\{ \sum_{(i,j) \in J} \xi_{ij} \delta_{ij} d_{ij}^{\chi^{1}} \right\}$$

$$= \max_{\xi \in U_{\Gamma}} \left\{ \sum_{(i,j) \in J} \xi_{ij} g_{ij}^{\chi^{2}} \right\} - \max_{\xi \in U_{\Gamma}} \left\{ \sum_{(i,j) \in J} \xi_{ij} g_{ij}^{\chi^{1}} \right\}.$$
(31)

Note that for the cases with  $i \neq a, b, j \neq a, b$ , we have  $g_{ij}^{X^2} = g_{ij}^{X^1}$ . Therefore, we only need to update  $g_{ij}$ , for other pairs of  $(i, j) \in J$  as follows.

$$\mathbf{g}_{ij}^{\chi^{2}} = \begin{cases} \delta_{ab}d_{\phi_{b}^{\chi^{1}}\phi_{a}^{\chi^{1}}}, & i = a, j = b \\ \delta_{ba}d_{\phi_{a}^{\chi^{1}}\phi_{b}^{\chi^{1}}}, & i = b, j = a \\ \delta_{aa}d_{\phi_{b}^{\chi^{1}}\phi_{b}^{\chi^{1}}}, & i = a, j = a \\ \delta_{bb}d_{\phi_{a}^{\chi^{1}}\phi_{b}^{\chi^{1}}}, & i = b, j = b \\ \delta_{aj}d_{\phi_{a}^{\chi^{1}}\phi_{a}^{\chi^{1}}}, & i = a, j \neq a, b \\ \delta_{bi}d_{\phi_{a}^{\chi^{1}}\phi_{b}^{\chi^{1}}}, & i = b, j \neq a, b \\ \delta_{bi}d_{\phi_{a}^{\chi^{1}}\phi_{b}^{\chi^{1}}}, & i \neq a, b, j = a \\ \delta_{bb}d_{\phi_{a}^{\chi^{1}}\phi_{b}^{\chi^{1}}}, & i \neq a, b, j = b. \end{cases}$$

To compute  $\Delta_{ab}^2$  in (31), we need to update  $g_{ij}^{X^2}$  for all  $(i, j) \in J$ , and then sort them. Afterward, by Proposition 1,  $RC(X^2)$  can be easily computed.

Suppose the algorithm is started from a random assignment and (29)–(31) are exploited to compute  $\Delta_{ab}$  for all  $a, b \in N, a \neq b$ . Let  $\Delta_{\min} = \min_{a,b} \{\Delta_{ab}\}$ . If  $\Delta_{\min} < 0$ , swap the locations of the corresponding facilities, otherwise terminate the algorithm. This procedure may be trapped in a local optimal solution. To remedy this issue, we can use multi random initial assignments or extend Taillard's general QAP tabu search [55] to solve RQAP.

# 4.2. Tabu search

Tabu search [53,54] is a technique to overcome local optimality in local search algorithms. For details of the general QAP tabu search, the reader is referred to [55]. A step of tabu search will be called a *swap* as it is based on an attempt to swap locations of two facilities. Next, we extend Taillard's tabu search procedure to solve RQAP. In this method, we use (29)–(31) to compute  $\Delta_{ab}$  and the rest of the procedure is similar to [55].

Note that the 2-Opt and tabu search algorithms do not provide any lower bound for RQAP. But, they provide upper bounds for RQAP while the robust (minmax regret) QAP tabu search in [45] provides neither a lower bound nor an upper bound.

## 5. Experimental results

### 5.1. Details of problem instances

We tested our proposed methods on uncertain QAP instances presented in [45]. These instances have been categorized into two main families.

• Euclidean Random (ER) instances: In this family, locations are chosen randomly on the 1000 × 1000 Euclidean grid as locations. For each pair of facilities *i*, *j* where  $i < j, f_{ij}^+ = f_{ij}^- = 0$  with probability 0.5, and with probability 0.5,  $f_{ij}^+ \in [0, 1000]$  and  $f_{ij}^- \in [0, f_{ij}^+]$  has been randomly generated. Based on the number of facilities and locations, *n*, these instances are categorized into groups ER-07, ER-08, ER-09, ER-10, ER-11, ER-12, ER-13, ER-15, ER-20, ER-25, and ER-30.

# QAPLIB instances: Instances of this family are generated from some classical QAP instances available in the QAPLIB library (http://www.seas.upenn.edu/qaplib/inst.html).

For more details on test instances see [45]. Similar to [45], 10 instances for each group of ER and QAPLIB families were used, and the average performance results are reported unless explicitly specified otherwise. The ER and QAPLIB groups were classified into "easy", "moderate" and "hard" categories as follows. We tried to solve the uncertain instance of the RQAP using CPLEX applied to MIP model (24)–(27) with time limit 7,200 s, and if proven optimality was achieved (respectively, not achieved) for all 10 instances and all tested values of  $\Gamma$ , we classified these instances as "easy" (respectively, "hard"). The remaining groups were classified as "moderate".

#### 5.2. Implementation details and parameters of the algorithms

The algorithms were coded in C++. IBM ILOG CPLEX, version 12.5.1, was used for solving the linearization of RQAP in the exact approaches. All experiments were conducted on a UNIX machine restricted to use of only one core with 2.27 GHz speed and 20 GB RAM.

In the 2-Opt algorithm, we started the method from a random assignment. The best possible swap was chosen to produce the largest reduction in the objective function. Swapping the locations of facilities was continued until there was no potential swap with cost reduction. The method restarted  $10 \times n^2$  times to reduce the chance of being trapped in a local optimal solution. In tabu search, we started the method from a random assignment with a termination criterion of  $100 \times n^2$  iterations.

In the Exact method, we used CPLEX (default parameters) to solve MIP model (24)–(27) with time limit 7,200 s and memory limit 20 GB. Relative optimality gap tolerance was set to  $10^{-6}$ . In addition, the CPLEX internal "MIPEmphasis" switch was set to "emphasize optimality over feasibility". For each instance RQAP was solved for all  $\Gamma \in \{1, 2, 4, 8, 16, 32, 64, 128, 256\}$  such that  $\Gamma \leq |J|$  for that instance.

#### 5.3. Illustrative example

The quality of the robust solutions are examined in this subsection. For this purpose, we compare the solutions of the RQAP model for an uncertain QAP with n = 9. The  $f^+$ ,  $f^-$  and d matrices are presented in Table 1. This example has |J| = 34 uncertain flows. The solutions of RQAP for all values of  $\Gamma \in [0, 34]$  are presented in Table 2. In this table,  $X^7$  is the minmax regret solution. In addition,  $Opt(f^-) = X^1$ ,  $Opt(f^=) = X^4$ , and  $Opt(f^+) = X^6$ .

Fig. 1 depicts robust cost, Z(X), of solutions  $X^1, \ldots, X^7$  versus  $\Gamma \in [0, 34]$ . In this figure, it is clear which assignment has the best (minimum) robust cost for each value of  $\Gamma$ . Note that Z(X) of  $X^4$  and  $X^5$  are very similar to  $X^3$ . Therefore, for clarity of the graphs in Fig. 2, the graphs corresponding to  $X^4$  and  $X^5$  are omitted.

For 10,000 runs we generated uncertain flow  $f_{ij}$ ,  $\forall (i, j) \in J$ , and simulated total cost. For this purpose, we used truncated normal, symmetric triangular, and uniform distributions in  $[f^-, f^+]$  to generate random flows. The results of the simulations are presented in Table 3 and Fig. 2. Let  $C(X) = \langle f, X \rangle$  be the random cost of assignment X, which depends on the realization of f. In Table 3, E(C(X)) and  $\sigma(C(X))$  are empirical estimations of the expectation and standard deviation of C(X), respectively. Moreover,  $C_{0.95}(X)$ and  $C_{0.99}(X)$  are the 0.95 and 0.99 quantiles of C(X), respectively.  $C_{max}(X)$  is the maximum observed C(X) in the simulation. For all three simulated distributions,  $X^4$  and  $X^6$  have the least E(C(X)) and  $\sigma(C(X))$ , respectively.  $X^4$  has also the least  $C_{0.95}(X)$  for normal and the least  $C_{0.95}(X)$  and  $C_{0.99}(X)$  for triangular distribution.  $X^3$  has the

Table 1	
Input data	of the illustrative example.

	Matrix f <sup>+</sup>						Matrix $f^-$				Matrix d																
	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
1	0	518	0	0	0	624	491	915	0	0	147	0	0	0	296	379	681	0	0	604	657	830	280	795	1340	246	593
2	518	0	577	0	0	0	183	0	0	147	0	94	0	0	0	122	0	0	604	0	515	544	404	633	1430	358	223
3	0	577	0	0	0	360	0	0	0	0	94	0	0	0	290	0	0	0	657	515	0	173	457	138	915	411	738
4	0	0	0	0	0	410	0	0	138	0	0	0	0	0	336	0	0	131	830	544	173	0	630	89	886	584	767
5	0	0	0	0	0	565	760	0	403	0	0	0	0	0	191	474	0	154	280	404	457	630	0	595	1060	130	593
6	624	0	360	410	565	0	217	0	683	296	0	290	336	191	0	25	0	547	795	633	138	89	595	0	797	549	856
7	491	183	0	0	760	217	0	128	592	379	122	0	0	474	25	0	6	302	1340	1430	915	886	1060	797	0	1190	1653
8	915	0	0	0	0	0	128	0	54	681	0	0	0	0	0	6	0	40	246	358	411	584	130	549	1190	0	463
9	0	0	0	138	403	683	592	54	0	0	0	0	131	154	547	302	40	0	593	223	738	767	593	856	1653	463	0

Table 2

Optimal solutions of the example for different protection levels.

Protection level	Solution
$0 \leq \Gamma \leq 0.40$	$X^1 = (2, 1, 7, 4, 5, 6, 8, 9, 3)$
$0.41 \le I' \le 1.04$ 1.05 < $\Gamma < 2.49$	$X^2 = (3, 4, 7, 1, 9, 5, 2, 6, 8)$ $X^3 = (8, 9, 2, 7, 3, 6, 5, 1, 4)$
$2.50 \le \Gamma \le 5.34$	$X^{4} = (8, 9, 2, 7, 3, 4, 5, 1, 6)$
$5.35 \leq \Gamma \leq 8.58$	$X^5 = (8, 2, 9, 7, 3, 4, 5, 1, 6)$
$8.59 \le I' \le 34$	$X^0 = (8, 2, 9, 7, 4, 5, 3, 1, 6)$
Minmax regret	$X^{7} = (2, 5, 1, 7, 3, 4, 8, 9, 6)$



**Fig. 1.** Robust cost of assignments  $x^1, \ldots, x^7$  versus protection level  $\Gamma$ .

minimum  $C_{0.99}(X)$  and  $C_{max}(X)$  for normal and triangular distributions, respectively. It has also the least  $C_{0.95}(X)$  and  $C_{0.99}(X)$  for uniform distribution. Minimum  $C_{max}(X)$  for normal and uniform distributions is found for  $X^5$ . Although  $Opt(f^+) = X^6$ ,  $X^6$  did not provide the minimum  $C_{max}(X)$  for any of the simulated distributions.

According to the results in Table 3 and Fig. 2, assignments  $X^3$ ,  $X^4$  and  $X^5$  outperform the other assignments almost in all of the metrics and distributions. As illustrated in this example, RQAP with budgeted uncertainty provides a pool of robust assignments depending on the value of  $\Gamma$ . Then, the decision makers can pick one of these assignments based on their risk preferences.

#### 5.4. Numerical results

In this subsection, we present the extensive results of implementing the proposed robust methods on "easy", "moderate" and "hard" instances. Recalling from Section 4, each of the proposed exact and heuristic methods for RQAP provides an upper bound for optimal robust cost of RQAP. Let  $ub_{EX}$ ,  $ub_{20}$  and  $ub_{TS}$  be the upper bounds obtained by Exact, 2-Opt and tabu search, respectively. Note that only the Exact method provides lower bounds, which we denote by  $lb_{EX}$ . We define  $gap_{EX}$  as

$$gap_{\rm EX} = \frac{ub_{\rm EX} - lb_{\rm EX}}{ub_{\rm EX}} \times 100.$$

Table 4 reports CPU time in seconds spent by CPLEX on Exact method to solve MIP model (24)–(27). This time is denoted by  $t_{\text{EX}}$ . |J| indicates the number of uncertain flows. In all of these instances, Exact method was able to find the optimal solution, i.e.  $gap_{\text{EX}} = 0$  and  $ub_{\text{EX}} = lb_{\text{EX}}$  for all easy instances.

According to Table 4, for almost all easy instances, increasing  $\Gamma$  results in a significant increase in  $t_{\text{EX}}$ . This observation is also true for moderate instances in Table 10. That means that increasing  $\Gamma$  loosens linearization (24)–(27) for RQAP.

By comparing  $t_{\text{EX}}$  in Table 4 with the time spent to obtain minmax regret solutions (Tables 2 and 3 in [45]), it is observed that even for large values of  $\Gamma$ ,  $t_{\text{EX}}$  is about 10%–30% of minmax regret solution time for most of the easy instances.

Table 5 presents the CPU time in seconds spent on 2-Opt  $(t_{20})$  and tabu search  $(t_{TS})$  to heuristically solve RQAP for easy instances. Unlike  $t_{EX}$ , increasing  $\Gamma$  did not affect  $t_{20}$  and  $t_{TS}$  systematically. Therefore, for moderate and hard instances, we only report average CPU times for different values of  $\Gamma$ .

To assess the quality of solutions obtained from heuristics, we used  $ub_{20}$  and  $ub_{TS}$ , upper bounds from heuristics, and  $lb_{EX}$ , the best lower bound from the Exact method. We defined  $gap_{20}$  and  $gap_{TS}$  as

$$gap_{20} = \frac{ub_{20} - lb_{EX}}{ub_{20}} \times 100$$
 and  $gap_{TS} = \frac{ub_{TS} - lb_{EX}}{ub_{TS}} \times 100.$ 

In group chr15c, for  $\Gamma = 1, 2, 8$  and 16 the  $gap_{TS}$  was 0.01%, 0.08%, 0.06% and 0.47%, respectively. This gap for chr15b and  $\Gamma = 8$  was 0.04%. For the rest of cases, tabu search found the optimal robust solutions. For chr15c and  $\Gamma = 1, 2, 4, 8$  and 16,  $gap_{20}$  was 0.01%, 0.03%, 0.33%, 0.52% and 0.12%, respectively. This gap for chr15a and  $\Gamma = 1, 2, 4$  and 8 was 0.10%, 0.07%, 0.24% and 0.38%, respectively. Moreover,  $gap_{20} = 0.18\%$  for chr12c and  $\Gamma = 8$ . For the rest of cases, 2-Opt found the optimal robust solutions.

Tables 6–11 represent the numerical results for moderate instances. For each group and  $\Gamma$ , Table 6 shows the number of instances (out of 10) which were solved optimally by Exact method within 7,200 s. Some moderate instances were not solved optimally, and a positive optimality gap remained for these cases. Table 7 reports the average optimality gap (among 10 instances) for each group and  $\Gamma$ . Tables 8 and 9 present the number (in thousands) of all nodes and the remaining nodes, respectively, in the branch-and-cut tree explored by CPLEX.

In all of the moderate instances,  $ub_{20} = ub_{TS} \le ub_{EX}$ . In other words, the exact method was not able to outperform the heuristics in any of the moderate instances. Moreover, in all of these instances, 2-Opt and tabu search got the same solutions, which outperform the solutions of Exact method.

Table 10 reports  $t_{EX}$  for moderate instances.  $t_{20}$  and  $t_{TS}$  for these instances are reported in Table 11. Moreover, the effective time



Fig. 2. Probability distributions of total cost for assignments x<sup>1</sup>, x<sup>2</sup>, x<sup>3</sup>, x<sup>6</sup>, and x<sup>7</sup> in simulation results from normal, triangular, and uniform distributions for uncertain material flows.

Table 3

Summary of the simulation for assignments  $x^1 - x^7$  from normal, triangular, and uniform distributions.

	Results fr	Results from truncated normal distribution				Results fro	om symmeti	ic triangula	r distributi	on	Results from uniform distribution				
Χ	E(C(X))	$\sigma(C(X))$	$C_{0.95}(X)$	$C_{0.99}(X)$	$C_{max}(X)$	E(C(X))	$\sigma(C(X))$	$C_{0.95}(X)$	$C_{0.99}(X)$	$C_{max}(X)$	E(C(X))	$\sigma(C(X))$	$C_{0.95}(X)$	$C_{0.99}(X)$	$C_{max}(X)$
$X^1$	4925943	183732	5231124	5353466	5593676	4918265	225385	5287349	5429806	5654092	4922488	318317	5450792	5633086	5983884
$X^2$	4625630	130569	4840172	4931579	5068223	4621308	161304	4887201	4984846	5196267	4623809	225544	4996801	5131568	5374541
$X^3$	4591398	91583	4741441	4801663	4917547	4587817	113873	4778583	4855632	5044375	4592330	157627	4853489	4941902	5160347
$X^4$	4588999	92627	4740427	4803665	4913413	4585379	115035	4778064	4852616	5047118	4590022	159355	4854413	4944834	5170011
$X^5$	4606422	89165	4753007	4812793	<u>4913326</u>	4603112	111119	4788751	4863401	5048748	4607485	153171	4860528	4947754	5154671
$X^{6}$	4674114	83334	4809191	4868794	4992954	4671868	102937	4840905	4907941	5108466	4672872	143334	4906776	5003099	5161551
$X^7$	4678934	89033	4824954	4881580	4990871	4675413	110900	4860870	4934662	5081722	4679470	153151	4931791	5026293	5235742

5

Table 4

CPU time spent on the Exact method  $(t_{EX})$  for easy instances.

		Protection level $\Gamma$								
Group	IJ	1	2	4	8	16	32	64		
chr12a	22	2	4	6	9	18	-	-		
chr12b	22	0.6	0.9	1	1.7	2.4	-	-		
chr12c	22	6	6	8	14	22	-	-		
chr15a	28	39	36	49	90	123	-	-		
chr15b	28	7	8	11	24	38	-	-		
chr15c	28	53	81	129	362	579	-	-		
esc16a	76	33	15	31	56	184	278	235		
esc16d	42	17	31	26	26	28	34	-		
esc16e	42	1.1	1.1	1.3	1.5	1.8	1.8	-		
esc16g	42	2	2.1	3	4	7	8	-		
esc16i	30	2	2	2.2	2.1	1.7	-	-		
esc16j	24	0.3	0.3	0.4	0.4	0.4	-	-		
ER07	20	0.1	0.2	0.3	0.4	0.6	-	-		
ER08	29	0.5	0.5	0.7	1.2	2.5	5	-		
ER09	31	2.3	3	4	7	11	22	-		
ER10	43	10	12	20	32	65	169	-		
ER11	55	83	95	114	129	235	647	2865		
scr12	56	8	10	10	15	30	68	-		
tai10a	90	41	46	54	67	115	175	397		
tai10b	90	5	7	8	8	11	15	15		
tai12b	127	11	12	13	17	24	17	13		

 $t_{20}^{\text{eff}}$  ( $t_{\text{TS}}^{\text{eff}}$ ) and the effective iteration iter $t_{20}^{\text{eff}}$  (iter $t_{\text{TS}}^{\text{eff}}$ ) for 2-Opt (tabu search), which are the time and iteration count of discovering the best solution in these algorithms, are reported in this table. We see that the robust cost was not improved by 2-Opt or tabu search after their effective time and iteration count. Based on the results in Table 11,  $t_{20}^{\text{eff}}$  and  $t_{\text{TS}}^{\text{eff}}$  are almost 5% of  $t_{20}$  and  $t_{\text{TS}}$ , respectively. Note that  $t_{20}$  and  $t_{TS}$  depend on iteration limits that we set for 2-Opt and tabu search, respectively. Similarly, we denoted the number of effective restarts in tabu search by Restart<sup>eff</sup><sub>20</sub>.

Hard instances were solved heuristically by 2-Opt and tabu search methods for all values of the protection level  $\Gamma \in \{1, 2, 4, ..., r\}$ 8, 16, 32, 64, 128, 256} such that  $\Gamma \leq \Gamma_{\text{max}}$ , and the results are reported in Table 12. We omitted Exact method for these instances, because there was no hope to solve them exactly. For these cases,

we reported 
$$gap_{20}^{\#}$$
 and  $gap_{TS}^{\#}$  in Table 12 which are defined as

$$gap_{20}^{\#} = \frac{ub_{20} - \min\{ub_{20}, ub_{TS}\}}{ub_{20}} \times 100$$
  
and

$$gap_{TS}^{\#} = \frac{ub_{TS} - \min\{ub_{20}, ub_{TS}\}}{ub_{TS}} \times 100.$$

Note that for the cases with the same value of *n*, increasing |J|results in a significant increase in  $t_{20}$  and  $t_{TS}$ . This is because, in each iteration of 2-Opt and tabu search, computing  $\Delta_{ab}^2$  for each pair of facilities  $a, b \in N$  in (31) requires sorting a vector of length |J|.

Recalling from Section 5.2, we used an iteration limit of  $100 \times n^2$ for tabu search and  $10 \times n^2$  restarts for 2-Opt. With this setting, according to Tables 11 and 12, for most of the moderate and hard instances,  $t_{20}$  and  $t_{TS}$  are very close to each other. But,  $t_{TS}^{eff}$  and iter $_{TS}^{eff}$  are almost half of  $t_{20}^{eff}$  and iter $_{20}^{eff}$ , respectively, in most cases. Furthermore, based on the values of  $gap_{20}^{\#}$  and  $gap_{TS}^{\#}$  in Table 12, for most hard instances, the quality of solutions obtained from tabu search is better than 2-Opt.

By comparing  $t_{20}$  and  $t_{TS}$  in Table 12, with the time spent to obtain approximate minmax regret solutions (Tables 6 and 7 in [45]), it is observed that  $t_{20}$  and  $t_{TS}$  are about 10%–30% of minmax regret solution time for most hard instances.

#### 6. Conclusions and future research

We studied the robust quadratic assignment problem where uncertain interfacility flows belong to a budgeted uncertainty set. Unlike minmax regret QAP, in the robust optimization approach with budgeted uncertainty framework, finding the worst-case scenario and evaluating the robust cost of a given assignment can be done in polynomial time. Therefore, the proposed heuristic and exact methods for RQAP in this paper are significantly faster than heuristic and exact methods, respectively, for minmax regret QAP. Moreover, the robust approach with budgeted uncertainty has an adjustable conservativeness, and the decision maker can get a pool of solutions with different levels of conservativeness. The main contributions of the paper are as follows:

### **Table 5** CPU time spent on 2-Opt $(t_{20})$ and tabu search $(t_{TS})$ for easy instances.

	Protect	tion level $\Gamma$														
	<i>t</i> <sub>20</sub> (s)								$t_{\rm TS}(s)$							
Group	1	2	4	8	16	32	64	Avg	1	2	4	8	16	32	64	Avg
chr12a	1.2	1.3	1.1	1.1	1.2	-	-	1.2	1.5	1.4	1.4	1.4	1.6	-	-	1.5
chr12b	1.3	1.3	1.1	1.2	1.4	-	-	1.3	1.5	1.5	1.4	1.5	1.6	-	-	1.5
chr12c	1.2	1.2	1.1	1.1	1	-	-	1.1	1.5	1.4	1.4	1.3	1.4	-	-	1.4
chr15a	5	4.7	4.3	4.9	4.7	-	-	4.7	4.4	4.2	4.2	4.4	4.4	-	-	4.3
chr15b	4.4	5.1	5.1	4.6	4.8	-	-	4.8	4.4	4.7	5	4.6	4.4	-	-	4.6
chr15c	4.2	3.8	4	4.2	4.1	-	-	4.1	4	4.3	4.1	4.8	4.7	-	-	4.4
esc16a	11	11	12	11	11	11	12	11	16	15	15	15	15	16	16	15
esc16d	6.8	6.8	6.5	7.1	6.7	6.5	-	6.7	8.1	7.8	8.6	8.1	8.7	7.7	-	8.2
esc16e	5.9	6	5.8	5.9	5.3	5.8	-	5.8	8.5	8.6	8.2	9.2	8.3	8.6	-	8.6
esc16g	5.1	4.9	5.1	5.7	5.1	5.5	-	5.2	7.7	7.3	8.2	8.9	8.4	8.5	-	8.2
esc16i	3.6	3.8	3.6	3.5	3.6	-	-	3.6	5.6	6.2	5.9	5.7	6.2	-	-	5.9
esc16j	2.9	2.6	2.9	2.8	2.7	-	-	2.8	5.2	5.1	5.4	5.6	5.5	-	-	5.4
ER07	0.05	0.05	0.05	0.05	0.05	-	-	0.05	0.1	0.1	0.1	0.1	0.1	-	-	0.1
ER08	0.1	0.1	0.1	0.1	0.1	0.2	-	0.1	0.2	0.2	0.2	0.2	0.2	0.3	-	0.2
ER09	0.3	0.3	0.3	0.3	0.3	0.3	-	0.3	0.5	0.4	0.5	0.5	0.5	0.6	-	0.5
ER10	0.9	0.8	0.8	1	0.9	0.9	-	0.9	1.3	1.3	1.2	1.4	1.3	1.3	-	1.3
ER11	1.9	1.8	1.7	2.1	2	2	2.7	1.9	2.5	2.3	2.3	2.7	2.4	2.7	3.3	2.5
scr12	2.7	2.8	2.8	2.5	2.8	2.8	-	2.7	3.3	3.5	3.7	3.1	3.7	3.8	-	3.5
tai10a	1.9	1.8	1.8	1.7	1.9	1.8	1.8	1.8	3.2	3.1	3.1	2.9	3.2	3.1	3.1	3.1
tai10b	1.9	2	1.9	1.9	1.8	1.7	1.9	1.9	2.8	3	2.9	2.9	2.7	2.6	2.9	2.8
tai12b	7.1	8.2	7.9	7.8	8.1	7.7	7.5	7.7	8.9	9.3	9.1	9	9.1	8.4	8.6	8.9

#### Table 6

Number of optimally solved moderate instances by the exact method.

28
20

### Table 7

Optimality gap (%) for moderate instances using exact method.

	Protection level $\Gamma$											
Group	1	2	4	8	16	32	64	128				
chr18b	0	0	0	0.21	6.1	11.6	-	-				
had12	0	0	0	0	0.2	1.1	15.2	26.6				
nug12	0	0	0	0	0	1.1	12.5	12.1				
rou12	0	0	0	0	0	2.4	12.9	19.9				
ER12	0	0	0	0	0	0.14	-	-				
ER13	0.08	0.17	0.24	0.68	1.7	7	18.8	-				
tai12a	0	0	0	0	0	0.81	7.9	13.9				

#### Table 8

Number of all branch-and-cut nodes (in thousands) for moderate instances.

	Protection level $\Gamma$												
Group	1	2	4	8	16	32	64	128					
chr18b	5.7	8.8	87	826	1720	2078	-	-					
had12	206	195	222	337	490	907	956	501					
nug12	361	403	536	629	864	1277	1009	1077					
rou12	342	402	438	565	788	1080	893	825					
ER12	241	218	203	297	551	1359	-	-					
ER13	1191	1349	1407	1396	1491	1705	1360	-					
tai12a	308	327	376	372	570	838	891	864					

- 1. A mathematical programming formulation of the RQAP under budgeted uncertainty.
- 2. A linearized MIP formulation for RQAP which was solved by CPLEX as an exact method.
- 3. Two heuristic methods (2-Opt and tabu search) to solve RQAP.
- 4. Extensive experimental results.

#### Table 9

Number of remaining branch-and-cut nodes (in thousands) for moderate instances.

	Protection level $\Gamma$												
Group	1	2	4	8	16	32	64	128					
chr18b	0	0	0	110	861	1395	-	-					
had12	0	0	0	0	84	176	814	465					
nug12	0	0	0	0	0	107	575	631					
rou12	0	0	0	0	0	221	581	691					
ER12	0	0	0	0	0	23	-	-					
ER13	55	107	93	220	389	897	1105	-					
tai12a	0	0	0	0	0	104	400	610					

Table 10 CPU time s	spent of	n Exact m	ethod ( <i>t</i> E	<sub>x</sub> ) for mo	derate in	stances.
	Prote	ection lev	el Γ			
Group	1	2	4	8	16	32

Group	1	2	4	8	16	32	64	128
chr18b	28	42	225	2245	5797	6948	-	-
had12	610	613	737	1136	1760	4140	7200	7200
nug12	971	1088	1486	2052	3325	5558	7200	7047
rou12	960	1179	1370	1833	3264	5673	7200	7200
ER12	419	364	391	613	1209	4300	-	-
ER13	2307	2494	2965	3052	3920	6198	7200	-
tai12a	955	985	1124	1167	2307	4209	7096	7200

Table 11	
Heuristic results for moderate instances.	

	2-Opt				tabu search			
Group	t <sub>20</sub>	$t_{20}^{\rm eff}$	iter <sup>eff</sup> <sub>20</sub>	Restart <sup>eff</sup>	t <sub>TS</sub>	$t_{\rm TS}^{\rm eff}$	$iter_{TS}^{eff}$	
chr18b	11.1	0.62	1605	165	11.7	0.31	932	
had12	9.3	0.37	454	71	7.1	0.22	325	
nug12	9	0.93	1234	214	6	0.65	1009	
rou12	9.3	0.49	592	99	6.7	0.25	356	
ER12	4.7	0.06	171	23	4	0.13	391	
ER13	6.9	0.17	338	41	6.6	0.19	457	
tai12a	9.9	0.54	663	110	7.1	0.21	302	

The main conclusions are as follows:

- 1. The Exact method is well-suited for easy instances (e.g. instances with  $n \leq 11$ ).
- 2. In the Exact method, CPU time significantly increases by increasing the protection level  $\Gamma$ .

#### Table 12

Experimental results for hard instances.

			2-Opt					tabu search			
Group	IJ	$\Gamma_{\rm max}$	t <sub>20</sub>	$t_{20}^{\rm eff}$	iter <sup>eff</sup> <sub>20</sub>	Restart <sup>eff</sup>	$gap_{20}^{\#}$	t <sub>TS</sub>	$t_{ m TS}^{ m eff}$	iter <sup>eff</sup>	$gap_{TS}^{\#}$
esc16b	184	128	39	0.5	274	34	0	43	0.3	164	0
esc16c	101	64	20	0.3	332	41	0	22	0.2	220	0
esc16h	229	128	55	0.4	155	17	0	54	0.5	221	0
had14	181	128	24	1.8	1120	139	0	26	0.7	530	0
had18	305	256	137	40	9770	947	0.07	120	11	2692	0
lipa20a	368	256	262	133	21017	1999	0.39	222	82	14634	0.01
lipa20b	358	256	242	117	20583	1931	0.73	210	43	8068	0.02
nug14	181	128	18	3.2	2433	340	0.03	22	1.7	1342	0
nug15	210	128	29	8.3	4960	640	0.12	32	3.3	2269	0
nug16a	240	128	45	15	7315	887	0.28	48	6.5	3496	0
nug16b	238	128	42	16	8079	985	0.34	45	5.6	3175	0
nug17	272	256	68	27	10182	1147	0.45	68	11	4721	0.01
rou15	210	128	29	7.4	4513	584	0.02	32	2.6	1836	0
rou20	378	256	245	128	22702	2098	0.74	205	58	11232	0.01
ER15	108	128	22	1	1012	102	0	20	0.8	908	0
ER20	192	128	177	24	8126	541	0	112	12	4109	0
ER25	291	256	802	238	37447	1850	0.03	378	69	11321	0
ER30	432	256	3671	1608	104384	3940	0.08	1351	305	20199	0.01
scr15	84	64	12	0.8	1448	150	0	11	1	2012	0
scr20	124	64	83	24	15921	1149	0.04	56	9.2	6509	0
tai15a	208	128	37	10	4909	625	0.02	42	4	2212	0
tai15b	210	128	42	2.3	1210	116	0	38	0.8	480	0
tai17a	266	256	81	35	10429	1206	0.32	84	17	5758	0
tai20a	374	256	239	127	22472	2128	0.94	208	59	11260	0.01
tai20b	380	256	262	117	24728	1775	1.47	176	57	12832	0.01

- 3. In 2-Opt and tabu search, for the cases with the same *n*, CPU times significantly increase by increasing the number of uncertain flows |*J*|.
- 4. In all moderate instances, 2-Opt and tabu search methods outperformed the Exact method.
- 5. In most hard instances, in the same time and number of iterations, tabu search outperformed 2-Opt. Thus, tabu search is recommended for larger instances.

A possible direction for future research may be developing tighter formulations for the linear equivalent of RQAP. Moreover, researchers can design other heuristic methods to solve RQAP and compare the results with 2-Opt and tabu search.

#### References

- [1] Koopmans TC, Beckmann M. Assignment problems and the location of economic activities. Econometrica 1957;25:53–76.
- [2] Lawler EL. The quadratic assignment problem. Manage Sci 1963;9(4):586–99.
- [3] Pierskalla WP. The multi-dimensional assignment and quadratic assignment problems. In: Technical memorandum no. 93. OH: Case Western Reserve University, Operations Research Department, School of Management Cleveland; 1967
- 1967.
  [4] Mezmaz M, Mehdi M, Bouvry P, Melab N, Talbi E-G, Tuyttens D. Solving the three dimensional quadratic assignment problem on a computational grid. Cluster Comput 2014;17(2):205–17.
- [5] Çela E. The quadratic assignment problem: theory and algorithms. 1998. Dordrecht: Kluwer Academic Publishers; 1998.
- [6] Burkard RE, Dell'Amico M, Martello S. Assignment problems. Philadelphia: SIAM; 2009. Revised Reprint.
- [7] Sahni S, Gonzalez T. P-complete approximation problems. J ACM (JACM) 1976; 23(3):555–65.
- [8] Adams W, Waddell L. Linear programming insights into solvable cases of the quadratic assignment problem. Discrete Optim 2014;14:46–60.
- [9] Laurent M, Seminaroti M. The quadratic assignment problem is easy for robinsonian matrices with toeplitz structure. Oper Res Lett 2015;43(1):103–9.
- [10] Muenvanichakul S, Charnsethikul P. The approximated dynamic programming approach to the dynamic quadratic assignment problem. Thammasat Int J Sci Tech 2007;12(2).
- [11] Ahmed ZH. Experimental analysis of crossover and mutation operators on the quadratic assignment problem. Ann Oper Res 2015;1–19.
- [12] Tosun U, Dokeroglu T, Cosar A. A robust island parallel genetic algorithm for the quadratic assignment problem. Int J Prod Res 2013;51(14):4117–33.
- [13] Tosun U. On the performance of parallel hybrid algorithms for the solution of the quadratic assignment problem. Eng Appl Artif Intell 2015;39:267–78.
- [14] Zhang Q, Sun J, Tsang E, Ford J. Estimation of distribution algorithm with 2opt local search for the quadratic assignment problem. In: Towards a new evolutionary computation. Springer: 2006, p. 281–92.
- evolutionary computation. Springer; 2006. p. 281–92.[15] Dokeroglu T. Hybrid teaching–learning-based optimization algorithms for the quadratic assignment problem. Comput Ind Eng 2015;85:86–101.

- [16] De Klerk E, Sotirov R. Exploiting group symmetry in semidefinite programming relaxations of the quadratic assignment problem. Math Program 2010;122(2): 225–46.
- [17] Peng J, Zhu T, Luo H, Toh K-C. Semi-definite programming relaxation of quadratic assignment problems based on nonredundant matrix splitting. Comput Optim Appl 2015;60(1):171–98.
- [18] Kaufman L, Broeckx F. An algorithm for the quadratic assignment problem using Bender's decomposition. Eur J Oper Res 1978;2(3):207–11.
- [19] Xia Y, Yuan Y-X. A new linearization method for quadratic assignment problems. Optim Methods Softw 2006;21(5):805–18.
- [20] Nyberg A, Westerlund T. A new exact discrete linear reformulation of the quadratic assignment problem. Eur J Oper Res 2012;220(2):314–9.
- [21] Nyberg A, Westerlund T, Lundell A. Improved discrete reformulations for the quadratic assignment problem. In: Integration of AI and OR techniques in constraint programming for combinatorial optimization problems. Springer; 2013. p. 193–203.
- [22] Zhang H, Beltran-Royo C, Ma L. Solving the quadratic assignment problem by means of general purpose mixed integer linear programming solvers. Ann Oper Res 2013;207(1):261–78.
- [23] Hahn PM, Zhu Y-R, Guignard M, Hightower WL, Saltzman MJ. A level-3 reformulation-linearization technique-based bound for the quadratic assignment problem. INFORMS J Comput 2012;24(2):202–9.
- [24] Rostami B, Malucelli F. A revised reformulation-linearization technique for the quadratic assignment problem. Discrete Optim 2014;14:97–103.
- [25] De Klerk E, Marianna E, Sotirov R, Truetsch U, et al. Symmetry in rlt-type relaxations for the quadratic assignment and standard quadratic optimization problems. Eur J Oper Res 2014;233(3):488–99.
- [26] Zhang H, Beltran-Royo C, Constantino M. Effective formulation reductions for the quadratic assignment problem. Comput Oper Res 2010;37(11):2007–16.
- [27] Fischetti M, Monaci M, Salvagnin D. Three ideas for the quadratic assignment problem. Oper Res 2012;60(4):954–64.
- [28] Steinberg L. The backboard wiring problem: A placement algorithm. SIAM Rev 1961;3(1):37–50.
- [29] Geoffrion AM, Graves G. Scheduling parallel production lines with changeover costs: Practical application of a quadratic assignment/LP approach. Oper Res 1976;24(4):595–610.
- [30] Heffley DR. The quadratic assignment problem: A note. Econometrica 1972; 1155–63.
- [31] Pollatschek M, Gershoni N, Radday Y. Optimization of the typewriter keyboard by simulation. Angew Inform 1976;17(0):438–9.
- [32] Dickey J, Hopkins J. Campus building arrangement using topaz. Transp Res 1972;6(1):59–68.
- [33] Elshafei AN. Hospital layout as a quadratic assignment problem. Oper Res Q 1977;28(1):167–79.
- [34] Francis RL, White JA, McGinnis LF. Facility layout and location: an analytical approach. vol. 31. NJ: Prentice-Hall Englewood Cliffs; 1974.
- [35] Alkaya AF, Duman E. Combining and solving sequence dependent traveling salesman and quadratic assignment problems in PCB assembly. Discrete Appl Math 2015.
- [36] Loiola EM, de Abreu NMM, Boaventura-Netto PO, Hahn P, Querido T. A survey for the quadratic assignment problem. Eur J Oper Res 2007;176(2):657–90.

- [37] Hahn PM, Zhu Y-R, Guignard M, Smith JM. Exact solution of emerging quadratic assignment problems. Int Trans Oper Res 2010; 17(5):525-52.
- [38] Drezner Z. The quadratic assignment problem. In: Location science. Springer; 2015. p. 345-63.
- Feizollahi MJ, Modarres M. Robust quadratic assignment problem with [39] uncertain locations. Iran J Oper Res 2012;3(2):46-65.
- [40] Madhusudanan Pillai V, Hunagund IB, Krishnan KK. Design of robust layout for dynamic plant layout problems. Comput Ind Eng 2011;61(3):813-23.
- [41] Liu L, Li Y. The fuzzy quadratic assignment problem with penalty: New models and genetic algorithm. Appl Math Comput 2006; 174(2): 1229-44.
- [42] Snyder LV. Facility location under uncertainty: a review. IIE Trans 2006;38(7): 547–64. [43] Zhao Y, Wallace SW. Integrated facility layout design and flow assignment
- problem under uncertainty. INFORMS | Comput 2014;26(4):798–808.
- Feyzollahi M, Shokouhi A, Yazdi MM, Tarokh M. Designing a model for optimal [44] hospital unit layout. Pejouhandeh 2009; 14(4): 191-8.
- Feizollahi MJ, Averbakh I. The robust (minmax regret) quadratic assignment [45] problem with interval flows. INFORMS J Comput 2014;26(2):321-35.
- [46] Bertsimas D, Sim M. Robust discrete optimization and network flows. Math Program 2003;98(1-3):49-71.

- [47] Bertsimas D, Sim M. The price of robustness. Oper Res 2004;52(1):35-53.
- [48] Feizollahi MJ, Modarres M. The robust deviation redundancy allocation problem with interval component reliabilities. IEEE Trans Reliab 2012;61(4): 057\_65
- [49] Soltani R, Sadjadi SJ, Tavakkoli-Moghaddam R. Robust cold standby redundancy allocation for nonrepairable series-parallel systems through Min-Max regret formulation and Benders' decomposition method. Proc Inst Mech Eng O 2014;228(3):254-64.
- [50] Feizollahi MI, Ahmed S, Modarres M, The robust redundancy allocation problem in series-parallel systems with budgeted uncertainty. IEEE Trans Reliab 2014:63(1):239-50.
- [51] Feizollahi MJ, Soltani R, Feyzollahi H. The robust cold standby redundancy allocation in series-parallel systems with budgeted uncertainty. IEEE Trans Reliab 2015:64(2):799-806.
- [52] Buffa ES, Armour GC, Vollmann TE. Allocating facilities with CRAFT. Harvard University; 1964.
- [53] Glover F. Tabu search part I. ORSA J Comput 1989;1(3):190–206.
   [54] Glover F. Tabu search part II. ORSA J Comput 1990;2(1):4–32.
- [55] Taillard E. Robust taboo search for the quadratic assignment problem. Parallel Comput 1991;17(4):443-55.