

Foster, Dean P.

Working Paper

A Proof of Calibration Via Blackwell's Approachability Theorem

Discussion Paper, No. 1182

Provided in Cooperation with:

Kellogg School of Management - Center for Mathematical Studies in Economics and Management Science, Northwestern University

Suggested Citation: Foster, Dean P. (1997) : A Proof of Calibration Via Blackwell's Approachability Theorem, Discussion Paper, No. 1182, Northwestern University, Kellogg School of Management, Center for Mathematical Studies in Economics and Management Science, Evanston, IL

This Version is available at:

<https://hdl.handle.net/10419/221538>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

Discussion Paper No. 1182

**A PROOF OF CALIBRATION VIA
BLACKWELL'S APPROACHABILITY THEOREM**

by

Dean P. Foster*

Department of Statistics

Wharton School, The University of Pennsylvania

February 27, 1997

* Foster is also a visiting professor in the Center for Mathematical Studies in Economics and Management Science at Northwestern University, Winter 1997.

A proof of Calibration via Blackwell's Approachability Theorem

Dean P. Foster*

February 27, 1997

Abstract

Over the past few years many proofs of calibration have been presented (Foster and Vohra (1991, 1997), Hart (1995), Fudenberg and Levine (1995), Hart and Mas-Colell (1996)). Does the literature really need one more? Probably not, but this algorithm for being calibrated is particularly simple and doesn't require a matrix inversion. Further the proof follows directly from Blackwell's approachability theorem. For these reasons it might be useful in the class room.

*This work was done while I was visiting the Center for Mathematical Studies in Economics and Management Science, Northwestern University. Permanent affiliation: Dept. of Statistics, The Wharton School, University of Pennsylvania, Philadelphia, PA 19104. Email: foster@bellsark.wharton.upenn.edu.

Thanks to Sergiu Hart who provided the proof of the only result in the paper. The algorithm is a modification of the original algorithm in Foster and Vohra (1991).

Suppose at time t a forecast, f_t , is made which takes on the value of the midpoint of each of the intervals $[0, 1/m]$, $[1/m, 2/m]$, \dots , $[(m-1)/m, 1]$, namely, $\frac{2i-1}{2m}$ for i equals 1 to m . Let A_t^i the vector of indicators as to which forecast is actually made:

$$A_t^i = \begin{cases} 1 & \text{if } f_t = \frac{2i-1}{2m} \\ 0 & \text{otherwise} \end{cases}$$

Let X_t be the outcome at time t . We can now define the empirical frequency ρ_t^i as:

$$\rho_T^i = \frac{\sum_{t=1}^T X_t A_t^i}{\sum_{t=1}^T A_t^i}$$

Hopefully, ρ_T^i lies in the interval $[\frac{i-1}{m}, \frac{i}{m}]$. If so, the forecast is approximately calibrated. If not, I will measure how far outside the interval it is by two distances: \bar{d}_T^i and \bar{e}_T^i (for deficit and excess) which are defined as:

$$\begin{aligned} \bar{d}_T^i &= \frac{1}{T} \sum_{t=1}^T \left(\frac{i-1}{m} - X_t \right) A_t^i = \left[\frac{i-1}{m} - \rho_T^i \right] \bar{A}_T^i \\ \bar{e}_T^i &= \frac{1}{T} \sum_{t=1}^T \left(X_t - \frac{i}{m} \right) A_t^i = \left[\rho_T^i - \frac{i}{m} \right] \bar{A}_T^i \end{aligned}$$

where $\bar{A}_T^i = \sum A_t^i / T$. I will show that the following forecasting rule will drive both of these distances to zero:

1. If there exist an i^* such that $\bar{e}_T^{i^*} \leq 0$ and $\bar{d}_T^{i^*} \leq 0$, then forecast $\frac{2i^*-1}{2m}$.
2. Otherwise, find an i^* such that $\bar{d}_T^{i^*} > 0$ and $\bar{e}_T^{i^*-1} < 0$ then randomly forecast either $\frac{2i^*-1}{2m}$ or $\frac{2i^*+1}{2m}$ with probabilities:

$$P \left(f_{T+1} = \frac{2i^*-1}{2m} \right) = 1 - P \left(f_{T+1} = \frac{2i^*+1}{2m} \right) = \frac{\bar{d}_T^{i^*}}{\bar{d}_T^{i^*} + \bar{e}_T^{i^*-1}}$$

It is clear that an i^* can be found in step 2, since $i = 1$ always under forecasts and $i = m$ always over-forecasts.

The L-1 calibration score:

$$C_{1,T} \equiv \sum_{i=0}^m |\rho_t^i - \frac{2i+1}{m}| \bar{A}_T^i = \frac{1}{2m} + \sum_{i=1}^m \max(\bar{d}_T^i, \bar{c}_T^i)$$

so showing that all the \bar{c}_T^i and \bar{d}_T^i converge to zero, implies that $C_{1,T}$ converges to $\frac{1}{2m}$.

Theorem 1 (Foster and Vohra) *For all $\epsilon > 0$, there exists a forecasting method which is calibrated in the sense that $C_{1,T} < \epsilon$ if T is sufficiently large. In particular the above algorithm will achieve this goal if $m \geq \frac{1}{\epsilon}$.*

Consider this as a game between a statistician and nature. The statistician picks the forecast f_t and nature picks the data sequence X_t . The statistician's goal is to force all of the \bar{c}^i and \bar{d}^i to be negative (or at least approach this in the limit). Nature's goal is to keep the statistician from doing this. This set up is a game of "approachability" which was studied by Blackwell. He found a necessary and sufficient condition for a set to be approachable.

Theorem 2 (Blackwell 1956) *Let L_{ij} be a vector valued payoff taking values in R^n , where the statistician picks an i from \mathcal{I} at round i and nature picks a strategy j from \mathcal{J} at time t . Let G be a convex subset of R^n . Let $a \in R^n$ and let $c \in G$ be the closest point in G to the point a . Then G is approachable by the statistician if for all such a , there exist a weight vector w_i such that for all $j \in \mathcal{J}$,*

$$\left(\sum_{i \in \mathcal{I}} w_i L_{ij} - c \right)' (a - c) \leq 0. \quad (1)$$

To prove Theorem 1, we need to translate the calibration game into a Blackwell approachability game. The set of strategies for the statistician, \mathcal{I} ,

is the set of the m different forecasts. The set of strategies for nature, \mathcal{J} , is the set $\{0, 1\}$. Define

$$\begin{aligned}c_X^i &= (X - \frac{i}{m})A^i \\d_X^i &= (\frac{i-1}{m} - X)A^i\end{aligned}$$

The vector loss is the vector of all the (d^i, ϵ^i) 's, in otherwords, it is a point in R^{2m} . The goal set $G \subset R^{2m}$ is $G = \{x \in R^{2m} | (\forall k)x_k \leq 0\}$. Let $\bar{\epsilon}_T^i = \frac{1}{T} \sum_{t=1}^T \epsilon_{X_t}^i$ and $\bar{d}_T^i = \frac{1}{T} \sum_{t=1}^T d_{X_t}^i$. The $(d_X^i, \epsilon_X^i)_i$ will be our L_{ij} in the Blackwell game, and $(\bar{d}^i, \bar{\epsilon}^i)_i$ will be the point c . The closest point in G to the current average $a = (\bar{d}^i, \bar{\epsilon}^i)_{i \in \mathcal{I}}$ is

$$c = \left((\bar{d}^i)^-, (\bar{\epsilon}^i)^- \right)_{i \in \mathcal{I}}.$$

where we have defined the positive and negative parts as $x^+ = \max(0, x)$ and $x^- = \min(0, x)$. The weight vector w is the vector of probability of forecasting i/k .

Proof: (Hart 1996) Now to check equation (1). Writing it in terms of d^i 's and ϵ^i 's equation (1) is:

$$\sum_{i=1}^m \left((w^i d^i - (\bar{d}^i)^-)(\bar{d}^i - (\bar{d}^i)^-) + (w^i \epsilon^i - (\bar{\epsilon}^i)^-)(\bar{\epsilon}^i - (\bar{\epsilon}^i)^-) \right) \leq 0$$

from $x - x^- = x^+$ equation (1) is equivalent to

$$\sum_{i=1}^m \left((w^i d^i - (\bar{d}^i)^-)(\bar{d}^i)^+ + (w^i \epsilon^i - (\bar{\epsilon}^i)^-)(\bar{\epsilon}^i)^+ \right) \leq 0$$

Since, $(x^-)(x^+) = 0$, it is sufficient to show:

$$\sum_{i=1}^m w^i (\epsilon^i (\bar{\epsilon}^i)^+ + d^i (\bar{d}^i)^+) \leq 0.$$

If step 1. of the algorithm is used the weight vector is just $w^{i^*} = 1$ if i^* is the forecast chosen and zero otherwise. So $w^i \neq 0$ only when both $(\bar{d}^i)^+$ and $(\bar{c}^i)^+$ are zero, so the entire sum is zero.

If step 2. is used, the non-zero terms are w^{i^*} and w^{i^*-1} . But, $(\bar{c}^{i^*})^+$ is zero and $(\bar{d}^{i^*-1})^+$ is zero. So, it is sufficient to show:

$$w^{i^*} d^{i^*} (\bar{d}^{i^*})^+ + w^{i^*-1} c^{i^*-1} (\bar{c}^{i^*-1})^+ \leq 0$$

But, $d^{i^*} = -c^{i^*-1}$, so it is sufficient to show:

$$w^{i^*} (\bar{d}^{i^*})^+ - w^{i^*-1} (\bar{c}^{i^*-1})^+ \leq 0$$

But, this follows (with equality) from the definition of our probabilities. \square

References

- Blackwell, D. (1956). "An analog of the minimax theorem for vector pay-offs." *Pacific Journal of Mathematics*, **6**, 1-8.
- Foster, D. and R. Vohra. (1991) "Asymptotic Calibration." technical report, Graduate School of Business, University of Chicago, Chicago IL.
- Foster, D. and R. Vohra. (1997) "Asymptotic Calibration." manuscript.
- Fudenberg, D. and D. Levine. (1997) "An easier way to calibrate." manuscript, 1995.
- Hart, S., personal communication, 1995.
- Hart, S. and A. Mas-Colell. (1997) "A simple adaptive procedure leading to correlated equilibrium."