

Wang, Jingnan; Korn, Ralf

Article

Numerical algorithms for reflected anticipated backward stochastic differential equations with two obstacles and default risk

Risks

Provided in Cooperation with:

MDPI – Multidisciplinary Digital Publishing Institute, Basel

Suggested Citation: Wang, Jingnan; Korn, Ralf (2020) : Numerical algorithms for reflected anticipated backward stochastic differential equations with two obstacles and default risk, Risks, ISSN 2227-9091, MDPI, Basel, Vol. 8, Iss. 3, pp. 1-30, <https://doi.org/10.3390/risks8030072>

This Version is available at:

<https://hdl.handle.net/10419/258025>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.



<https://creativecommons.org/licenses/by/4.0/>

Article

Numerical Algorithms for Reflected Anticipated Backward Stochastic Differential Equations with Two Obstacles and Default Risk

Jingnan Wang ^{1,*}  and Ralf Korn ² 

¹ Department of Mathematics, University of Kaiserslautern, 67663 Kaiserslautern, Germany

² Department of Mathematics, University of Kaiserslautern, Fraunhofer ITWM, 67663 Kaiserslautern, Germany; korn@mathematik.uni-kl.de

* Correspondence: wang@mathematik.uni-kl.de; Tel.: +49-0631-205-2595

Received: 27 April 2020; Accepted: 12 June 2020; Published: 1 July 2020



Abstract: We study numerical algorithms for reflected anticipated backward stochastic differential equations (RABSDEs) driven by a Brownian motion and a mutually independent martingale in a defaultable setting. The generator of a RABSDE includes the present and future values of the solution. We introduce two main algorithms, a discrete penalization scheme and a discrete reflected scheme basing on a random walk approximation of the Brownian motion as well as a discrete approximation of the default martingale, and we study these two methods in both the implicit and explicit versions respectively. We give the convergence results of the algorithms, provide a numerical example and an application in American game options in order to illustrate the performance of the algorithms.

Keywords: numerical algorithm; reflected anticipated backward stochastic differential equations; discrete penalization scheme; discrete reflected scheme

1. Introduction

The backward stochastic differential equation (BSDE) theory plays a significant role in financial modeling. Given a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, where $B := (B_t)_{t \geq 0}$ is a d -dimensional standard Brownian motion, $\mathcal{F} := (\mathcal{F}_t)_{t \geq 0}$ is the associated natural filtration of B , $\mathcal{F}_t = \sigma(B_s; 0 \leq s \leq t)$, and \mathcal{F}_0 contains all P -null sets of \mathcal{F} . We first consider the following form of BSDE with the generator f and the terminal value ξ :

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dB_s, \quad t \in [0, T]. \quad (1)$$

The setting of this problem is to find a pair of \mathcal{F}_t -adapted processes $(Y, Z) \in \mathcal{S}_{\mathcal{F}}^2(0, T; \mathbb{R}) \times \mathcal{L}_{\mathcal{F}}^2(0, T; \mathbb{R}^d)$ satisfying BSDE (1).

Linear BSDE was first introduced by Bismut (1973), when he studied maximum principle in stochastic optimal control. Pardoux and Peng (1990) studied the general nonlinear BSDEs under a smooth square integrability assumptions on the coefficient and the terminal value, and a Lipschitz condition for the generator f . Duffie and Epstein (1992) independently used a class of BSDEs to describe the stochastic differential utility function theory in uncertain economic environments. Tang and Li (1994) considered the BSDEs driven by a Brownian motion and an independent Poisson jump. Barles et al. (1997) completed the theoretical proofs of BSDE with Poisson jump. Cordoni and Di Persio (2014) studied the hedging, option pricing and insurance problems in a BSDE approach.

Cvitanic and Karatzas (1996) first studied reflected BSDEs with continuous lower obstacle and continuous upper obstacle under the smooth square integrability assumption and Lipschitz condition. A quadruple $(Y, Z, K^+, K^-) := (Y_t, Z_t, K_t^+, K_t^-)_{0 \leq t \leq T}$ is a solution RBSDE with the generator f , the terminal value ζ and the obstacles L and V :

$$\begin{cases} Y_t = \zeta + \int_t^T f(s, Y_s, Z_s) ds + (K_T^+ - K_t^+) - (K_T^- - K_t^-) - \int_t^T Z_s dB_s, & t \in [0, T]; \\ V_t \geq Y_t \geq L_t, & t \in [0, T]; \\ \int_0^T (Y_t - L_t) dK_t^+ = \int_0^T (V_t - Y_t) dK_t^- = 0. \end{cases} \quad (2)$$

where K^+ and K^- are continuous increasing processes, K^+ is to keep Y above L , while K^- is to keep Y under V in a minimal way. When $V \equiv \infty$ and $K^- \equiv 0$ (resp. $L \equiv \infty$ and $K^+ \equiv 0$), we obtain a reflected anticipated backward stochastic differential equation (RABSDE) with one lower (resp. upper) obstacle. The existence of the solution of RBSDE with two obstacles can be obtained under one of the following assumptions: (1) one of the obstacles L and V are regular (see e.g., Cvitanic and Karatzas 1996; Hamadène et al. 1997); (2) Mokobodski's condition (see e.g., Hamadène and Lepeltier 2000; Lepeltier and Xu 2007), which means the existence of a difference of non-negative super-martingales between obstacles L and V . However, both of them have disadvantages, assumption (1) is somewhat restrictive, (2) is difficult to verify in practice. In this paper, we use the Assumption 5 for the obstacles.

Peng and Yang (2009) studied a new type of BSDE, anticipated BSDE (ABSDE) whose generator includes the values of both the present and the future,

$$\begin{cases} Y_t = \zeta_T + \int_t^T f(s, Y_s, \mathbb{E}^{\mathcal{G}_s}[Y_{s+\delta}], Z_s, \mathbb{E}^{\mathcal{G}_s}[Z_{s+\delta}]) ds - \int_t^T Z_s dB_s, & t \in [0, T]; \\ Y_t = \zeta_t, & t \in (T, T + T^\delta]; \\ Z_t = \alpha_t, & t \in (T, T + T^\delta]. \end{cases} \quad (3)$$

under the smooth square integrability assumption of the anticipated processes ζ and α , and Lipschitz condition of the generator f . Peng and Yang (2009) gave the existence and uniqueness theorem and the comparison theorem of anticipated BSDE (3). Øksendal et al. (2011) extended this topic to ABSDEs driven by a Brownian motion and an independent Poisson random measure. Jeanblanc et al. (2017) studied ABSDEs driven by a Brownian motion and a single jump process.

Default risk is the risk that an investor suffers a loss due to the inability of getting back the initial investment, it arises from a borrower failing to make required payments. This loss may be complete or partial (more see Kusuoka 1999). Peng and Xu (2009) introduced BSDE with default risk and gave the relative existence and uniqueness theorem and comparison theorem. Jiao and Pham (2011) studied the optimal investment with counterparty risk. Jiao et al. (2013) continued the research on the optimal investment under multiple default risk through a BSDE approach. Cordoni and Di Persio (2016) studied the BSDE with delayed generator in a defaultable setting. In this paper, we focus on the study of reflected anticipated BSDE with two obstacles and default risk.

For the numerical methods of BSDEs, Peng and Xu (2011) studied numerical algorithms for BSDEs driven by Brownian motion. Xu (2011) introduced a discrete penalization scheme and a discrete reflected scheme for RBSDE with two obstacles. Later Dumitrescu and Labart (2016) extended to RBSDE with two obstacles driven by Brownian motion and an independent compensated Poisson process. Lin and Yang (2014) studied the discrete BSDE with random terminal horizon.

The paper is organized as follows, we first introduce the basics of the defaultable model in Section 1.1 and the reflected anticipated BSDE (4) with two obstacles and default risk in Section 1.3. Section 2 illustrates the discrete time framework. We study the implicit and the explicit methods of two discrete schemes, i.e., the discrete penalization scheme in Section 3 and the discrete reflected scheme in Section 4. Section 5 completes the convergence results of the numerical algorithms which were provided in the previous sections. In Section 6, we illustrate the performance of the algorithms

by a simulation example and an application in American game options in the defaultable setting. The proofs of the convergence results in Section 5 can be found in the Appendix A.

1.1. Basics of the Defaultable Model

Let $\tau = \{\tau_i; i = 1, \dots, k\}$ be k non-negative random variables on a probability space $(\Omega, \mathcal{G}, \mathbb{P})$ satisfying

$$\mathbb{P}(\tau_i > 0) = 1; \quad \mathbb{P}(\tau_i > t) > 0, \forall t > 0; \quad \mathbb{P}(\tau_i = \tau_j) = 0, i \neq j.$$

For each $i = 1, \dots, k$, we define a right-continuous default process $H^i := (H_t^i)_{t \geq 0}$, where $H_t^i := 1_{\{\tau_i \leq t\}}$, denote by $\mathcal{H}^i := (\mathcal{H}_t^i)_{t \geq 0}$ the associated filtration $\mathcal{H}_t^i := \sigma(H_s^i; 0 \leq s \leq t)$. We assume that \mathcal{F}_0 is trivial (it follows that \mathcal{G}_0 is trivial as well). For a fixed terminal time $T \geq 0$, there are two kinds of information: one is from the asset prices, denoted by $\mathcal{F} := (\mathcal{F}_t)_{0 \leq t \leq T}$; the other is from the default times $\{\tau_i; i = 1, \dots, k\}$, denoted by $\{\mathcal{H}^i; i = 1, \dots, k\}$.

The enlarged filtration considered is denoted by $\mathcal{G} := (\mathcal{G}_t)_{0 \leq t \leq T}$, where $\mathcal{G}_t = \mathcal{F}_t \vee \mathcal{H}_t^1 \vee \dots \vee \mathcal{H}_t^k$. Generally, a \mathcal{G} -stopping time is not necessarily a \mathcal{F} -stopping time. Let $G := G_t$, where $G_t = \mathbb{P}(\tau > t | \mathcal{F}_t)$, i.e., $G_t^i = \mathbb{P}(\tau_i > t | \mathcal{F}_t)$, for each $i = 1, \dots, k$. In the following, G^i is assumed to be continuous, then the random default time τ_i is totally inaccessible \mathcal{G} -stopping time. The processes H^i ($i = 1, \dots, k$) are obviously \mathcal{G} -adapted, but they are not necessarily \mathcal{F} -adapted. We need the following assumptions (see Kusuoka 1999; Bielecki et al. 2007):

Assumption 1. There exist \mathcal{F} -adapted processes $\gamma^i \geq 0$ ($i = 1, \dots, k$) such that

$$M_t^i = H_t^i - \int_0^t 1_{\{\tau_i > s\}} \gamma_s^i ds$$

are \mathcal{G} -martingales under \mathbb{P} . γ^i is the \mathcal{G} -intensity of the default time τ_i :

$$\gamma_t^i = \lim_{\Delta \rightarrow 0} \frac{\mathbb{P}(t < \tau_i \leq t + \Delta | \mathcal{F}_t)}{\Delta \mathbb{P}(\tau_i > t | \mathcal{F}_t)}, \quad t \in [0, T];$$

Assumption 2. Every \mathcal{F} -local martingale is a \mathcal{G} -local martingale.

1.2. Basic Notions

- $\mathcal{L}^2(\mathcal{G}_T; \mathbb{R}) := \{\varphi \in \mathbb{R} \mid \varphi \text{ is a } \mathcal{G}_T\text{-measurable random variable and } \mathbb{E}|\varphi|^2 < \infty\};$
- $\mathcal{L}_{\mathcal{G}}^2(0, t; \mathbb{R}^d) := \{\varphi : \Omega \times [0, t] \rightarrow \mathbb{R}^d \mid \varphi_t \text{ is } \mathcal{G}_t\text{-progressively measurable and } \mathbb{E} \int_0^t |\varphi_s|^2 ds < \infty\};$
- $\mathcal{S}_{\mathcal{G}}^2(0, t; \mathbb{R}) := \{\varphi : \Omega \times [0, t] \rightarrow \mathbb{R} \mid \varphi_t \text{ is } \mathcal{G}_t\text{-progressively measurable rcll process and } \mathbb{E} \left[\sup_{0 \leq s \leq t} |\varphi_s|^2 \right] < \infty\};$
- $\mathcal{L}_{\mathcal{G}}^{2, \tau}(0, t; \mathbb{R}^k) := \{\varphi : \Omega \times [0, t] \rightarrow \mathbb{R}^k \mid \varphi_t \text{ is } \mathcal{G}_t\text{-progressively measurable and satisfies } \mathbb{E} \left[\int_0^t |\varphi_s|^p 1_{\{\tau > s\}} \gamma_s ds \right] = \mathbb{E} \left[\int_0^t \sum_{i=1}^k |\varphi_{i,s}|^2 1_{\{\tau_i > s\}} \gamma_s ds \right] < \infty\};$
- $\mathcal{A}_{\mathcal{G}}^2(0, T; \mathbb{R}) := \{K : \Omega \times [0, T] \rightarrow \mathbb{R} \mid K_t \text{ is a } \mathcal{G}_t\text{-adapted rcll increasing process and } K_0 = 0, K_T \in \mathcal{L}^p(\mathcal{G}_T; \mathbb{R})\};$

1.3. Reflected Anticipated BSDEs with Two Obstacles and Default Risk

Consider the RABSDE below with two obstacles and default risk with coefficient (f, ξ, δ, L, V) . $(Y, Z, U, K^+, K^-) := (Y_t, Z_t, U_t, K_t^+, K_t^-)_{0 \leq t \leq T + \delta}$ is a solution for RABSDE with the generator f , the terminal value ξ_T , the anticipated processes ξ , the anticipated time δ ($\delta > 0$ is a constant), and the obstacles L and V , such that

$$\left\{ \begin{array}{l} Y_t = \xi_T + \int_t^T f(s, Y_s, \mathbb{E}^{G_s}[Y_{s+\delta}], Z_s, U_s) ds + (K_T^+ - K_t^+) - (K_T^- - K_t^-) \\ \quad - \int_t^T Z_s dB_s - \int_t^T U_s dM_s, \quad t \in [0, T]; \\ Y_t \geq Y_t \geq L_t, \quad t \in [0, T]; \\ Y_t = \xi_t, \quad t \in (T, T + \delta]; \\ \int_0^T (Y_t - L_t) dK_t^+ = \int_0^T (V_t - Y_t) dK_t^- = 0, \end{array} \right. \tag{4}$$

where $Y \in \mathcal{S}_{\mathcal{G}}^2(0, T + \delta; \mathbb{R})$, $Z \in \mathcal{L}_{\mathcal{G}}^2(0, T; \mathbb{R}^d)$, $U \in \mathcal{L}_{\mathcal{G}}^{2,\tau}(0, T; \mathbb{R}^k)$, $K^\pm \in \mathcal{A}_{\mathcal{G}}^2(0, T; \mathbb{R})$. We further state the following assumptions for RABSDE (4):

Assumption 3. The anticipated process $\xi \in \mathcal{L}_{\mathcal{G}}^2(T, T + \delta; \mathbb{R}^d)$, $\alpha \in \mathcal{L}_{\mathcal{G}}^2(T, T + T^\delta; \mathbb{R}^d)$, here ξ is a given process, and ξ_T is the terminal value;

Assumption 4. The generator $f(w, t, y, \bar{y}_r, z, \bar{z}_r, u, \bar{u}_r) : \Omega \times [0, T + T^\delta] \times \mathbb{R} \times \mathcal{S}_{\mathcal{G}}^2(t, T + T^\delta; \mathbb{R}) \times \mathbb{R}^d \times \mathcal{L}_{\mathcal{G}}^2(t, T + T^\delta; \mathbb{R}^d) \times \mathbb{R}^k \rightarrow \mathcal{L}_{\mathcal{G}}^{2,\tau}(t, T + T^\delta; \mathbb{R}^k)$ satisfies:

- (a) $f(\cdot, 0, 0, 0, 0) \in \mathcal{L}_{\mathcal{G}}^2(0, T + \delta; \mathbb{R})$;
- (b) **Lipschitz condition:** for any $t \in [0, T]$, $r \in [t, T + \delta]$, $y, y' \in \mathbb{R}$, $z, z' \in \mathbb{R}^d$, $u, u' \in \mathbb{R}^k$, $\bar{y}, \bar{y}' \in \mathcal{L}_{\mathcal{G}}^2(t, T + \delta; \mathbb{R})$, there exists a constant $L \geq 0$ such that

$$|f(t, y, \bar{y}_r, z, u) - f(t, y', \bar{y}'_r, z', u')| \leq L \left(|y - y'| + \mathbb{E}^{G_t} |\bar{y}_r - \bar{y}'_r| + |z - z'| + |u - u'| 1_{\{\tau > t\}} \sqrt{\gamma_t} \right);$$

- (c) for any $t \in [0, T]$, $r \in [t, T + \delta]$, $y, y' \in \mathbb{R}$, $z, z' \in \mathbb{R}^d$, $u, u' \in \mathbb{R}^k$, $\bar{y}, \bar{y}' \in \mathcal{L}_{\mathcal{G}}^2(t, T + T^\delta; \mathbb{R})$, the following holds:

$$\frac{f(t, y, \bar{y}_r, z, \bar{u}^{i-1}) - f(t, y, \bar{y}_r, z, \bar{u}^i)}{(u^i - \bar{u}^i) 1_{\{\tau^i > t\}} \gamma_t^i} > -1,$$

where $\bar{u}^i = (\bar{u}^1, \bar{u}^2, \dots, \bar{u}^i, u^{i+1}, \dots, u^k)$, u^i is the i -th element of u .

Assumption 5. The obstacle processes satisfy $L, V \in \mathcal{S}_{\mathcal{G}}^2(0, T + \delta; \mathbb{R})$:

- (a) for any $t \in [0, T]$, $V_T \geq \xi \geq L_T$, L and V are separated, i.e., $V_t > L_t$, $\mathbb{P} - a.s.$;
- (b) L and V are rcll and their jumping times are totally inaccessible and satisfy

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} (L_t^+)^2 \right] < \infty, \quad \mathbb{E} \left[\sup_{0 \leq t \leq T} (V_t^-)^2 \right] < \infty;$$

- (c) there exists a process of the following form:

$$X_t = X_0 - \int_0^t \sigma_s^{(1)} dB_s - \int_0^t \sigma_s^{(2)} dM_s + A_t^+ - A_t^-,$$

where $X_T = \xi_T$, $\sigma^{(1)} \in \mathcal{L}_{\mathcal{G}}^2(0, T; \mathbb{R}^d)$, $\sigma^{(2)} \in \mathcal{L}_{\mathcal{G}}^{2,\tau}(0, T; \mathbb{R}^k)$, A^+ and A^- are \mathcal{G} -adapted increasing processes, $\mathbb{E}[|A_T^+|^2 + |A_T^-|^2] < \infty$, such that

$$V_t \geq X_t \geq L_t, \quad t \in [0, T], \quad \mathbb{P} - a.s.$$

2. Discrete Time Framework

In order to discretize $[0, T]$, for $n \in \mathbb{N}$, we introduce $\Delta^n := \frac{T}{n}$ and an equidistant time grid $(t_i)_{i=0,1,\dots,n,\dots,n^\delta}$ with step size Δ^n , where $t_i := i\Delta^n$, $n^\delta = n + \lceil \frac{\delta}{\Delta^n} \rceil$.

2.1. Random Walk Approximation of the Brownian Motion

We use a random walk to approximate the 1-dimensional standard Brownian motion:

$$B_0^n = 0; \quad B_t^n := \sqrt{\Delta^n} \sum_{j=1}^{[t/\Delta^n]} \epsilon_j^n, \quad t \in (0, T];$$

$$\Delta B_i^n := B_i^n - B_{i-1}^n = \sqrt{\Delta^n} \epsilon_i^n, \quad i \in [1, n^\delta],$$

where $(\epsilon_i^n)_{i=1, \dots, n}$ is a $\{-1, 1\}$ -value i.i.d. Bernoulli sequence with $\mathbb{P}(\epsilon_i^n = 1) = \mathbb{P}(\epsilon_i^n = -1) = \frac{1}{2}$. Denote $\mathcal{F}_i^n = \sigma\{\epsilon_1^n, \dots, \epsilon_i^n\}$, for any $i \in [1, n^\delta]$. By Donsker's invariance principle and the Skorokhod representation theorem, there exists a probability space, such that $\sup_{0 \leq t \leq T+\delta} |B_t^n - B_t| \rightarrow 0$, in $\mathcal{L}^2(\mathcal{G}_{T+\delta})$, as $n \rightarrow \infty$.

2.2. Approximation of the Defaultable Model

We consider a defaultable model of a single uniformly distributed random default time $\tau \in (0, T]$. We define the discrete default process $h_i^n = h_{t_i}^n = 1_{\{\tau \leq t_i\}}$ ($i \in [1, n]$). Particularly, when $i \in [n+1, n^\delta]$, $h_i^n = 1$ (since default case already happened). We have the conditional expectations of h_i^n in \mathcal{G}_{i-1}^n :

$$\mathbb{E}[h_i^n = 1 | h_{i-1}^n = 1] = \mathbb{P}(\tau \leq t_i | \tau \leq t_{i-1}) = 1,$$

$$\mathbb{E}[h_i^n = 1 | h_{i-1}^n = 0] = \mathbb{P}(\tau \leq t_i | \tau > t_{i-1}) = \frac{\Delta^n}{T - t_{i-1}},$$

$$\mathbb{E}[h_i^n = 0 | h_{i-1}^n = 0] = \mathbb{P}(\tau > t_i | \tau > t_{i-1}) = \frac{T - t_i}{T - t_{i-1}}, \quad i \in [1, n].$$

We have the following approximation for the discrete martingale M_i^n directly based on the definition of the martingale M (Assumption 1):

$$M_0^n = 0; \quad M_t^n := h_{[t/\Delta^n]}^n - \Delta^n \sum_{j=1}^{[t/\Delta^n]} (1 - h_j^n) \gamma_j^n, \quad t \in (0, T];$$

$$\Delta M_i^n := h_i^n - h_{i-1}^n - \Delta^n (1 - h_i^n) \gamma_i^n, \quad i \in [1, n],$$
(5)

where the discrete intensity process $\gamma_i^n = \gamma_{t_i}^n \geq 0$ is an \mathcal{F}_i^n -adapted process. Denote $\mathbb{G} := \{\mathcal{G}_i^n; i \in [0, n^\delta]\}$, $\mathcal{G}_0^n = \{\Omega, \emptyset\}$, for $i \in [1, n]$, $\mathcal{G}_i^n = \sigma\{\epsilon_1^n, \dots, \epsilon_i^n, h_i^n\}$; for $i \in [n+1, n^\delta]$, $\mathcal{G}_i^n = \sigma\{\epsilon_1^n, \dots, \epsilon_i^n, h_n^n\}$, where h_i is independent from $\epsilon_1^n, \dots, \epsilon_i^n$. From the martingale property of M_i , we can get

$$\mathbb{E}^{\mathcal{G}_{i-1}^n} [\Delta M_i^n] = \mathbb{E}^{\mathcal{G}_{i-1}^n} [h_i^n - h_{i-1}^n - \Delta^n (1 - h_i^n) \gamma_i^n] = 0, \quad i \in [1, n],$$

therefore, the discrete intensity process has the following form (by the projection on \mathcal{F}_{i-1}^n):

$$\gamma_i^n = \frac{\mathbb{P}(t_{i-1} < \tau \leq t_i | \mathcal{F}_{i-1}^n)}{\Delta^n \mathbb{P}(\tau > t_i | \mathcal{F}_{i-1}^n)} = \frac{1}{T - t_i}, \quad i \in [1, \left\lceil \frac{\tau}{\Delta^n} \right\rceil].$$

Note that $\gamma_i^n = 0$, when $i = 0$ and $i \in \left[\left\lceil \frac{\tau}{\Delta^n} \right\rceil + 1, n\right]$. If we set $\hat{\gamma}_t^n = \gamma_{[t/\Delta^n]}^n$ ($t \in [0, T]$), then as $n \rightarrow \infty$, it follows that $\hat{\gamma}_t^n$ converges to γ_t .

2.3. Computing the Conditional Expectations

When $i \in [1, n-1]$, we use the following formula to compute the conditional expectation for the function $f : \mathbb{R}^{i+2} \rightarrow \mathbb{R}$:

$$\begin{aligned} & \mathbb{E}^{\mathcal{G}_i^n} [f(\epsilon_1^n, \dots, \epsilon_{i+1}^n, h_{i+1}^n)] \\ &= \frac{1}{2} \left(f(\epsilon_1^n, \dots, \epsilon_i^n, 1, 1) \Big|_{\epsilon_{i+1}^n=1, h_i^n=1} + f(\epsilon_1^n, \dots, \epsilon_i^n, -1, 1) \Big|_{\epsilon_{i+1}^n=-1, h_i^n=1} \right) \\ & \quad + \frac{\Delta^n}{2(T-t_i)} \left(f(\epsilon_1^n, \dots, \epsilon_i^n, 1, 1) \Big|_{\epsilon_{i+1}^n=1, h_i^n=0, h_{i+1}^n=1} + f(\epsilon_1^n, \dots, \epsilon_i^n, -1, 1) \Big|_{\epsilon_{i+1}^n=-1, h_i^n=0, h_{i+1}^n=1} \right) \\ & \quad + \frac{T-t_{i+1}}{2(T-t_i)} \left(f(\epsilon_1^n, \dots, \epsilon_i^n, 1, 0) \Big|_{\epsilon_{i+1}^n=1, h_i^n=0, h_{i+1}^n=0} + f(\epsilon_1^n, \dots, \epsilon_i^n, -1, 0) \Big|_{\epsilon_{i+1}^n=-1, h_i^n=0, h_{i+1}^n=0} \right). \end{aligned}$$

When $i \in [n, n^\delta]$, we have the following conditional expectation for the function $f : \mathbb{R}^{i+2} \rightarrow \mathbb{R}$:

$$\mathbb{E}^{\mathcal{G}_i^n} [f(\epsilon_1^n, \dots, \epsilon_{i+1}^n, h_i^n)] = \frac{1}{2} f(\epsilon_1^n, \dots, \epsilon_i^n, 1, 1) \Big|_{\epsilon_{i+1}^n=1} + \frac{1}{2} f(\epsilon_1^n, \dots, \epsilon_i^n, -1, 1) \Big|_{\epsilon_{i+1}^n=-1}.$$

2.4. Approximations of the Anticipated Processes and the Generator

Consider the approximation ζ_n^n of the terminal value ζ , we have the following assumption:

Assumption 6. $(\zeta_i^n)_{i \in [n, n^\delta]}$ is \mathcal{G}_i^n -measurable, $\Psi : \{1, -1\}^n \rightarrow \mathbb{R}$ is a real analytic function, such that

$$\zeta_i^n = \Psi(\epsilon_1^n, \dots, \epsilon_i^n, h_i^n), \quad i \in [n, n^\delta],$$

particularly, the terminal value $\zeta_n^n = \Psi(\epsilon_1^n, \dots, \epsilon_n^n, h_i^n)$ is \mathcal{G}_n^n -measurable.

For the approximation $(f^n(t_i, y, \bar{y}, z, u))_{i \in [0, n]}$ of the generator f :

Assumption 7. for any $i \in [0, n]$, $f^n(t_i, y, \bar{y}, z, u)$ is \mathcal{G}_i^n -adapted, and satisfies:

(a) there exists a constant $C > 0$, such that for all $n > 1 + 2L + 4L^2$,

$$\mathbb{E} \left[\Delta^n \sum_{i=0}^{n-1} |f^n(\cdot, 0, 0, 0, 0)|^2 \right] < C.$$

(b) for any $i \in [0, n - 1]$, $y, y' \in \mathbb{R}, z, z' \in \mathbb{R}, u, u' \in \mathbb{R}, \bar{y}, \bar{y}' \in \mathcal{S}_G^2(t, T + \delta; \mathbb{R})$, there exists a constant $L \geq 0$, such that

$$|f^n(t_i, y, \bar{y}_i, z, u) - f^n(t_i, y', \bar{y}'_i, z', u')| \leq L \left(|y - y'| + \mathbb{E}^{\mathcal{G}_t} |\bar{y}_i - \bar{y}'_i| + |z - z'| + |u - u'| 1_{\{\tau > t_i\}} \sqrt{\gamma_i} \right),$$

where $\bar{y}_i = \mathbb{E}^{\mathcal{G}_i^n} [y_{\bar{i}}], \bar{i} = i + \lfloor \frac{\delta}{\Delta^n} \rfloor$.

As $n \rightarrow \infty$, $f^n(\lfloor \frac{t}{\Delta^n} \rfloor, y, \bar{y}, z, u)$ converges to $f(t, y, \bar{y}, z, u)$ in $\mathcal{S}_G^2(0, T + \delta; \mathbb{R})$.

2.5. Approximation of the Obstacles

$(L_i^n)_{i \in [0, n]}$ and $(V_i^n)_{i \in [0, n]}$ are the discrete versions of L and V , by Assumption 5, we can have the following approximations:

$$\begin{aligned} L_i^n &= L_0 + \Delta^n \sum_{j=0}^{i-1} l_j^{(1)} + \sum_{j=0}^{i-1} l_j^{(2)} \Delta B_{j+1}^n + \sum_{j=0}^{i-1} l_j^{(3)} \Delta M_{j+1}^n; \\ V_i^n &= V_0 + \Delta^n \sum_{j=0}^{i-1} v_j^{(1)} + \sum_{j=0}^{i-1} v_j^{(2)} \Delta B_{j+1}^n + \sum_{j=0}^{i-1} v_j^{(3)} \Delta M_{j+1}^n, \end{aligned}$$

where $l_j^{(k)} = l_{t_j}^{(k)}, v_j^{(k)} = v_{t_j}^{(k)}$ ($k = 1, 2, 3$). By the Burkholder–Davis–Gundy inequality, it follows

$$V_i^n \geq L_i^n, \quad \sup_n \mathbb{E} \left[\sup_i (L_i^n)^{+2} + \sup_i (V_i^n)^{-2} \right] < \infty.$$

We introduce the discrete version of Assumption 5 (c):

Assumption 8. *There exists a process X_i^n with the following form:*

$$X_i^n = X_0^n - \sum_{j=0}^{i-1} \sigma_j^{(1)} \Delta B_{i+1}^n - \sum_{j=0}^{i-1} \sigma_j^{(2)} \Delta M_{i+1}^n + A_i^{+n} - A_i^{-n},$$

where A_i^{+n} and A_i^{-n} are \mathcal{G}_i^n -adapted increasing processes, $\mathbb{E} [|A_n^{+n}|^2 + |A_n^{-n}|^2] < \infty$, such that

$$L_i^n \leq X_i^n \leq V_i^n, \quad i \in [0, n].$$

We introduce two numerical algorithms below, discrete penalization scheme in Section 3 and discrete reflected scheme in Section 4. For each scheme, we study the implicit and explicit versions.

3. Discrete Penalization Scheme

We first use the methodology of penalization for the discrete scheme below. [El Karoui et al. \(1997a\)](#) proved the existence of RBSDE with one obstacle under a smooth square integrability assumption and Lipschitz condition through penalization method. [Lepeltier and Martín \(2004\)](#) used the similar penalization method to prove the existence theorem of RBSDE with two obstacles and Poisson jump. Similarly to Lemma 4.3.1 in [Wang \(2020\)](#), we consider the following special case of the penalized ABSDE for RABSDE (2):

$$\begin{aligned} -dY_t^p &= f^n(t, Y_t^p, \mathbb{E}^{\mathcal{G}_t}[Y_{t+\delta}^p], Z_t^p, U_t^p)dt + dK_t^{+p} - dK_t^{-p} - Z_t^p dB_t - U_t^p dM_t, \quad t \in [0, T], \\ Y_T^p &= \xi_T, \quad t \in [T, T + \delta], \end{aligned} \tag{6}$$

where

$$K_t^{+p} = p \int_t^T (Y_s^p - L_s)^- ds, \quad K_t^{-p} = p \int_t^T (Y_s^p - V_s)^+ ds.$$

By the existence and uniqueness theorem for ABSDEs with default risk (Theorem 4.3.3 in [Wang 2020](#)), there exists the unique solution for this penalized ABSDE (6). We will give the convergence of penalized ABSDE (6) to RABSDE (2) in Theorem 1 below.

3.1. Implicit Discrete Penalization Scheme

We first introduce the implicit discrete penalization scheme. In this scheme, p represents the penalization parameter. In practice, we can choose p which is independent of n and much larger than n , this will be illustrated in the simulation Section 6.

$$\left\{ \begin{aligned} y_i^{p,n} &= y_{i+1}^{p,n} + f^n(t_i, y_i^{p,n}, \bar{y}_i^{p,n}, z_i^{p,n}, u_i^{p,n})\Delta^n + k_i^{+p,n} - k_i^{-p,n} \\ &\quad - z_i^{p,n} \Delta B_{i+1}^n - u_i^{p,n} \Delta M_{i+1}^n, \quad i \in [0, n-1]; \\ k_i^{+p,n} &= p\Delta^n (y_i^{p,n} - L_i^n)^-, \quad i \in [0, n-1]; \\ k_i^{-p,n} &= p\Delta^n (y_i^{p,n} - V_i^n)^+, \quad i \in [0, n-1]; \\ y_i^{p,n} &= \xi_i^n, \quad i \in [n, n^\delta], \end{aligned} \right. \tag{7}$$

where $\bar{y}_i^{p,n} = \mathbb{E}^{\mathcal{G}_i^n}[y_{\bar{i}}^{p,n}]$, $\bar{i} = i + \lceil \frac{\delta}{\Delta^n} \rceil$.

For the theoretical convergence results in Section 5, we first prove the convergence (Theorem 2) of implicit discrete penalization scheme (7) to the penalized ABSDE (6), then combining with Theorem A2, we can get the convergence of the explicit discrete penalization scheme. By Theorem A3 and Theorem 1, we can prove the convergence of the implicit discrete reflected scheme (13).

From Section 2.3, taking conditional expectation in \mathcal{G}_i^n , we can calculate $y_i^{p,n}$ as follows:

$$\mathbb{E}^{\mathcal{G}_i^n} [y_i^{p,n}] = \begin{cases} \frac{1}{2} (y_i^{p,n} |_{\epsilon_i^n=1, h_i^n=1} + y_i^{p,n} |_{\epsilon_i^n=-1, h_i^n=1}) \\ + \frac{\delta}{2(T-t_i)} (y_i^{p,n} |_{\epsilon_i^n=1, h_i^n=0, h_i^n=1} + y_i^{p,n} |_{\epsilon_i^n=-1, h_i^n=0, h_i^n=1}) \\ + \frac{T-t_i-\delta}{2(T-t_i)} (y_i^{p,n} |_{\epsilon_i^n=1, h_i^n=0, h_i^n=0} + y_i^{p,n} |_{\epsilon_i^n=-1, h_i^n=0, h_i^n=0}), & \bar{i} \in [i, n-1]; \\ \frac{1}{2} (\zeta_i^n |_{\epsilon_i^n=1} + \zeta_i^n |_{\epsilon_i^n=-1}), & \bar{i} \in [n, n^\delta]. \end{cases} \tag{8}$$

Similarly, $z_i^{p,n}$ and $u_i^{p,n}$ ($i \in [0, n-1]$) are given by

$$\begin{aligned} z_i^{p,n} &= \frac{\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^{p,n} \Delta B_{i+1}^n]}{\mathbb{E}^{\mathcal{G}_i^n} [(\Delta B_{i+1}^n)^2]} = \frac{\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^{p,n} \sqrt{\Delta^n} \epsilon_{i+1}^n]}{\mathbb{E}^{\mathcal{G}_i^n} [(\sqrt{\Delta^n} \epsilon_{i+1}^n)^2]} \\ &= \frac{1}{\sqrt{\Delta^n}} \mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^{p,n} \epsilon_{i+1}^n] = \frac{y_{i+1}^{p,n} |_{\epsilon_{i+1}=1} - y_{i+1}^{p,n} |_{\epsilon_{i+1}=-1}}{2\sqrt{\Delta^n}}; \\ u_i^{p,n} &= \frac{\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^{p,n} \Delta M_{i+1}^n]}{\mathbb{E}^{\mathcal{G}_i^n} [(\Delta M_{i+1}^n)^2]} = \frac{\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^{p,n} (h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})]}{\mathbb{E}^{\mathcal{G}_i^n} [(h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})^2]} \\ &= \frac{y_{i+1}^{p,n} \Delta^n |_{h_i=0, h_{i+1}=1} - y_{i+1}^{p,n} (T-t_{i+1}) \Delta^n \gamma_{i+1} |_{h_i=0, h_{i+1}=0}}{\Delta^n + (T-t_{i+1})(\Delta^n \gamma_{i+1})^2}. \end{aligned} \tag{9}$$

Note that $u_i^{p,n}$ only exists on $[0, [\frac{\tau}{\Delta^n}] - 1]$ (i.e., before the default event happens). By taking the conditional expectation of (7) in \mathcal{G}_i^n , it follows:

$$\left\{ \begin{aligned} y_i^{p,n} &= (\Phi^{p,n})^{-1} (\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^n]), & i \in [0, n-1]; \\ k_i^{+p,n} &= p\Delta^n (y_i^{p,n} - L_i^n)^-, & i \in [0, n-1]; \\ k_i^{-p,n} &= p\Delta^n (y_i^{p,n} - V_i^n)^+, & i \in [0, n-1]; \\ y_i^{p,n} &= \zeta_i^n, & i \in [n, n^\delta]; \\ z_i^{p,n} &= \frac{1}{\sqrt{\Delta^n}} \mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^{p,n} \epsilon_{i+1}^n], & i \in [0, n-1]; \\ u_i^{p,n} &= \frac{\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^{p,n} (h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})]}{\mathbb{E}^{\mathcal{G}_i^n} [(h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})^2]}, & i \in [0, [\frac{\tau}{\Delta^n}] - 1]. \end{aligned} \right. \tag{10}$$

where $\Phi^{p,n}(y) = y - f^n(t_i, y, \bar{y}_i^{p,n}, z_i^{p,n}, u_i^{p,n})\Delta^n - p\Delta^n(y - L_i^n)^- + p\Delta^n(y - V_i^n)^+$. For the continuous time version $(Y_t^{p,n}, Z_t^{p,n}, U_t^{p,n}, K_t^{+p,n}, K_t^{-p,n})_{0 \leq t \leq T}$:

$$\begin{aligned} Y_t^{p,n} &:= y_{[t/\Delta^n]}^{p,n}, & Z_t^{p,n} &:= z_{[t/\Delta^n]}^{p,n}, & U_t^{p,n} &:= u_{[t/\Delta^n]}^{p,n}, \\ K_t^{+p,n} &:= \sum_{i=0}^{[t/\Delta^n]} k_i^{+p,n}, & K_t^{-p,n} &:= \sum_{i=0}^{[t/\Delta^n]} k_i^{-p,n}. \end{aligned}$$

3.2. Explicit Discrete Penalization Scheme

In many cases, the inverse of mapping Φ is not easy to get directly, for example, if f is not a linear function on y . We replace $y_i^{p,n}$ in f^n by $\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^{p,n}]$ in (7), it follows

$$\left\{ \begin{array}{l} \tilde{y}_i^{p,n} = \tilde{y}_{i+1}^{p,n} + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}], \tilde{y}_i^{p,n}, \tilde{z}_i^{p,n}, \tilde{u}_i^{p,n})\Delta^n + \tilde{k}_i^{+p,n} - \tilde{k}_i^{-p,n} \\ \quad - \tilde{z}_i^{p,n} \Delta B_{i+1}^n - \tilde{u}_i^{p,n} \Delta M_{i+1}^n, \quad i \in [0, n-1]; \\ \tilde{k}_i^{+p,n} = p\Delta^n (\tilde{y}_i^{p,n} - L_i^n)^-, \quad i \in [0, n-1]; \\ \tilde{k}_i^{-p,n} = p\Delta^n (\tilde{y}_i^{p,n} - V_i^n)^+, \quad i \in [0, n-1]; \\ \tilde{y}_i^{p,n} = \xi_i^n, \quad i \in [n, n^\delta], \end{array} \right. \tag{11}$$

where $\tilde{y}_i^{p,n}$, $\tilde{z}_i^{p,n}$ and $\tilde{u}_i^{p,n}$ can be calculated as (8) and (9). By Section 2.3, we computer $\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^{p,n}]$ as follows:

$$\begin{aligned} \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}] &= \frac{1}{2} \left(\tilde{y}_{i+1}^{p,n} |_{\epsilon_{i+1}^n=1, h_i^n=1} + \tilde{y}_{i+1}^{p,n} |_{\epsilon_{i+1}^n=-1, h_i^n=1} \right) \\ &\quad + \frac{\Delta^n}{2(T-t_i)} \left(\tilde{y}_{i+1}^{p,n} |_{\epsilon_{i+1}^n=1, h_i^n=0, h_{i+1}^n=1} + \tilde{y}_{i+1}^{p,n} |_{\epsilon_{i+1}^n=-1, h_i^n=0, h_{i+1}^n=1} \right) \\ &\quad + \frac{T-t_{i+1}}{2(T-t_i)} \left(\tilde{y}_{i+1}^{p,n} |_{\epsilon_{i+1}^n=1, h_i^n=0, h_{i+1}^n=0} + \tilde{y}_{i+1}^{p,n} |_{\epsilon_{i+1}^n=-1, h_i^n=0, h_{i+1}^n=0} \right). \end{aligned}$$

By taking the conditional expectation of (11) in \mathcal{G}_i^n , we have the following explicit penalization scheme:

$$\left\{ \begin{array}{l} \tilde{y}_i^{p,n} = \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}] + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}], \tilde{y}_i^{p,n}, \tilde{z}_i^{p,n}, \tilde{u}_i^{p,n})\Delta^n + \tilde{k}_i^{+p,n} - \tilde{k}_i^{-p,n}, \quad i \in [0, n-1]; \\ \tilde{k}_i^{+p,n} = \frac{p\Delta^n}{1+p\Delta^n} \left(\mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}] + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}], \tilde{y}_i^{p,n}, \tilde{z}_i^{p,n}, \tilde{u}_i^{p,n})\Delta^n - L_i^n \right)^-, \quad i \in [0, n-1]; \\ \tilde{k}_i^{-p,n} = \frac{p\Delta^n}{1+p\Delta^n} \left(\mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}] + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}], \tilde{y}_i^{p,n}, \tilde{z}_i^{p,n}, \tilde{u}_i^{p,n})\Delta^n - V_i^n \right)^+, \quad i \in [0, n-1]; \\ \tilde{y}_i^{p,n} = \xi_i^n, \quad i \in [n, n^\delta]; \\ \tilde{z}_i^{p,n} = \frac{1}{\sqrt{\Delta^n}} \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n} \epsilon_{i+1}^n], \quad i \in [0, n-1]; \\ \tilde{u}_i^{p,n} = \frac{\mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n} (h_{i+1}^n - h_i^n - \Delta^n (1-h_{i+1}^n)\gamma_{i+1})]}{\mathbb{E}^{\mathcal{G}_i^n} [(h_{i+1}^n - h_i^n - \Delta^n (1-h_{i+1}^n)\gamma_{i+1})^2]}, \quad i \in [0, [\frac{\tau}{\Delta^n}] - 1]. \end{array} \right. \tag{12}$$

For the continuous time version $(\tilde{Y}_t^{p,n}, \tilde{Z}_t^{p,n}, \tilde{U}_t^{p,n}, \tilde{K}_t^{+p,n}, \tilde{K}_t^{-p,n})_{0 \leq t \leq T}$:

$$\begin{aligned} \tilde{Y}_t^{p,n} &:= \tilde{y}_{[t/\Delta^n]}^{p,n}, & \tilde{Z}_t^{p,n} &:= \tilde{z}_{[t/\Delta^n]}^{p,n}, & \tilde{U}_t^{p,n} &:= \tilde{u}_{[t/\Delta^n]}^{p,n}, \\ \tilde{K}_t^{+p,n} &:= \sum_{i=0}^{[t/\Delta^n]} \tilde{k}_i^{+p,n}, & \tilde{K}_t^{-p,n} &:= \sum_{i=0}^{[t/\Delta^n]} \tilde{k}_i^{-p,n}. \end{aligned}$$

Remark 1. We give the following explanations of the derivation of $\tilde{k}_i^{+p,n}$ and $\tilde{k}_i^{-p,n}$:

- If $V_i^n > \tilde{y}_i^{p,n} > L_i^n$, we can get $\tilde{k}_i^{+p,n} = \tilde{k}_i^{-p,n} = 0$;
- If $\tilde{y}_i^{p,n} \leq L_i^n$, we can get $\tilde{k}_i^{+p,n} = \frac{p\Delta^n}{1+p\Delta^n} \left(\mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}] + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}], \tilde{y}_i^{p,n}, \tilde{z}_i^{p,n}, \tilde{u}_i^{p,n})\Delta^n - L_i^n \right)^-$, $\tilde{k}_i^{-p,n} = 0$. From (12), we know that p should be much larger than n to keep $\tilde{y}_i^{p,n}$ above the lower obstacle L_i^n ;
- If $\tilde{y}_i^{p,n} \geq V_i^n$, we can get $\tilde{k}_i^{-p,n} = \frac{p\Delta^n}{1+p\Delta^n} \left(\mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}] + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^{p,n}], \tilde{y}_i^{p,n}, \tilde{z}_i^{p,n}, \tilde{u}_i^{p,n})\Delta^n - V_i^n \right)^+$, $\tilde{k}_i^{+p,n} = 0$. From (12), we know that p should be much larger than n to keep $\tilde{y}_i^{p,n}$ under the upper obstacle V_i^n .

4. Discrete Reflected Scheme

We can obtain the solution Y by reflecting between the two obstacles and get the increasing processes K^+ and K^- directly.

4.1. Implicit Discrete Reflected Scheme

We have the following implicit discrete reflected scheme,

$$\left\{ \begin{array}{l} y_i^n = y_{i+1}^n + f^n(t_i, y_i^n, \bar{y}_i^n, z_i^n, u_i^n)\Delta^n + k_i^{+n} - k_i^{-n} - z_i^n \Delta B_{i+1}^n - u_i^n \Delta M_{i+1}^n, \quad i \in [0, n-1]; \\ V_i^n \geq y_i^n \geq L_i^n, \quad i \in [0, n-1]; \\ k_i^{+n} \geq 0, \quad k_i^{-n} \geq 0, \quad k_i^{+n} k_i^{-n} = 0, \quad i \in [0, n-1]; \\ (y_i^n - L_i^n) k_i^{+n} = (y_i^n - V_i^n) k_i^{-n} = 0, \quad i \in [0, n-1]; \\ y_i^n = \zeta_i^n, \quad i \in [n, n^\delta]. \end{array} \right. \tag{13}$$

where \bar{y}_i^n, z_i^n and u_i^n can be calculated as (8) and (9). By taking conditional expectation of (13) in \mathcal{G}_i^n , it follows

$$\left\{ \begin{array}{l} y_i^n = \mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^n] + f^n(t_i, y_i^n, \bar{y}_i^n, z_i^n, u_i^n)\Delta^n + k_i^{+n} - k_i^{-n}, \quad i \in [0, n-1]; \\ V_i^n \geq y_i^n \geq L_i^n, \quad i \in [0, n-1]; \\ k_i^{+n} \geq 0, \quad k_i^{-n} \geq 0, \quad k_i^{+n} k_i^{-n} = 0, \quad i \in [0, n-1]; \\ (y_i^n - L_i^n) k_i^{+n} = (y_i^n - V_i^n) k_i^{-n} = 0, \quad i \in [0, n-1]; \\ y_i^n = \zeta_i^n, \quad i \in [n, n^\delta]; \\ z_i^n = \frac{1}{\sqrt{\Delta^n}} \mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^n \epsilon_{i+1}^n], \quad i \in [0, n-1]; \\ u_i^n = \frac{\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^n (h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})]}{\mathbb{E}^{\mathcal{G}_i^n} [(h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})^2]}, \quad i \in [0, [\frac{\tau}{\Delta^n}] - 1]. \end{array} \right. \tag{14}$$

If Δ^n is small enough, similarly to Section 4.1 in Xu (2011), (14) is equivalent to

$$\left\{ \begin{array}{l} y_i^n = \Phi^{-1} \left(\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^n] + k_i^{+n} - k_i^{-n} \right), \quad i \in [0, n-1]; \\ k_i^{+n} = \left(\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^n] + f^n(t_i, L_i^n, \bar{L}_i^n, z_i^n, u_i^n)\Delta^n - L_i^n \right)^-, \quad i \in [0, n-1]; \\ k_i^{-n} = \left(\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^n] + f^n(t_i, V_i^n, \bar{V}_i^n, z_i^n, u_i^n)\Delta^n - V_i^n \right)^+, \quad i \in [0, n-1]; \\ y_i^n = \zeta_i^n, \quad i \in [n, n^\delta]; \\ z_i^n = \frac{1}{\sqrt{\Delta^n}} \mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^n \epsilon_{i+1}^n], \quad i \in [0, n-1]; \\ u_i^n = \frac{\mathbb{E}^{\mathcal{G}_i^n} [y_{i+1}^n (h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})]}{\mathbb{E}^{\mathcal{G}_i^n} [(h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})^2]}, \quad i \in [0, [\frac{\tau}{\Delta^n}] - 1]. \end{array} \right. \tag{15}$$

here $\Phi(y) = y - f^n(t, y, \bar{y}_i^n, z_i^n, u_i^n)\Delta^n$. For the continuous time version $(Y_t^n, Z_t^n, U_t^n, K_t^{+n}, K_t^{-n})_{0 \leq t \leq T}$:

$$\begin{aligned} Y_t^n &:= y_{[t/\Delta^n]}^n, & Z_t^n &:= z_{[t/\Delta^n]}^n, & U_t^n &:= u_{[t/\Delta^n]}^n, \\ K_t^{+n} &:= \sum_{i=0}^{[t/\Delta^n]} k_i^{+n}, & K_t^{-n} &:= \sum_{i=0}^{[t/\Delta^n]} k_i^{-n}. \end{aligned}$$

4.2. Explicit Discrete Reflected Scheme

We introduce the following explicit discrete reflected scheme by replacing y_i^n in the generator f^n by $\mathbb{E}[y_{i+1}^n | \mathcal{G}_i^n]$ in (13).

$$\left\{ \begin{array}{l} \tilde{y}_i^n = \tilde{y}_{i+1}^n + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n], \tilde{y}_i^n, z_i^n, \tilde{u}_i^n) \Delta^n + \tilde{k}_i^{+n} - \tilde{k}_i^{-n} - z_i^n \Delta B_{i+1}^n \\ \quad - \tilde{u}_i^n \Delta M_{i+1}^n, \quad i \in [0, n-1]; \\ V_i^n \geq \tilde{y}_i^n \geq L_i^n, \quad i \in [0, n-1]; \\ \tilde{k}_i^{+n} \geq 0, \quad \tilde{k}_i^{-n} \geq 0, \quad \tilde{k}_i^{+n} \tilde{k}_i^{-n} = 0, \quad i \in [0, n-1]; \\ (\tilde{y}_i^n - L_i^n) \tilde{k}_i^{+n} = (\tilde{y}_i^n - V_i^n) \tilde{k}_i^{-n} = 0, \quad i \in [0, n-1]; \\ \tilde{y}_i^n = \zeta_i^n, \quad i \in [n, n^\delta]. \end{array} \right. \tag{16}$$

where \tilde{y}_i^n, z_i^n and \tilde{u}_i^n can be calculated as (8) and (9). By taking conditional expectation of (16) in \mathcal{G}_i^n :

$$\left\{ \begin{array}{l} \tilde{y}_i^n = \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n] + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n], \tilde{y}_i^n, z_i^n, \tilde{u}_i^n) \Delta^n + \tilde{k}_i^{+n} - \tilde{k}_i^{-n}, \quad i \in [0, n-1]; \\ V_i^n \geq \tilde{y}_i^n \geq L_i^n, \quad i \in [0, n-1]; \\ \tilde{k}_i^{+n} \geq 0, \quad \tilde{k}_i^{-n} \geq 0, \quad \tilde{k}_i^{+n} \tilde{k}_i^{-n} = 0, \quad i \in [0, n-1]; \\ (\tilde{y}_i^n - L_i^n) \tilde{k}_i^{+n} = (\tilde{y}_i^n - V_i^n) \tilde{k}_i^{-n} = 0, \quad i \in [0, n-1]; \\ \tilde{y}_i^n = \zeta_i^n, \quad i \in [n, n^\delta]; \\ z_i^n = \frac{1}{\sqrt{\Delta^n}} \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n \epsilon_{i+1}^n], \quad i \in [0, n-1]; \\ \tilde{u}_i^{p,n} = \frac{\mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n (h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})]}{\mathbb{E}^{\mathcal{G}_i^n} [(h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})^2]}, \quad i \in [0, [\frac{\tau}{\Delta^n}] - 1]. \end{array} \right. \tag{17}$$

Similarly to the implicit reflected case, we can obtain

$$\left\{ \begin{array}{l} \tilde{y}_i^n = \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n] + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n], \tilde{y}_i^n, z_i^n, \tilde{u}_i^n) \Delta^n + \tilde{k}_i^{+n} - \tilde{k}_i^{-n}, \quad i \in [0, n-1]; \\ \tilde{k}_i^{+n} = \left(\mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n] + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n], \tilde{y}_i^n, z_i^n, \tilde{u}_i^n) \Delta^n - L_i^n \right)^-, \quad i \in [0, n-1]; \\ \tilde{k}_i^{-n} = \left(\mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n] + f^n(t_i, \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n], \tilde{y}_i^n, z_i^n, \tilde{u}_i^n) \Delta^n - V_i^n \right)^+, \quad i \in [0, n-1]; \\ \tilde{y}_i^n = \zeta_i^n, \quad i \in [n, n^\delta]; \\ z_i^n = \frac{1}{\sqrt{\Delta^n}} \mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n \epsilon_{i+1}^n], \quad i \in [0, n-1]; \\ \tilde{u}_i^{p,n} = \frac{\mathbb{E}^{\mathcal{G}_i^n} [\tilde{y}_{i+1}^n (h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})]}{\mathbb{E}^{\mathcal{G}_i^n} [(h_{i+1}^n - h_i^n - \Delta^n (1 - h_{i+1}^n) \gamma_{i+1})^2]}, \quad i \in [0, [\frac{\tau}{\Delta^n}] - 1]. \end{array} \right. \tag{18}$$

For the continuous time version $(\tilde{Y}_t^n, \tilde{Z}_t^n, \tilde{U}_t^n, \tilde{K}_t^{+n}, \tilde{K}_t^{-n})_{0 \leq t \leq T}$:

$$\begin{aligned} \tilde{Y}_t^n &:= \tilde{y}_{\lfloor t/\Delta^n \rfloor}^n, & \tilde{Z}_t^n &:= z_{\lfloor t/\Delta^n \rfloor}^n, & \tilde{U}_t^n &:= \tilde{u}_{\lfloor t/\Delta^n \rfloor}^n, \\ \tilde{K}_t^{+n} &:= \sum_{i=0}^{\lfloor t/\Delta^n \rfloor} \tilde{k}_i^{+n}, & \tilde{K}_t^{-n} &:= \sum_{i=0}^{\lfloor t/\Delta^n \rfloor} \tilde{k}_i^{-n}. \end{aligned}$$

5. Convergence Results

We first state the convergence result from the Penalized ABSDE (19) to RABSDE (2) in Theorem 1, which is the basis of the following convergence results of the discrete schemes we have studied above. We prove the convergence (Theorem 2) from the implicit discrete penalization scheme (7) to the penalized ABSDE (6) with the help of Lemma 1. Combining with Theorem A2, we can get the convergence (Theorem 3) of the explicit discrete penalization scheme (11). By Theorem A3, Lemma 1 and Theorem 1, we can prove the convergence of the implicit discrete reflected scheme (13). By Theorem A3, Theorem A4 and Lemma A4, the convergence (Theorem 5) of the explicit penalization discrete scheme (16) then follows. The proofs of Theorem 1, Lemma 1, Theorem 2 and Theorem 4 can be seen in Appendix A.

5.1. Convergence of the Penalized ABSDE to RABSDE (2)

Theorem 1. Suppose that the anticipated process ξ , the generator f satisfy Assumption 3 and Assumption 4, $f(t, y, \bar{y}_r, z, u)$ is increasing in \bar{y} , the obstacles L and V satisfy Assumption 5. We can consider the following special case of the penalized ABSDE for RABSDE (2):

$$\left\{ \begin{array}{l} -dY_t^p = f^n(t, Y_t^p, \mathbb{E}^{\mathcal{G}_t}[Y_{t+\delta}^p], Z_t^p, U_t^p)dt + dK_t^{+p} - dK_t^{-p} - Z_t^p dB_t - U_t^p dM_t, \quad t \in [0, T]; \\ K_t^{+p} = \int_0^t p(Y_s^p - L_s)^- ds, \quad t \in [0, T]; \\ K_t^{-p} = \int_0^t p(Y_s^p - V_s)^+ ds, \quad t \in [0, T]; \\ Y_t^p = \xi_t, \quad t \in [T, T + \delta]. \end{array} \right. \quad (19)$$

Then we have the limiting process (Y, Z, U, K^+, K^-) of $(Y^p, Z^p, U^p, K^{+p}, K^{-p})$, i.e., as $p \rightarrow \infty$, $Y_t^p \rightarrow Y_t$ in $\mathcal{S}_{\mathcal{G}}^2(0, T + \delta; \mathbb{R})$, $Z_t^p \rightarrow Z_t$ weakly in $\mathcal{L}_{\mathcal{G}}^2(0, T; \mathbb{R})$, $U_t^p \rightarrow U_t$ weakly in $\mathcal{L}_{\mathcal{G}}^{2,\tau}(0, T; \mathbb{R})$, $K_t^{+p} (K_t^{-p}) \rightarrow K_t^+ (K_t^-)$ weakly in $\mathcal{A}_{\mathcal{G}}^2(0, T; \mathbb{R})$. Moreover, there exists a constant $C_{\xi, f, L, V}$ depending on $\xi, f(t, 0, 0, 0, 0)$, L and V , such that

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^p - Y_t|^2 + \int_0^T |Z_t^p - Z_t|^2 dt + \int_0^T |U_t^p - U_t|^2 \mathbf{1}_{\{\tau > t\}} \gamma_t dt + \sup_{0 \leq t \leq T} |(K_t^{+p} - K_t^+) - (K_t^{-p} - K_t^-)|^2 \right] \leq \frac{1}{\sqrt{p}} C_{\xi, f, L, V}.$$

5.2. Convergence of the Implicit Discrete Penalization Scheme

We first introduce the following lemma to prove the convergence result of the penalized ABSDE (19) to the implicit penalization scheme.

Lemma 1. Under Assumption 6 and Assumption 7, $(Y_t^{p,n}, Z_t^{p,n}, U_t^{p,n})$ converges to (Y_t^p, Z_t^p, U_t^p) in the following sense:

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^{p,n} - Y_t^p|^2 + \int_0^T |Z_t^{p,n} - Z_t^p|^2 dt + \int_0^T |U_t^{p,n} - U_t^p|^2 \mathbf{1}_{\{\tau > t\}} \gamma_t dt \right] = 0, \quad (20)$$

for any $t \in [0, T]$, as $n \rightarrow \infty$, $K_t^{+p,n} - K_t^{-p,n} \rightarrow K_t^{+p} - K_t^{-p}$ in $\mathcal{L}_{\mathcal{G}}^2(0, T; \mathbb{R})$.

Theorem 2. (Convergence of the implicit discrete penalization scheme) Under Assumption 3 and Assumption 7, $(Y_t^{p,n}, Z_t^{p,n}, U_t^{p,n})$ converges to (Y_t, Z_t, U_t) in the following sense:

$$\lim_{p \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^{p,n} - Y_t|^2 + \int_0^T |Z_t^{p,n} - Z_t|^2 dt + \int_0^T |U_t^{p,n} - U_t|^2 \mathbf{1}_{\{\tau > t\}} \gamma_t dt \right] = 0, \quad (21)$$

for any $t \in [0, T]$, as $p \rightarrow \infty, n \rightarrow \infty$, $K_t^{+p,n} - K_t^{-p,n} \rightarrow K_t^+ - K_t^-$ in $\mathcal{L}_{\mathcal{G}}^2(0, T; \mathbb{R})$.

5.3. Convergence of the Explicit Discrete Penalization Scheme

By Theorem 2 and Theorem A2, we can obtain the following convergence result of the explicit penalization discrete scheme.

Theorem 3. (Convergence of the explicit discrete penalization scheme) Under Assumption 3 and Assumption 7, $(\tilde{Y}_t^{p,n}, \tilde{Z}_t^{p,n}, \tilde{U}_t^{p,n})$ converges to (Y_t, Z_t, U_t) in the following sense:

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{0 \leq t \leq T} |\tilde{Y}_t^{p,n} - Y_t|^2 + \int_0^T |\tilde{Z}_t^{p,n} - Z_t|^2 dt + \int_0^T |\tilde{U}_t^{p,n} - U_t|^2 \mathbf{1}_{\{\tau > t\}} \gamma_t dt \right] = 0, \quad (22)$$

for any $t \in [0, T]$, as $n \rightarrow \infty$, $\tilde{K}_t^{+p,n} - \tilde{K}_t^{-p,n} \rightarrow K_t^+ - K_t^-$ in $\mathcal{L}_G^2(0, T; \mathbb{R})$.

5.4. Convergence of the Implicit Discrete Reflected Scheme

Theorem 4. (Convergence of the implicit discrete reflected scheme) Under Assumption 7 and Assumption 3, (Y_t^n, Z_t^n, U_t^n) converges to (Y_t, Z_t, U_t) in the following sense:

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^n - Y_t|^2 + \int_0^T |Z_t^n - Z_t|^2 dt + \int_0^T |U_t^n - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] = 0, \tag{23}$$

and for any $t \in [0, T]$, as $n \rightarrow \infty$, $K_t^{+n} - K_t^{-n} \rightarrow K_t^+ - K_t^-$ in $\mathcal{L}_G^2(0, T; \mathbb{R})$.

5.5. Convergence of the Explicit Discrete Reflected Scheme

By Theorem A3, Theorem A4 and Lemma A4, we can get the convergence result of the explicit penalization discrete scheme.

Theorem 5. (Convergence of the explicit discrete reflected scheme) Under Assumption 3 and Assumption 7, $(\tilde{Y}_t^n, \tilde{Z}_t^n, \tilde{U}_t^n)$ converges to (Y_t, Z_t, U_t) in the following sense:

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{0 \leq t \leq T} |\tilde{Y}_t^n - Y_t|^2 + \int_0^T |\tilde{Z}_t^n - Z_t|^2 dt + \int_0^T |\tilde{U}_t^n - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] = 0, \tag{24}$$

for any $t \in [0, T]$, as $n \rightarrow \infty$, $\tilde{K}_t^{+n} - \tilde{K}_t^{-n} \rightarrow K_t^+ - K_t^-$ in $\mathcal{L}_G^2(0, T; \mathbb{R})$.

6. Numerical Calculations and Simulations

6.1. One Example of RABSDE with Two Obstacles and Default Risk

For the convenience of computation, we consider the case when the terminal time $T = 1$, the calculation begins from $y_n^n = \zeta^n$, and proceeds backward to solve $(y_i^n, z_i^n, u_i^n, k_i^{+n}, k_i^{-n})$ for $i = n - 1, n - 2, \dots, 1, 0$. We use Matlab for the simulation. We consider a simple situation: the terminal value $\zeta_T = \Phi(B_T, M_T)$ and anticipated process $\zeta_t = \Phi(B_t)$ ($t \in (T, T + \delta]$); the obstacles $L_t = \Psi_1(t, B_t, M_t)$ and $V_t = \Psi_2(t, B_t, M_t)$, where Φ, Ψ_2 and Ψ_3 are real analytic functions defined on $\mathbb{R}, [0, T] \times \mathbb{R}$ and $[0, T] \times \mathbb{R}$ respectively. We take the following example ($n = 200$, anticipated time $\delta = 0.3$):

$$\begin{aligned} f(t, y, \bar{y}, z, u) &= \left| \frac{y}{2} + \frac{\bar{y}}{2} \right| + z + u, & t \in [0, T]; \\ \Phi(B_t) &= |B_t| + M_T, & t \in [T, T + \delta]; \\ \Psi_1(t, B_t, M_t) &= |B_t| + M_t + T - t, & t \in [0, T]; \\ \Psi_2(t, B_t, M_t) &= |B_t| + M_t + 2(T - t), & t \in [0, T]; \end{aligned}$$

This example satisfies the Assumption 3, Assumption 4 and Assumption 5 in the theoretical Section 1.3. We choose the default time τ as a uniformly distributed random variable.

As the inverse for both implicit schemes in (10) and (15) is not easy to get directly, we only use explicit schemes below. We are going to illustrate the behaviors of the explicit reflected scheme by looking at the pathwise behavior for $n = 400$. Further, we will compare the explicit reflected scheme with the explicit penalization scheme for different values of the penalization parameter.

Figure 1 represents one path of the Brownian motion, Figures 2 and 3 represent one path of the Brownian motion and one path of the default martingale when the default time $\tau = 0.7$ and 0.2 respectively.




Figure 1. One path of the Brownian motion.




Figure 2. One path of the default martingale ($\tau = 0.7$).




Figure 3. One path of the default martingale ($\tau = 0.2$).

Figures 4 and 5 represent the paths of the solution \tilde{y}^n , increasing processes \tilde{K}^{+n} and \tilde{K}^{-n} in the explicit reflected scheme where the random default time $\tau = 0.7$. We can see that for all i , \tilde{y}_i^n stays between the lower obstacle L_i^n and the upper obstacle V_i^n , the increasing process \tilde{K}_i^{+n} (resp. \tilde{K}_i^{-n}) pushes \tilde{y}_i^n upward (resp. downward), and they can not increase at the same time. In this example for $n = 400$, default time $\tau = 0.7$, we can get the reflected solution $\tilde{y}_0^n = 1.2563$ from the explicit reflected scheme.




Figure 4. One path of \tilde{y}^n in the explicit reflected scheme ($\tau = 0.7$).




Figure 5. The paths of the increasing processes in the explicit reflected scheme ($\tau = 0.7$).

Figures 4 and 6 illustrate the influence of the jump on the solution \tilde{y}^n at the different random default times, the reflected solution \tilde{y}^n moves downwards after the default time (which can not be shown in Figure 7). From the approximation of the default martingale (5), M^n is larger with a larger default time.




Figure 6. One path of \tilde{y}^n in the explicit reflected scheme ($\tau = 0.2$).



Figure 7. One path of \tilde{y}^n in the explicit reflected scheme without default risk.

Table 1 and contains the comparison between the explicit reflected scheme and the explicit penalization scheme by the values of \tilde{y}_0^n and $\tilde{y}_0^{p,n}$ with respect to the parameters n and p . As n increases, the reflected solution \tilde{y}_0^n increases because of the choice of the coefficient. For fixed n , as the penalization parameter p increases, the penalization solution $\tilde{y}_0^{p,n}$ converges increasingly to the reflected solution \tilde{y}_0^n , which is obvious from the comparison theorem of BSDE with default risk. If p and n have a smaller difference (when $n = 10^3, p = 10^3$), the penalization solution $\tilde{y}_0^{p,n}$ is far from the reflected solution \tilde{y}_0^n . Hence, the penalization parameter p should be chosen as large as possible. Table 2 illustrates the comparison between the reflected solution \tilde{y}_0^n and \tilde{y}_0^{n*} . Figure 7 represents the situation without the default risk, the reflected solution \tilde{y}_0^{n*} has a larger value than in the situation when the default case happens (Figure 4).

Table 1. The values of the penalization solution $\tilde{y}_0^{p,n}$ ($\tau = 0.7$).

$\tilde{y}_0^{p,n}$	$p = 10^3$	$p = 10^4$	$p = 10^5$	$p = 10^6$
$n = 200$	1.2369	1.2394	1.2428	1.2452
$n = 400$	1.2458	1.2482	1.2496	1.2511
$n = 1000$	1.2343	1.2497	1.2527	1.2630

Table 2. The values of the reflected solution \tilde{y}_0^n ($\tau = 0.7$) and \tilde{y}_0^{n*} .

	\tilde{y}_0^n	\tilde{y}_0^{n*}
$n = 200$	1.2469	1.5451
$n = 400$	1.2563	1.5507
$n = 1000$	1.2644	1.5614

6.2. Application in American Game Options in a Defaultable Setting

6.2.1. Model Description

Hamadène (2006) studied the relation between American game options and RBSDE with two obstacles driven by Brownian motion. In our paper, we consider the case with default risk. An American game option contract with maturity T involves a broker c_1 and a trader c_2 :

- The broker c_1 has the right to cancel the contract at any time before the maturity T , while the trader c_2 has the right to early exercise the option;
- the trader c_2 pays an initial amount (the price of this option) which ensures an income L_{τ_1} from the broker c_1 , where $\tau_1 \in [0, T]$ is an \mathcal{G} -stopping time;
- the broker has the right to cancel the contract before T and needs to pay V_{τ_2} to c_2 . Here, the payment amount of the broker c_1 should be greater than his payment to the trader c_2 (if trader

decides for early exercise), i.e. $V_{\tau_2} \geq L_{\tau_2}$, $V_{\tau_2} - L_{\tau_2}$ is the premium that the broker c_1 pays for his decision of early cancellation. $\tau_2 \in [0, T]$ is an \mathcal{G} -stopping time;

- if c_1 and c_2 both decide to stop the contract at the same time τ , then the trader c_2 gets an income equal to $Q_\tau 1_{\{\tau < T\}} + \zeta 1_{\{\tau = T\}}$.

6.2.2. The Hedge for the Broker

Consider a financial market \mathcal{M} , we have a riskless asset $C_t \in \mathbb{R}$ with risk-free rate r :

$$\begin{cases} dC_t = rC_t dt, & t \in (0, T], \\ C_0 = c_0, & t = 0; \end{cases} \tag{25}$$

one risky asset $S_t \in \mathbb{R}$:

$$\begin{cases} dS_t = S_t (\mu dt + \sigma dB_t + \chi dM_t), & t \in (0, T], \\ S_0 = s_0, & t = 0, \end{cases} \tag{26}$$

where B_t is a 1-dimensional Brownian Motion, μ is the expected return, σ is the volatility, χ is the parameter related to the default risk.

Consider a self-financing portfolio $\pi \in \mathbb{R}^d$ with strategy $\pi = (\beta_s^{(1)}, \beta_s^{(2)})_{s \in [t, T]}$ trading on C and S respectively on the time interval $[t, T]$. $A^{\pi, \alpha}$ is the wealth process with the value α^A at time t , here is a non-negative \mathcal{F}_t -measurable random variable.

$$\begin{aligned} A_s^{\pi, \alpha} &= \beta_s^{(1)} C_s + \beta_s^{(2)} S_s, & s \in [t, T]; \\ A_s^{\pi, \alpha} &= \alpha^A + \int_t^s \beta_u^{(1)} dC_u + \int_t^s \beta_u^{(2)} dS_u, & s \in [t, T]; \\ \int_t^T [|\beta_u^{(1)}| + (\beta_u^{(2)} S_u)^2] du &< \infty; \end{aligned} \tag{27}$$

Let $L^{\pi, \alpha}$ be a positive local martingale with the following form:

$$\begin{cases} dL_t^{\pi, \alpha} = -L_{t-}^{\pi, \alpha} \sigma^{-1} (\mu - r) dB_t, & t \in (0, T], \\ L_0^{\pi, \alpha} = 1, & t = 0, \end{cases}$$

By Girsanov's theorem, let $\mathbb{P}^{\pi, \alpha}$ be the equivalent measure of \mathbb{P} :

$$\frac{\mathbb{P}^{\pi, \alpha}}{\mathbb{P}} \Big|_{\mathcal{G}_T} = L_T^{\pi, \alpha} = \exp \left\{ -\sigma^{-1} (\mu - r) B_T - \frac{1}{2} \left(-\sigma^{-1} (\mu - r) \right)^2 T \right\},$$

here let $\mathbb{E}^{\pi, \alpha}$ be the expectation, $B^{\pi, \alpha}$ and $M^{\pi, \alpha}$ be the Brownian motion and the default martingale under the measure $\mathbb{P}^{\pi, \alpha}$:

$$\begin{aligned} B_t^{\pi, \alpha} &:= B_t + \sigma^{-1} (\mu - r) t; \\ M_t^{\pi, \alpha} &:= M_t. \end{aligned}$$

Hence, the risky asset S_t defined in (26) can be converted into the following form under measure $\mathbb{P}^{\pi, \alpha}$:

$$\begin{cases} dS_t = S_t (r dt + \sigma dB_t^{\pi, \alpha} + \chi dM_t^{\pi, \alpha}), & t \in (0, T], \\ S_0 = s_0, & t = 0. \end{cases} \tag{28}$$

Denote by (π, θ) a hedge for the broker against the American game option after t , where π is defined in (27), $\theta \in [t, T]$ is a stopping time, satisfying

$$A_s^{\pi, \alpha} \geq R(s, \theta) := V_\theta 1_{\{\theta < s\}} + L_s 1_{\{s < \theta\}} + Q_s 1_{\{\theta = s < T\}} + \zeta 1_{\{\theta = s = T\}}, \quad s \in [t, T], \quad \mathbb{P} - a.s. \tag{29}$$

here $R(s, \theta)$ is the amount that the broker c_1 has to pay if the option is exercised by c_2 at s or canceled at time θ . Similarly to [El Karoui et al. \(1997b\)](#) and [Karatzas and Shreve \(1998\)](#), we define the value of the option at time t by J_t , where $(J_t)_{0 \leq t \leq T}$ is an rcl (right continuous with left limits) process, for any $t \in [0, T]$,

$$J_t := \text{ess inf} \left\{ \alpha^A \geq 0; \mathcal{G}_t\text{-measurable such that there exists a hedge } (\pi, \theta) \text{ after } t, \right. \\ \left. \text{where } \pi \text{ is a self-financing portfolio after } t \text{ whose value at } t \text{ is } \alpha^A. \right\} \quad (30)$$

Consider the following RBSDE with two obstacles and default risk, for any $t \in [0, T]$, there exist a stopping time θ_t , a process $(Z_s^{\pi, \alpha})_{t \leq s \leq T}$ and increasing processes $(K_s^{\pi, \alpha+})_{t \leq s \leq T}$ and $(K_s^{\pi, \alpha-})_{t \leq s \leq T}$, such that

$$\left\{ \begin{array}{l} Y_t^{\pi, \alpha} = Y_{\theta_t}^{\pi, \alpha} + \left(K_{\theta_t}^{\pi, \alpha+} - K_s^{\pi, \alpha+} \right) - \left(K_{\theta_t}^{\pi, \alpha-} - K_s^{\pi, \alpha-} \right) - \int_s^{\theta_t} Z_u^{\pi, \alpha} dB_u^{\pi, \alpha} \\ \quad - \int_s^{\theta_t} U_u^{\pi, \alpha} dM_u^{\pi, \alpha}, \quad s \in [t, \theta_t]; \\ Y_T^{\pi, \alpha} = e^{-rT} \xi; \\ e^{-rs} L_s \leq Y_s^{\pi, \alpha} \leq e^{-rs} V_s, \quad s \in [t, T]; \\ \int_t^{\theta_t} (Y_u^{\pi, \alpha} - e^{-ru} L_u) dK_u^{\pi, \alpha+} = \int_t^{\theta_t} (e^{-ru} V_u - Y_u^{\pi, \alpha}) dK_u^{\pi, \alpha-} = 0; \end{array} \right. \quad (31)$$

For any $s \in [t, T]$, $e^{rt} Y_t^{\pi, \alpha}$ is a hedge for the broker c_1 against the game option, i.e. $J_t = e^{rt} Y_t^{\pi, \alpha}$ (see [Theorem A5](#) in the Appendix). Similarly to [Proposition 4.3](#) in [Hamadène \(2006\)](#), we set

$$\theta_t^* := \inf \{ s \geq t; Y_s^{\pi, \alpha} = e^{-rs} V_s \} \wedge T = \inf \{ s \geq t; K_s^{\pi, \alpha-} > 0 \}; \\ v_t^* := \inf \{ s \geq t; Y_s^{\pi, \alpha} = e^{-rs} L_s \} \wedge T = \inf \{ s \geq t; K_s^{\pi, \alpha+} > 0 \}, \quad (32)$$

therefore, we can get

$$\hat{R}_t(v, \theta_t^*) \leq Y_t^{\pi, \alpha} = \hat{R}_t(v_t^*, \theta_t^*) \leq \hat{R}_t(v_t^*, \theta),$$

where

$$\hat{R}_t(v, \theta) := \mathbb{E}^{\pi, \alpha} \left[e^{-r\theta} V_{\theta} 1_{\{\theta < v\}} + e^{-rv} L_v 1_{\{v < \theta\}} + e^{-r\theta} Q_{\theta} 1_{\{\theta = v < T\}} + e^{-rT} \xi 1_{\{\theta = v = T\}} \mid \mathcal{G}_t \right], \quad \mathbb{P} - a.s.$$

6.2.3. Numerical Simulation

We use the same calculation method as in [Section 6.1](#), starting from $Y_n^{\pi, \alpha} = \xi$, and proceeding backward to solve $(Y_i^{\pi, \alpha}, Z_i^{\pi, \alpha}, U_i^{\pi, \alpha}, K_i^{\pi, \alpha+}, K_i^{\pi, \alpha-})$ for $i = n - 1, \dots, 1, 0$ with step size Δ^n . The forward SDEs [\(25\)](#) and [\(26\)](#) can be numerically approximated by the Euler scheme on the time grid $(t_i)_{i=0,1,\dots,n}$:

$$C_{i+1} = C_i + rC_i \Delta^n; \\ S_{i+1} = S_i + S_i (\mu \Delta^n + \sigma \Delta B_i^n + \chi \Delta M_i^n).$$

In this case, we consider parameters as below:

$$s_0 = 1.5, \quad T = 1, \quad r = 1.1, \quad \mu = 1.5, \quad \sigma = 0.5, \quad \chi = 0.2, \\ L_t = (S_t - 1)^+, \quad V_t = 2(S_t - 1)^+, \quad \xi = 1.2(S_T - 1)^+,$$

In the case $n = 400$, [Figure 8](#) represents one path of the Brownian motion, [Figures 9](#) and [10](#) represent the paths of the solution $Y^{\pi, \alpha}$, increasing processes $K^{\pi, \alpha+}$ and $K^{\pi, \alpha-}$ in the explicit reflected scheme where the random default time $\tau = 0.2$. We can see that $Y_t^{\pi, \alpha}$ stays between the lower obstacle $e^{-rt} L_t$ and the upper obstacle $e^{-rt} V_t$. In this example for $n = 400$, default time $\tau = 0.2$, we can get the solution $Y_0^{\pi, \alpha} = 0.6857$ from the explicit reflected scheme, i.e. the hedge for the broker c_1 against the game option at $t = 0$ in the defaultable model. In the case without the default risk, $Y_0^{\pi, \alpha} = 0.7704$,

which means the occurrence of the default event could reduce the value of $Y^{\tau, \kappa}$. Figure 11 represents the situation without the default risk, the solution $Y^{\tau, \kappa}$ has a larger value than in the situation when the default case happens (Figure 9).




Figure 8. One path of the Brownian motion.




Figure 9. One path of $Y^{\tau, \kappa}$ in the explicit reflected scheme ($\tau = 0.2$).




Figure 10. One path of the increasing processes in the explicit reflected scheme ($\tau = 0.2$).




Figure 11. The paths of $Y^{\pi, \alpha}$ in the explicit reflected scheme without default risk.

Author Contributions: Both two authors have contributed to the final version of the manuscript, and have read and agreed to the published version of the manuscript. This paper is one part of Jingnan Wang’s doctoral dissertation (Wang 2020), Ralf Korn is Jingnan Wang’s PhD supervisor. For this manuscript, Jingnan Wang developed the theoretical formalism and completed the numerical simulations through Matlab. Ralf Korn supervised the work, discussed several versions of the research with Jingnan Wang and assisted in the transformation of the research to this article format. All authors have read and agreed to the published version of the manuscript.

Funding: Jingnan Wang’s work has been carried out as a member of the DFG Research Training Group 1932 “Stochastic Models for Innovations in the Engineering Sciences. The funding by DFG is gratefully acknowledged.

Acknowledgments: We acknowledge the PhD seminar held in our department every week to exchange research progress and bring new insights.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A

Lemma A1. (Discrete Gronwall’s Inequality) (Lemma 2.2 in *Mémin et al. 2008*)

Suppose that a, b and c are positive constants, $b\Delta < 1$, $(\beta_i)_{i \in \mathbb{N}}$ is a sequence with positive values, such that

$$\beta_i + c \leq a + b\Delta \sum_{j=1}^i \beta_j, \quad i \in \mathbb{N},$$

then it follows

$$\sup_{i \leq n} \beta_i + c \leq aF_{\Delta}(b),$$

where $F_{\Delta}(b)$ is a convergent series with the following form:

$$F_{\Delta}(b) = 1 + \sum_{n=1}^{\infty} \frac{b^n}{n} (1 + \Delta) \dots (1 + (n - 1)\Delta).$$

Theorem A1. (Itô’s formula for rcll semi-martingale) (*Protter 2005*)

Let $X := (X_t)_{0 \leq t \leq T}$ be a rcll semi-martingale, g is a real value function in C^2 , therefore, $g(X)$ is also a semi-martingale, such that

$$g(X_t) = g(X_0) + \int_0^t g'(X_{s-}) dX_s + \frac{1}{2} \int_0^t g''(X_s) d[X]_s^c + \sum_{0 < s \leq t} [g(X_s) - g(X_{s-}) - g'(X_{s-}) \Delta X_s].$$

where $[X]$ is the second variation of X , $[X]^c$ is the continuous part of $[X]$, $\Delta X_s = X_s - X_{s-}$.

We give the proofs of Theorem 1, Lemma 1, Theorem 2 and Theorem 4 can be seen in Section 5.

Proof of Theorem 1. Firstly, we introduce the following ABSDE:

$$\begin{aligned}
 -dY_t^{p,q} = & f(t, Y_t^{p,q}, \mathbb{E}^{G_t}[Y_{t+\delta}^{p,q}], Z_t^{p,q}, U_t^{p,q})dt + q(Y_t^{p,q} - L_t)^- dt \\
 & - p(Y_t^{p,q} - V_t)^+ dt - Z_t^{p,q}dB_t - U_t^{p,q}dM_t, \quad t \in [0, T].
 \end{aligned}
 \tag{A1}$$

By the existence and uniqueness theorem for ABSDEs with default risk (Theorem 4.3.3 in Wang 2020), there exists the unique solution for ABSDE (A1). It follows that as $q \rightarrow \infty$, $Y_t^{p,q} \uparrow \underline{Y}_t^p$ in $\mathcal{S}_G^2(0, T + \delta; \mathbb{R})$, $Z_t^{p,q} \rightarrow \underline{Z}_t^p$ in $\mathcal{L}_G^2(0, T; \mathbb{R})$, $U_t^{p,q} \rightarrow \underline{U}_t^p$ in $\mathcal{L}_G^{2,\tau}(0, T; \mathbb{R})$, $\int_0^t q(Y_s^{p,q} - L_s)^- ds \rightarrow \underline{K}_t^p$ in $\mathcal{A}_G^2(0, T; \mathbb{R})$. $(\underline{Y}^p, \underline{Z}^p, \underline{U}^p, \underline{K}^p)$ is a solution of the following RABSDE with one obstacle L :

$$\begin{aligned}
 -d\underline{Y}_t^p = & f(t, \underline{Y}_t^p, \mathbb{E}^{G_t}[\underline{Y}_{t+\delta}^p], \underline{Z}_t^p, \underline{U}_t^p)dt + d\underline{K}_t^p \\
 & - p(\underline{Y}_t^p - V_t)^+ dt - \underline{Z}_t^p dB_t - \underline{U}_t^p dM_t, \quad t \in [0, T].
 \end{aligned}
 \tag{A2}$$

Let $p \rightarrow \infty$, it follows that $\underline{Y}_t^p \downarrow Y_t$ in $\mathcal{S}_G^2(0, T + \delta; \mathbb{R})$, $\underline{Z}_t^p \rightarrow Z_t$ in $\mathcal{L}_G^2(0, T; \mathbb{R})$, $\underline{U}_t^p \rightarrow U_t$ in $\mathcal{L}_G^{2,\tau}(0, T; \mathbb{R})$. By the comparison theorem for ABSDEs with default risk (Theorem 2.3.1 in Wang 2020), we know that \underline{K}_t^p is increasing, then $\underline{K}_T^p \uparrow K_T^{+p}$ and $\underline{K}_T^{p+1} - \underline{K}_T^p \geq \sup_{0 \leq t \leq T} [K_t^{p+1} - K_t^p] \geq 0$, therefore, $\underline{K}_t^p \rightarrow K_t^{+p}$ in $\mathcal{A}_G^2(0, T; \mathbb{R})$. Hence, there exists a constant C_1 depending on $\xi, f(t, 0, 0, 0, 0), \delta, L$ and V , such that

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} |\underline{Y}_t^p - Y_t|^2 + \int_0^T |\underline{Z}_t^p - Z_t|^2 dt + \int_0^T |\underline{U}_t^p - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] \leq \frac{C_1}{\sqrt{p}}.$$

Similarly, let $p \rightarrow \infty$ in (A1), it follows that $Y_t^{p,q} \downarrow \bar{Y}_t^q$ in $\mathcal{S}_G^2(0, T + \delta; \mathbb{R})$, $Z_t^{p,q} \rightarrow \bar{Z}_t^q$ in $\mathcal{L}_G^2(0, T; \mathbb{R})$, $U_t^{p,q} \rightarrow \bar{U}_t^q$ in $\mathcal{L}_G^{2,\tau}(0, T; \mathbb{R})$, $\int_0^t p(Y_s^p - V_s)^+ ds \rightarrow \bar{K}_t^q$ in $\mathcal{A}_G^2(0, T; \mathbb{R})$. $(\bar{Y}^q, \bar{Z}^q, \bar{U}^q, \bar{K}^q)$ is a solution of the following RABSDE with one obstacle V :

$$-d\bar{Y}_t^q = f(t, \bar{Y}_t^q, \mathbb{E}^{G_t}[\bar{Y}_{t+\delta}^q], \bar{Z}_t^q, \bar{U}_t^q)dt + q(\bar{Y}_t^q - L_t)^- dt - d\bar{K}_t^q - \bar{Z}_t^q dB_t - \bar{U}_t^q dM_t, \quad t \in [0, T].
 \tag{A3}$$

Letting $q \rightarrow \infty$, it follows that $\bar{Y}_t^q \downarrow Y_t$ in $\mathcal{S}_G^2(0, T + \delta; \mathbb{R})$, $\bar{Z}_t^q \rightarrow Z_t$ in $\mathcal{L}_G^2(0, T; \mathbb{R})$, $\bar{U}_t^q \rightarrow U_t$ in $\mathcal{L}_G^{2,\tau}(0, T; \mathbb{R})$, $\bar{K}_t^q \rightarrow K_t^{-p}$ in $\mathcal{A}_G^2(0, T; \mathbb{R})$. Moreover, there exists a constant C_2 depending on $\xi, f(t, 0, 0, 0, 0), \delta, L$ and V , such that

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} |\bar{Y}_t^q - Y_t|^2 + \int_0^T |\bar{Z}_t^q - Z_t|^2 dt + \int_0^T |\bar{U}_t^q - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] \leq \frac{C_2}{\sqrt{q}}.$$

By the comparison theorem for ABSDEs with default risk, it follows that $\underline{Y}_t^p \leq Y_t^p \leq \bar{Y}_t^p$, for any $t \in [0, T]$. Therefore,

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^p - Y_t|^2 \right] \leq \frac{C_3}{\sqrt{p}},$$

where $C_3 \geq 0$ is a constant. Applying Itô formula for rcll semi-martingale (Theorem A1), we can obtain

$$\mathbb{E} \left[\int_0^T |Z_t^p - Z_t|^2 dt + \int_0^T |U_t^p - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] \leq \frac{C_4}{\sqrt{p}},$$

where $C_4 \geq 0$ is a constant. Since

$$\begin{aligned}
 K_t^{+p} - K_t^{-p} = & Y_0^p - Y_t^p - \int_0^t f(s, Y_s^p, \mathbb{E}^{G_s}[Y_{s+\delta}^p], Z_s^p, U_s^p)ds - \int_0^t Z_s^p dB_s - \int_0^t U_s^p dM_s; \\
 K_t^+ - K_t^- = & Y_0 - Y_t - \int_0^t f(s, Y_s, \mathbb{E}^{G_s}[Y_{s+\delta}], Z_s, U_s)ds - \int_0^t Z_s dB_s - \int_0^t U_s dM_s.
 \end{aligned}$$

By the convergence of Y^p, Z^p, U^p and the Lipschitz condition of f , it follows

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq t \leq T} |(K_t^{+p} - K_t^+) - (K_t^{-p} - K_t^-)|^2 \right] \\ & \leq \lambda \mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^p - Y_t|^2 + \int_0^T |Z_t^p - Z_t|^2 dt + \int_0^T |U_t^p - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] \leq \frac{C_5}{\sqrt{p}}, \end{aligned}$$

where $\lambda, C_5 \geq 0$ are constants. Since $\mathbb{E}[|K_T^{+p}|^2 + |K_T^{-p}|^2] < \infty$, there exist processes \hat{K}^+ and \hat{K}^- in $\mathcal{A}_{\mathcal{G}}^2(0, T; \mathbb{R})$ are the weak limits of K^{+p} and K^{-p} respectively. Since for any $t \in [0, T], \underline{Y}_t^p \leq Y_t^p \leq \bar{Y}_t^p$, we can get

$$\begin{aligned} dK_t^{+p} &= p(Y_t^p - L_t)^- dt \leq p(\bar{Y}_t^p - L_t)^- dt = d\bar{K}_t^{+p}; \\ dK_t^{-p} &= p(Y_t^p - V_t)^+ dt \geq p(\underline{Y}_t^p - V_t)^+ dt = d\underline{K}_t^{-p}. \end{aligned}$$

Therefore, $d\hat{K}_t^+ \leq dK_t^+, d\hat{K}_t^- \geq dK_t^-$, it follows that $d\hat{K}_t^+ - d\hat{K}_t^- \leq dK_t^+ - dK_t^-$. On the other hand, the limit of Y^p is Y , so $d\hat{K}_t^+ - d\hat{K}_t^- = dK_t^+ - dK_t^-$, it follows that $d\hat{K}_t^+ = dK_t^+, d\hat{K}_t^- = dK_t^-$, then $\hat{K}_t^+ = K_t^+, \hat{K}_t^- = K_t^-$. \square

Proof of Lemma 1. Step 1. Firstly, we consider the continuous and discrete time equations by Picard’s method.

In the continuous case, we set $Y^{p,\infty,0} = Z^{p,\infty,0} = U^{p,\infty,0} = 0$, $(Y_t^{p,\infty,m+1}, Z_t^{p,\infty,m+1}, U_t^{p,\infty,m+1})$ is the solution of the following BSDE:

$$\left\{ \begin{aligned} Y_t^{p,\infty,m+1} &= \xi_T + \int_t^T f^n(s, Y_s^{p,\infty,m}, \mathbb{E}^{\mathcal{G}_s}[Y_{s+\delta}^{p,\infty,m}], Z_s^{p,\infty,m}, U_s^{p,\infty,m}) ds \\ &\quad + \int_t^T q(Y_s^{p,\infty,m} - L_s)^- ds - \int_t^T p(Y_s^{p,\infty,m} - V_s)^+ ds \\ &\quad - \int_t^T Z_s^{p,\infty,m+1} dB_s - \int_t^T U_s^{p,\infty,m+1} dM_s, \quad t \in [0, T]; \\ Y_t^{p,\infty,m+1} &= \xi_t, \quad t \in (T, T + \delta], \end{aligned} \right. \tag{A4}$$

where $(Y_t^{p,\infty,m}, Z_t^{p,\infty,m}, U_t^{p,\infty,m})$ is the Picard approximation of (Y_t^p, Z_t^p, U_t^p) .

In the discrete case, we set $y_i^{p,n,0} = z_i^{p,n,0} = u_i^{p,n,0} = 0$ (for any $i = 0, 2, \dots, n$), $(y_i^{p,n,m+1}, z_i^{p,n,m+1}, u_i^{p,n,m+1})$ is the solution of the following BSDE:

$$\left\{ \begin{aligned} y_i^{p,n,m+1} &= y_{i+1}^{p,n,m+1} + f^n(t_i, y_i^{p,n,m}, \bar{y}_i^{p,n,m}, z_i^{p,n,m}, u_i^{p,n,m}) \Delta^n \\ &\quad - z_i^{p,n,m+1} \Delta B_{i+1}^n - u_i^{p,n,m+1} \Delta M_{i+1}^n + p \Delta^n (y_i^{p,n} - L_i^n)^- \\ &\quad - p \Delta^n (y_i^{p,n} - V_i^n)^+, \quad i \in [0, n - 1]; \\ y_i^{p,n,m+1} &= \xi_i^n, \quad i \in [n, n^\delta]. \end{aligned} \right. \tag{A5}$$

here $(Y_t^{p,n,m}, Z_t^{p,n,m}, U_t^{p,n,m})$ is the continuous time version of the discrete Picard approximation of $(y_i^{p,n,m}, z_i^{p,n,m}, u_i^{p,n,m})$.

Step 2. Then, we consider the following decomposition:

$$Y^{p,n} - Y^p = (Y^{p,n} - Y^{p,n,m}) + (Y^{p,n,m} - Y^{p,\infty,m}) + (Y^{p,\infty,m} - Y^p).$$

From Proposition 1 and Proposition 3 in Lejay et al. (2014) and the definition of L_i^n and V_i^n , it follows (20). \square

Proof of Theorem 2. By Lemma 1 and Theorem 1, as $p \rightarrow \infty, n \rightarrow \infty$, it follows

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^{p,n} - Y_t|^2 + \int_0^T |Z_t^{p,n} - Z_t|^2 dt + \int_0^T |U_t^{p,n} - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] \\ & \leq 2\mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^{p,n} - Y_t^p|^2 + \int_0^T |Z_t^{p,n} - Z_t^p|^2 dt + \int_0^T |U_t^{p,n} - U_t^p|^2 1_{\{\tau > t\}} \gamma_t dt \right] \\ & + 2\mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^p - Y_t|^2 + \int_0^T |Z_t^p - Z_t|^2 dt + \int_0^T |U_t^p - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] \rightarrow 0. \end{aligned}$$

For the increasing processes $K^{+p,n}$ and $K^{-p,n}$, by Theorem 1, we can obtain

$$\begin{aligned} & \mathbb{E} \left[\left(K_t^{+p,n} - K_t^{-p,n} \right) - \left(K_t^+ - K_t^- \right) \right]^2 \\ & \leq 2\mathbb{E} \left[\left(K_t^{+p,n} - K_t^{-p,n} \right) - \left(K_t^{+p} - K_t^{-p} \right) \right]^2 + \frac{C}{\sqrt{p}}, \end{aligned}$$

where $C \geq 0$ is a constant depending on $\xi, f(t, 0, 0, 0, 0), \delta, L$ and V . For each fixed p ,

$$\begin{aligned} K_t^{+p,n} - K_t^{-p,n} &= Y_0^{p,n} - Y_t^{p,n} - \int_0^t f^n(s, Y_s^{p,n}, \mathbb{E}^{\mathcal{G}_s}[Y_{s+\delta}^{p,n}], Z_s^{p,n}, U_s^{p,n}) ds - \int_0^t Z_s^{p,n} dB_s^n - \int_0^t U_s^{p,n} dM_s^n; \\ K_t^{+p} - K_t^{-p} &= Y_0^p - Y_t^p - \int_0^t f(s, Y_s^p, \mathbb{E}^{\mathcal{G}_s}[Y_{s+\delta}^p], Z_s^p, U_s^p) ds - \int_0^t Z_s^p dB_s - \int_0^t U_s^p dM_s. \end{aligned}$$

From Corollary 14 in Briand et al. (2002), we know that as $n \rightarrow \infty, \int_0^\cdot Z_s^{p,n} dB_s^n \rightarrow \int_0^\cdot Z_s^p dB_s$ in $\mathcal{S}_{\mathcal{G}}^2(0, T; \mathbb{R}), \int_0^\cdot U_s^{p,n} dM_s^n \rightarrow \int_0^\cdot U_s^p dM_s$ in $\mathcal{S}_{\mathcal{G}}^2(0, T; \mathbb{R})$. By the Lipschitz condition of f and the convergence of $Y^{p,n}$, it follows that $K_t^{+p,n} - K_t^{-p,n} \rightarrow K_t^+ - K_t^-$ in $\mathcal{L}_{\mathcal{G}}^2(0, T; \mathbb{R})$. \square

Proof of Theorem 4. Firstly, we prove (23).

From Theorem A3, Lemma 1 and Theorem 1. For fixed $p \in \mathbb{N}$, as $n \rightarrow \infty$, it follows

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^n - Y_t|^2 + \int_0^T |Z_t^n - Z_t|^2 dt + \int_0^T |U_t^n - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] \\ & \leq 3\mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^n - Y_t^{p,n}|^2 + \int_0^T |Z_t^n - Z_t^{p,n}|^2 dt + \int_0^T |U_t^n - U_t^{p,n}|^2 1_{\{\tau > t\}} \gamma_t dt \right] \\ & + 3\mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^{p,n} - Y_t^p|^2 + \int_0^T |Z_t^{p,n} - Z_t^p|^2 dt + \int_0^T |U_t^{p,n} - U_t^p|^2 1_{\{\tau > t\}} \gamma_t dt \right] \\ & + 3\mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^p - Y_t|^2 + \int_0^T |Z_t^p - Z_t|^2 dt + \int_0^T |U_t^p - U_t|^2 1_{\{\tau > t\}} \gamma_t dt \right] \\ & \leq 3\mathbb{E} \left[\sup_{0 \leq t \leq T} |Y_t^{p,n} - Y_t^p|^2 + \int_0^T |Z_t^{p,n} - Z_t^p|^2 dt + \int_0^T |U_t^{p,n} - U_t^p|^2 1_{\{\tau > t\}} \gamma_t dt \right] \\ & + \frac{3}{\sqrt{p}} C_{\xi, f, L, V} + \frac{3}{\sqrt{p}} \lambda_{L, T, \delta} C_{f^n, \xi^n, L^n, V^n} \rightarrow 0. \end{aligned}$$

For increasing processes, for fixed $p \in \mathbb{N}$, as $n \rightarrow \infty$,

$$\begin{aligned} & \mathbb{E} \left[\left| (K_t^{+n} - K_t^{-n}) - (K_t^+ - K_t^-) \right|^2 \right] \\ & \leq 3\mathbb{E} \left[\left| (K_t^{+n} - K_t^{-n}) - (K_t^{+p,n} - K_t^{-p,n}) \right|^2 \right] + 3\mathbb{E} \left[\left| (K_t^{+p,n} - K_t^{-p,n}) - (K_t^{+p} - K_t^{-p}) \right|^2 \right] \\ & \quad + 3\mathbb{E} \left[\left| (K_t^{+p} - K_t^{-p}) - (K_t^+ - K_t^-) \right|^2 \right] \\ & \leq 3\mathbb{E} \left[\left| (K_t^{+p,n} - K_t^{-p,n}) - (K_t^{+p} - K_t^{-p}) \right|^2 \right] + \frac{3}{\sqrt{p}} C_{\xi, f, L, V} + \frac{3}{\sqrt{p}} \lambda_{L, T, \delta} C_{f^n, \xi^n, L^n, V^n} \rightarrow 0. \end{aligned}$$

□

We give the proof of the following Lemma A2. Lemma A3 and Lemma A4 below have the similar proof method.

Lemma A2. (Estimation result of implicit discrete penalization scheme) Under Assumption 6 and Assumption 7, for each $p \in \mathbb{N}$ and Δ^n , when $(\Delta^n + 3\Delta^n L + 4\Delta^n L^2 + (\Delta^n L)^2) < 1$, there exists a constant $\lambda_{L, T, \delta}$ depending on the Lipschitz coefficient L , T and δ , such that

$$\begin{aligned} & \mathbb{E} \left[\sup_i |y_i^{p,n}|^2 + \Delta^n \sum_{j=0}^{n-1} |z_j^{p,n}|^2 + \Delta^n \sum_{j=0}^{n-1} |u_j^{p,n}|^2 (1 - h_{j+1}^n) \gamma_{j+1} \right. \\ & \quad \left. + \frac{1}{p\Delta^n} \sum_{j=0}^{n-1} (|k_j^{+p,n}|^2 + |k_j^{-p,n}|^2) \right] \leq \lambda_{L, T, \delta} C_{\xi^n, f^n, L^n, V^n}, \end{aligned} \tag{A6}$$

where $C_{\xi^n, f^n, L^n, V^n} \geq 0$ is a constant depending on ξ^n , $f^n(t_j, 0, 0, 0, 0)$, $(L^n)^+$ and $(V^n)^-$.

Proof of Lemma A2. By the definition of implicit penalization discrete scheme (7), applying Itô formula for rcl semi-martingale (Theorem A1) to $|y_j^{p,n}|^2$ on $j \in [i, n - 1]$, it follows

$$\begin{aligned} & \mathbb{E} \left[|y_i^{p,n}|^2 + \Delta^n \sum_{j=i}^{n-1} |z_j^{p,n}|^2 + \Delta^n \sum_{j=i}^{n-1} |u_j^{p,n}|^2 (1 - h_{j+1}^n) \gamma_{j+1} \right] \\ & = \mathbb{E} |\xi_i^n|^2 + 2\Delta^n \mathbb{E} \sum_{j=i}^{n-1} \left[y_j^{p,n} \left| f^n(t_j, y_j^{p,n}, \bar{y}_j^{p,n}, z_j^{p,n}, u_j^{p,n}) \right| \right] + 2\mathbb{E} \sum_{j=i}^{n-1} \left[y_j^{p,n} k_j^{+p,n} - y_j^{p,n} k_j^{-p,n} \right], \end{aligned} \tag{A7}$$

since

$$\begin{aligned} y_j^{p,n} k_j^{+p,n} &= -\frac{1}{p\Delta^n} (k_j^{+p,n})^2 + L_j^n k_j^{+p,n}; \\ y_j^{p,n} k_j^{-p,n} &= \frac{1}{p\Delta^n} (k_j^{-p,n})^2 + V_j^n k_j^{-p,n}. \end{aligned}$$

Moreover, by the Lipschitz condition of f^n , we can obtain

$$\begin{aligned} & \mathbb{E} \left[|y_i^{p,n}|^2 + \frac{\Delta^n}{2} \sum_{j=i}^{n-1} |z_j^{p,n}|^2 + \frac{\Delta^n}{2} \sum_{j=i}^{n-1} |u_j^{p,n}|^2 (1 - h_{j+1}^n) \gamma_{j+1} + \frac{2}{p\Delta^n} \sum_{j=i}^{n-1} (|k_j^{+p,n}|^2 + |k_j^{-p,n}|^2) \right] \\ & \leq \mathbb{E} \left[|\xi_n^n|^2 + \Delta^n L \sum_{j=n+1}^{n^\delta-1} |\xi_j^n|^2 \right] + \Delta^n \mathbb{E} \left[\sum_{j=i}^{n-1} |f^n(t_j, 0, 0, 0, 0)|^2 \right] \\ & \quad + \left(\Delta^n + 3\Delta^n L + 4\Delta^n L^2 + (\Delta^n L)^2 \right) \mathbb{E} \left[\sum_{j=i}^{n-1} |y_j^n|^2 \right] + \frac{1}{\lambda_1} \mathbb{E} \left(\sum_{j=i}^{n-1} k_j^{+p,n} \right)^2 \\ & \quad + \lambda_1 \mathbb{E} \sup_{i \leq j \leq n-1} ((L_j^n)^+)^2 + \frac{1}{\lambda_1} \mathbb{E} \left(\sum_{j=i}^{n-1} k_j^{-p,n} \right)^2 + \lambda_1 \mathbb{E} \sup_{i \leq j \leq n-1} ((V_j^n)^-)^2. \end{aligned}$$

By **Assumption 8**, applying techniques of stopping times for the discrete case, it follows

$$\mathbb{E} \left(\sum_{j=i}^{n-1} k_j^{+p,n} \right)^2 + \mathbb{E} \left(\sum_{j=i}^{n-1} k_j^{-p,n} \right)^2 \leq C_{\xi^n, f^n, X^n} \left[1 + \Delta^n \mathbb{E} \sum_{j=i}^{n-1} (|z_j^{p,n}|^2 + |u_j^{p,n}|^2 (1 - h_{j+1}^n) \gamma_{j+1}) \right].$$

where $C_{\xi^n, f^n, X^n} \geq 0$ is a constant depending on ξ^n , $f^n(t_j, 0, 0, 0, 0)$, X^n . Since X^n can be dominated by L^n and V^n , we can replace it by L^n and V^n . By the discrete Gronwall's inequality (Lemma A1), when $(\Delta^n + 3\Delta^n L + 4\Delta^n L^2 + (\Delta^n L)^2) < 1$, we can obtain

$$\begin{aligned} & \sup_i \mathbb{E} \left[|y_i^{p,n}|^2 \right] + \mathbb{E} \left[\Delta^n \sum_{j=0}^n |z_j^{p,n}|^2 + \Delta^n \sum_{j=0}^n |u_j^{p,n}|^2 (1 - h_{j+1}^n) \gamma_{j+1} \right. \\ & \quad \left. + \frac{1}{p\Delta^n} \sum_{j=0}^n (|k_j^{+p,n}|^2 + |k_j^{-p,n}|^2) \right] \leq \lambda_{L,T,\delta} C_{\xi^n, f^n, L^n, V^n}, \end{aligned}$$

where $C_{\xi^n, f^n, L^n, V^n} \geq 0$ is a constant depending on ξ^n , $f^n(t_j, 0, 0, 0, 0)$, $(L^n)^+$ and $(V^n)^-$. Reconsidering (A7), we take square, sup and sum over j , then take expectation, by Burkholder-Davis-Gundy inequality for the martingale parts, it follows

$$\mathbb{E} \left[\sup_i |y_i^{p,n}|^2 \right] \leq C_{\xi^n, f^n, L^n, V^n} + C\Delta^n \left[\sum_{i=0}^{n-1} \mathbb{E} |y_i^{p,n}|^2 \right] \leq C_{\xi^n, f^n, L^n, V^n} + CT \sup_i \mathbb{E} |y_i^{p,n}|^2.$$

It follows (A6). \square

We present the proof of the following Theorem A2. Theorem A3 and Theorem A4 below have the similar proof method.

Theorem A2. (Distance between implicit discrete penalization and explicit discrete penalization schemes) Under Assumption 6 and Assumption 7, for any $p \in \mathbb{N}$:

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq t \leq T} |\check{Y}_t^{p,n} - Y_t^{p,n}|^2 + \int_0^T |\check{Z}_t^{p,n} - Z_t^{p,n}|^2 dt + \int_0^T |\check{U}_t^{p,n} - U_t^{p,n}|^2 1_{\{\tau > t\}} \gamma_t dt \right. \\ & \quad \left. + \left| (\check{K}_t^{+p,n} - \check{K}_t^{-p,n}) - (K_t^{+p,n} - K_t^{-p,n}) \right|^2 \right] \leq \lambda_{L,T,\delta} C_{f^n, \xi^n, \Delta^n, L^n, V^n, p} (\Delta^n)^2. \end{aligned} \tag{A8}$$

where $\lambda_{L,T,\delta} \geq 0$ is a constant depending on the Lipschitz coefficient L , the terminal T and δ , $C_{f^n, \xi^n, \Delta^n, L^n, V^n, p} \geq 0$ is a constant depending on $f^n(t_j, 0, 0, 0, 0)$, ξ^n , $(L^n)^+$, $(V^n)^-$ and p .

Proof of Theorem A2. From the definitions of implicit discrete penalization scheme (7) and explicit discrete penalization scheme (11), and the Lipschitz condition of f^n , it follows

$$\begin{aligned} & \mathbb{E} \left[\left| \tilde{y}_j^{p,n} - y_j^{p,n} \right|^2 + \Delta^n \left| \tilde{z}_j^{p,n} - z_j^{p,n} \right|^2 + \Delta^n \left| \tilde{u}_j^{p,n} - u_j^{p,n} \right|^2 (1 - h_{j+1}^n) \gamma_{j+1} \right] \\ & \leq 2\Delta^n L \mathbb{E} \left[\left(\left| \mathbb{E}^{G_j^n} [\tilde{y}_{j+1}^{p,n}] - y_j^{p,n} \right| + \left| \tilde{y}_j^{p,n} - \bar{y}_j^{p,n} \right| + \left| \tilde{z}_j^{p,n} - z_j^{p,n} \right| \right. \right. \\ & \quad \left. \left. + \left| \tilde{u}_j^{p,n} - u_j^{p,n} \right| (1 - h_{j+1}^n) \sqrt{\gamma_{j+1}} \right) \left(\tilde{y}_j^{p,n} - y_j^{p,n} \right) \right], \end{aligned} \tag{A9}$$

Summing from $j = i, \dots, n - 1$, it follows

$$\begin{aligned} & \mathbb{E} \left[\left| \tilde{y}_i^{p,n} - y_i^{p,n} \right|^2 + \frac{\Delta^n}{2} \sum_{j=i}^{n-1} \left| \tilde{z}_j^{p,n} - z_j^{p,n} \right|^2 + \frac{\Delta^n}{2} \sum_{j=i}^{n-1} \left| \tilde{u}_j^{p,n} - u_j^{p,n} \right|^2 (1 - h_{j+1}^n) \gamma_{j+1} \right] \\ & \leq 2\Delta^n L \mathbb{E} \sum_{j=i}^{n-1} \left[\left| \mathbb{E}^{G_j^n} [\tilde{y}_{j+1}^{p,n}] - y_j^{p,n} \right| \left| \tilde{y}_j^{p,n} - y_j^{p,n} \right| \right] + \Delta^n (2L + 4L^2) \mathbb{E} \left[\sum_{j=i}^{n-1} \left| \tilde{y}_j^{p,n} - y_j^{p,n} \right|^2 \right]. \end{aligned}$$

By (17), (11) and the Lipschitz condition of f^n , we can obtain

$$\begin{aligned} & 2\Delta^n L \mathbb{E} \sum_{j=i}^{n-1} \left[\left| \mathbb{E}^{G_j^n} [\tilde{y}_{j+1}^{p,n}] - y_j^{p,n} \right| \left| \tilde{y}_j^{p,n} - y_j^{p,n} \right| \right] \\ & \leq (\Delta^n L)^2 \mathbb{E} \sum_{j=i}^{n-1} \left[2 \left| \tilde{y}_j^{p,n} \right|^2 + \left| \tilde{z}_j^{p,n} \right|^2 + \left| \tilde{u}_j^{p,n} \right|^2 (1 - h_{i+1}^n) \gamma_{i+1} \right] + (\Delta^n)^2 \mathbb{E} \sum_{j=i}^{n-1} \left[\left| f^n(t_j, 0, 0, 0, 0) \right|^2 \right] \\ & \quad + (p\Delta^n)^2 \mathbb{E} \sum_{j=i}^{n-1} \left[\left((\tilde{y}_i^{p,n} - L_i^n)^- \right)^2 + \left((\tilde{y}_i^{p,n} - V_i^n)^+ \right)^2 \right] \\ & \quad + \left(8(\Delta^n L)^2 + 2\Delta^n L \right) \mathbb{E} \left[\sum_{i=j}^{n-1} \left| \tilde{y}_j^{p,n} - y_j^{p,n} \right|^2 \right] + (\Delta^n L)^2 \mathbb{E} \left[\sum_{i=n}^{n^\delta-1} \left| \xi_j^n \right|^2 \right], \end{aligned}$$

hence, there exists a constant $C_{f^n, \xi^n, L^n, V^n} \geq 0$ depending on $f^n(t_j, 0, 0, 0, 0)$, ξ^n , $(L^n)^+$ and $(V^n)^-$, such that

$$\begin{aligned} & \mathbb{E} \left[\left| \tilde{y}_i^{p,n} - y_i^{p,n} \right|^2 + \frac{\Delta^n}{2} \sum_{j=i}^{n-1} \left| \tilde{z}_j^{p,n} - z_j^{p,n} \right|^2 + \frac{\Delta^n}{2} \sum_{j=i}^{n-1} \left| \tilde{u}_j^{p,n} - u_j^{p,n} \right|^2 (1 - h_{j+1}^n) \gamma_{j+1} \right] \\ & \leq C_{f^n, \xi^n, L^n, V^n} (\Delta^n)^2 + \left(8(\Delta^n L)^2 + 4\Delta^n L + 4\Delta^n L^2 \right) \mathbb{E} \left[\sum_{j=i}^{n-1} \left| \tilde{y}_j^{p,n} - y_j^{p,n} \right|^2 \right]. \end{aligned}$$

By the discrete Gronwall's inequality (Lemma A1), when $(8(\Delta^n L)^2 + 4\Delta^n L + 4\Delta^n L^2) < 1$, we can get

$$\sup_i \mathbb{E} \left| \tilde{y}_i^{p,n} - y_i^{p,n} \right|^2 \leq C_{f^n, \xi^n, L^n, V^n} (\Delta^n)^2 e^{(8\Delta^n L^2 + 4L + 4L^2)T},$$

therefore, it follows

$$\mathbb{E} \left[\sum_{j=i}^{n-1} \left| \tilde{z}_j^{p,n} - z_j^{p,n} \right|^2 + \sum_{j=i}^{n-1} \left| \tilde{u}_j^{p,n} - u_j^{p,n} \right|^2 (1 - h_{j+1}^n) \gamma_{j+1} \right] \leq C_{f^n, \xi^n, L^n, V^n} (\Delta^n)^2.$$

Reconsidering (A9), we take square, sup and sum over j , then take expectation, by Burkholder-Davis-Gundy inequality for the martingale parts, we can get

$$\mathbb{E} \left[\sup_i \left| \tilde{y}_i^{p,n} - y_i^{p,n} \right|^2 \right] \leq C\Delta^n \left[\sum_{i=0}^{n-1} \mathbb{E} \left| \tilde{y}_i^{p,n} - y_i^{p,n} \right|^2 \right] \leq CT \sup_i \mathbb{E} \left| \tilde{y}_i^{p,n} - y_i^{p,n} \right|^2,$$

hence,

$$\begin{aligned} & \mathbb{E} \left[\sup_i \left| \tilde{y}_i^{p,n} - y_i^{p,n} \right|^2 + \Delta^n \sum_{j=0}^{n-1} \left| \tilde{z}_j^{p,n} - z_j^{p,n} \right|^2 + \Delta^n \sum_{j=0}^{n-1} \left| \tilde{u}_j^{p,n} - u_j^{p,n} \right|^2 (1 - h_{j+1}^n) \gamma_{j+1} \right] \\ & \leq \lambda_{L,T,\delta} C_{f^n, \xi^n, L^n, V^n, p} (\Delta^n)^2. \end{aligned} \tag{A10}$$

For the increasing processes, by the Lipschitz condition of f^n and (A10), it follows, for each fixed p ,

$$\mathbb{E} \left[\left| \left(\tilde{K}_t^{+p,n} - \tilde{K}_t^{-p,n} \right) - \left(K_t^{+p,n} - K_t^{-p,n} \right) \right|^2 \right] \leq \lambda_{L,T,\delta} C_{f^n, \xi^n, L^n, V^n, p} (\Delta^n)^2.$$

It follows (A8). \square

Similarly to the proof method of Lemma A2, we can get the following Lemma A3.

Lemma A3. (Estimation result of implicit discrete reflected scheme) Under Assumption 6 and Assumption 7, for each $p \in \mathbb{N}$ and Δ^n , when $\Delta^n + 3\Delta^n L + 4\Delta^n L^2 + (\Delta^n)^2 L^2 < 1$, there exists a constant $\lambda_{L,T,\delta}$ depending on the Lipschitz coefficient L and the terminal time T , such that

$$\mathbb{E} \left[\sup_i \left| y_i^n \right|^2 + \Delta^n \sum_{j=0}^{n-1} \left| z_j^n \right|^2 + \Delta^n \sum_{j=0}^{n-1} \left| u_j^n \right|^2 (1 - h_{j+1}^n) \gamma_{j+1} + \left| \sum_{j=0}^{n-1} k_j^{+n} \right|^2 + \left| \sum_{j=0}^{n-1} k_j^{-n} \right|^2 \right] \leq \lambda_{L,T,\delta} C_{\xi^n, f^n, L^n, V^n}, \tag{A11}$$

where $C_{\xi^n, f^n, L^n, V^n} \geq 0$ is a constant depending on $\xi^n, f^n(t_j, 0, 0, 0, 0), (L^n)^+$ and $(V^n)^-$.

Similarly to the proof method of Theorem A2, we can obtain the Theorem A3 below.

Theorem A3. (Distance between implicit discrete penalization and implicit discrete reflected schemes) Under Assumption 6 and Assumption 7, for any $p \in \mathbb{N}$:

$$\begin{aligned} & \mathbb{E} \left[\sup_{0 \leq t \leq T} \left| Y_t^n - Y_t^{p,n} \right|^2 + \int_0^T \left| Z_t^n - Z_t^{p,n} \right|^2 dt + \int_0^T \left| U_t^n - U_t^{p,n} \right|^2 \mathbf{1}_{\{\tau > t\}} \gamma_t dt \right. \\ & \left. + \left| \left(K_t^{+n} - K_t^{-n} \right) - \left(K_t^{+p,n} - K_t^{-p,n} \right) \right|^2 \right] \leq \lambda_{L,T,\delta} C_{f^n, \xi^n, L^n, V^n} \frac{1}{\sqrt{p}}, \end{aligned} \tag{A12}$$

where $\lambda_{L,T,\delta} \geq 0$ is a constant depending on the Lipschitz coefficient L , the terminal time T and δ , $C_{f^n, \xi^n, L^n, V^n} \geq 0$ is a constant depending on $f^n(t_j, 0, 0, 0, 0), \xi^n, L^n$ and V^n .

Similarly to the proof of Lemma A2, we can get the following Lemma A4.

Lemma A4. (Estimation result of explicit discrete reflected scheme) Under Assumption 6 and Assumption 7, for each $p \in \mathbb{N}$ and Δ^n , when $\frac{7\Delta^n}{4} + 2\Delta^n L + 12\Delta^n L^2 + 10(\Delta^n L)^2 < 1$, there exists a constant $\lambda_{L,T,\delta}$ depending on the Lipschitz coefficient L , T and δ , such that

$$\mathbb{E} \left[\sup_i \left| \tilde{y}_i^n \right|^2 + \Delta^n \sum_{j=0}^{n-1} \left| \tilde{z}_j^n \right|^2 + \Delta^n \sum_{j=0}^{n-1} \left| \tilde{u}_j^n \right|^2 (1 - h_{j+1}^n) \gamma_{j+1} + \left| \sum_{j=0}^{n-1} \tilde{k}_j^{+n} \right|^2 + \left| \sum_{j=0}^{n-1} \tilde{k}_j^{-n} \right|^2 \right] \leq \lambda_{L,T,\delta} C_{\xi^n, f^n, L^n, V^n}, \tag{A13}$$

where $C_{\xi^n, f^n, L^n, V^n} \geq 0$ is a constant depending on $\xi^n, f^n(t_j, 0, 0, 0, 0), (L^n)^+$ and $(V^n)^-$.

Similarly to the proof method of Theorem A2, we can obtain the Theorem A4 below.

Theorem A4. (Distance between implicit discrete reflected and explicit discrete reflected schemes)
Under Assumption 3 and Assumption 7, for any $p \in \mathbb{N}$:

$$\begin{aligned} \mathbb{E} \left[\sup_{0 \leq t \leq T} |\tilde{Y}_t^n - Y_t^n|^2 + \int_0^T |\tilde{Z}_t^n - Z_t^n|^2 dt + \int_0^T |\tilde{U}_t^n - U_t^n|^2 1_{\{\tau > t\}} \gamma_t dt \right. \\ \left. + |(\tilde{K}_t^{+n} - \tilde{K}_t^{-n}) - (K_t^{+n} - K_t^{-n})|^2 \right] \leq \lambda_{L,T,\delta} C_{f^n, \xi^n, L^n, V^n, p} (\Delta^n)^2. \end{aligned} \tag{A14}$$

where $\lambda_{L,T,\delta} \geq 0$ is a constant depending on Lipschitz coefficient L, T and $\delta, C_{f^n, \xi^n, \Delta^n, L^n, V^n, p} \geq 0$ is a constant depending on $f^n(t_j, 0, 0, 0, 0), \xi^n, L^n, V^n$ and p .

Theorem A5. For any $s \in [t, T], e^{rt} Y_t^{\pi, \alpha}$ is a hedge for the broker c_1 against the game option, i.e., $J_t = e^{rt} Y_t^{\pi, \alpha}$.

Proof of Theorem A5. Step 1. We first prove $J_t \geq e^{rt} Y_t^{\pi, \alpha}$.

Similarly to the proof method of Theorem 5.1 in Hamadène (2006), for any fixed time $t \in [0, T], (\pi, \theta)$ is a hedge after t for the broker against the American game option. By (29) and (30), it follows that $\theta \geq t, \pi = (\beta_s^{(1)}, \beta_s^{(2)})_{s \in [t, T]}$ is a self-financing portfolio whose value at time t is A , satisfying $A_s^{\pi, \alpha} \geq R(s, \theta)$, here $s \in [t, T]$. By (27) and Itô formula for rcll semi-martingale (Theorem A1), we can obtain

$$e^{-r(s \wedge \theta)} A_{s \wedge \theta}^{\pi, \alpha} = e^{-rt} \alpha^A + \int_t^{s \wedge \theta} \beta_u^{(2)} e^{-ru} S \sigma_u dB_u^{\pi, \alpha} + \int_t^{s \wedge \theta} \beta_u^{(2)} e^{-ru} \chi S_u dM_u^{\pi, \alpha} \geq e^{-r(s \wedge \theta)} R(s, \theta) \tag{A15}$$

Let $v \in [t, T]$ be a \mathcal{G} -stopping time, setting $s = v$ and taking the conditional expectation in (A15), it follows

$$e^{-rt} \alpha^A \geq \mathbb{E}^{\pi, \alpha} \left[e^{-r(v \wedge \theta)} R(v, \theta) \middle| \mathcal{G}_t \right].$$

Hence, similarly to the result of Proposition 4.3 in Hamadène (2006),

$$\begin{aligned} e^{-rt} \alpha^A &\geq \text{ess sup}_{v \geq t} \mathbb{E}^{\pi, \alpha} \left[e^{-r(v \wedge \theta)} R(v, \theta) \middle| \mathcal{G}_t \right] \\ &\geq \text{ess inf}_{\theta \geq t} \text{ess sup}_{v \geq t} \mathbb{E}^{\pi, \alpha} \left[e^{-r(v \wedge \theta)} R(v, \theta) \middle| \mathcal{G}_t \right] = Y_t^{\pi, \alpha}. \end{aligned}$$

It follows $J_t \geq e^{rt} Y_t^{\pi, \alpha}$.

Step 2. Then prove $J_t \leq e^{rt} Y_t^{\pi, \alpha}$.

By the definition of θ_t^* in (32), it follows

$$Y_{s \wedge \theta_t^*}^{\pi, \alpha} = Y_t^{\pi, \alpha} - K_{s \wedge \theta_t^*}^{\pi, \alpha, +} + \int_t^{s \wedge \theta_t^*} Z_u^{\pi, \alpha} dB_u^{\pi, \alpha} + \int_t^{s \wedge \theta_t^*} U_u^{\pi, \alpha} dM_u^{\pi, \alpha}. \tag{A16}$$

Since

$$Y_{\theta_t^*}^{\pi, \alpha} 1_{\{\theta_t^* < T\}} = e^{-r\theta_t^*} U_{\theta_t^*} 1_{\{\theta_t^* < T\}} \geq e^{-r\theta_t^*} Q_{\theta_t^*} 1_{\{\theta_t^* < T\}},$$

therefore, by (A16), we can obtain

$$\begin{aligned} Y_{s \wedge \theta_t^*}^{\pi, \alpha} &= Y_s^{\pi, \alpha} 1_{\{s < \theta_t^*\}} + Y_{\theta_t^*}^{\pi, \alpha} 1_{\{\theta_t^* < s\}} + Y_{\theta_t^*}^{\pi, \alpha} 1_{\{s = \theta_t^* < T\}} + \zeta 1_{\{s = \theta_t^* < T\}} \\ &\geq e^{-rs} L_s 1_{\{s < \theta_t^*\}} + e^{-r\theta_t^*} U_{\theta_t^*} 1_{\{\theta_t^* < T\}} + e^{-r\theta_t^*} Q_{\theta_t^*} 1_{\{s = \theta_t^* < T\}} + \zeta 1_{\{s = \theta_t^* < T\}} \\ &= e^{-r(s \wedge \theta_t^*)} R(s, \theta_t^*). \end{aligned}$$

It follows

$$Y_t^{\pi, \alpha} + \int_t^{s \wedge \theta_t^*} Z_u^{\pi, \alpha} dB_u^{\pi, \alpha} + \int_t^{s \wedge \theta_t^*} U_u^{\pi, \alpha} dM_u^{\pi, \alpha} \geq e^{-r(s \wedge \theta_t^*)} R(s, \theta_t^*), \quad s \in [t, T].$$

Set

$$\begin{aligned} \bar{A}_s &= e^{rs} \left(Y_t^{\pi, \alpha} + \int_t^{s \wedge \theta_t^*} Z_u^{\pi, \alpha} dB_u^{\pi, \alpha} + \int_t^{s \wedge \theta_t^*} U_u^{\pi, \alpha} dM_u^{\pi, \alpha} \right); \\ \beta_s^2 &= e^{rs} \left(Z_s^{\pi, \alpha} (\sigma S_s)^{-1} + U_s^{\pi, \alpha} (\chi S_s)^{-1} \right) 1_{\{s \leq \theta_t^*\}}; \\ \beta_s^1 &= \left(\bar{A}_s - \beta_s^{(2)} S_s \right) C_s^{-1}. \end{aligned}$$

Obviously, $\bar{A}_s = \beta_s^{(1)} C_s + \beta_s^{(2)} S_s$. Applying Itô formula for rcll semi-martingale (Theorem A1), we can obtain

$$\begin{aligned} \bar{A}_s &= e^{rt} Y_t^{\pi, \alpha} + \int_t^s r \bar{A}_u du + \int_t^s e^{ru} Z_u^{\pi, \alpha} 1_{\{s \leq \theta_t^*\}} dB_u^{\pi, \alpha} + \int_t^s e^{ru} U_u^{\pi, \alpha} 1_{\{s \leq \theta_t^*\}} dM_u^{\pi, \alpha} \\ &= e^{rs} Y_t^{\pi, \alpha} + \int_t^s e^{ru} \beta_u^{(1)} dC_u + \int_t^s e^{ru} \beta_u^{(2)} dS_u. \end{aligned}$$

So $\pi = (\beta_s^{(1)}, \beta_s^{(2)})_{s \in [t, T]}$ is a self-financing portfolio with value $e^{rt} Y_t^{\pi, \alpha}$ at time t . Since for any $s \in [t, T]$, $\bar{A}_{s \wedge \theta_t^*} \geq R(s, \theta_t^*)$, then (π, θ_t^*) is a hedge strategy against this American game option, it follows that $J_t \leq e^{rt} Y_t^{\pi, \alpha}$. \square

References

- Barles, Guy, Rainer Buckdahn, and Étienne Pardoux. 1997. Backward stochastic differential equations and integral-partial differential equations. *Stochastics: An International Journal of Probability and Stochastic Processes* 60: 57–83. [\[CrossRef\]](#)
- Bielecki, Tomasz, Monique Jeanblanc, and Marek Rutkowski. 2007. *Introduction to Mathematics of Credit Risk Modeling*. Stochastic Models in Mathematical Finance. Marrakech: CIMPÀ–UNESCO–MOROCCO School, pp. 1–78.
- Bismut, Jean Michel. 1973. Conjugate convex functions in optimal stochastic control. *Journal of Mathematical Analysis and Applications* 44: 384–404. [\[CrossRef\]](#)
- Briand, Philippe, Bernard Delyon, and Jean Mémin. 2002. On the robustness of backward stochastic differential equations. *Stochastic Processes and their Applications* 97: 229–53. [\[CrossRef\]](#)
- Cordoni, Francesco, and Luca Di Persio. 2014. Backward stochastic differential equations approach to hedging, option pricing, and insurance problems. *International Journal of Stochastic Analysis* 2014: 152389. [\[CrossRef\]](#)
- Cordoni, Francesco, and Luca Di Persio. 2016. A bsde with delayed generator approach to pricing under counterparty risk and collateralization. *International Journal of Stochastic Analysis* 2016: 1059303. [\[CrossRef\]](#)
- Cvitanic, Jaksa, and Ioannis Karatzas. 1996. Backward stochastic differential equations with reflection and dynkin games. *Annals of Probability* 24: 2024–56. [\[CrossRef\]](#)
- Duffie, Darrell, and Larry Epstein. 1992. Stochastic differential utility. *Econometrica: Journal of the Econometric Society* 60: 353–94. [\[CrossRef\]](#)
- Dumitrescu, Roxana, and Céline Labart. 2016. Numerical approximation of doubly reflected bsdes with jumps and rcll obstacles. *Journal of Mathematical Analysis and Applications* 442: 206–43. [\[CrossRef\]](#)
- El Karoui, Nicole, Christophe Kapoudjian, Étienne Pardoux, Shige Peng, and Marie Claire Quenez. 1997a. Reflected solutions of backward sdes and related obstacle problems for pdes. *Annals of Probability* 25: 702–37. [\[CrossRef\]](#)
- El Karoui, Nicole, Étienne Pardoux, and Marie Claire Quenez. 1997b. Reflected backward sdes and american options. *Numerical Methods in Finance* 13: 215–31.
- Hamadène, Saïd. 2006. Mixed zero-sum stochastic differential game and american game options. *SIAM Journal on Control and Optimization* 45: 496–518.

- Hamadène, Said, Jean Pierre Lepeltier, and Anis Matoussi. 1997. Double barrier backward sdes with continuous coefficient. *Pitman Research Notes in Mathematics Series* 41: 161–76.
- Hamadène, Said, and Jean Pierre Lepeltier. 2000. Reflected bsdes and mixed game problem. *Stochastic Processes and Their Applications* 85: 177–88. [CrossRef]
- Jeanblanc, Monique, Thomas Lim, and Nacira Agram. 2017. Some existence results for advanced backward stochastic differential equations with a jump time. *ESAIM: Proceedings and Surveys* 56: 88–110. [CrossRef]
- Jiao, Ying, Idris Kharroubi, and Huyen Pham. 2013. Optimal investment under multiple defaults risk: A bsde-decomposition approach. *The Annals of Applied Probability* 23: 455–91. [CrossRef]
- Jiao, Ying, and Huyên Pham. 2011. Optimal investment with counterparty risk: A default-density model approach. *Finance and Stochastics* 15: 725–53. [CrossRef]
- Karatzas, Ioannis, and Steven Shreve. 1998. *Methods of Mathematical Finance*. New York: Springer, vol. 39.
- Kusuoka, Shigeo. 1999. A remark on default risk models. *Advances in Mathematical Economics* 1: 69–82.
- Lejay, Antoine, Ernesto Mordecki, and Soledad Torres. 2014. Numerical approximation of backward stochastic differential equations with jumps. Research report. inria-00357992v3. Available online: <https://hal.inria.fr/inria-00357992v3/document> (accessed on 30 June 2020).
- Lepeltier, Jean Pierre, and Jaime San Martín. 2004. Backward sdes with two barriers and continuous coefficient: An existence result. *Journal of Applied Probability* 41: 162–75. [CrossRef]
- Lepeltier, Jean Pierre, and Mingyu Xu. 2007. Reflected backward stochastic differential equations with two roll barriers. *ESAIM: Probability and Statistics* 11: 3–22. [CrossRef]
- Lin, Yin, and Hailiang Yang. 2014. Discrete-time bsdes with random terminal horizon. *Stochastic Analysis and Applications* 32: 110–27. [CrossRef]
- Mémin, Jean, Shige Peng, and Mingyu Xu. 2008. Convergence of solutions of discrete reflected backward sdes and simulations. *Acta Mathematicae Applicatae Sinica, English Series* 24: 1–18. [CrossRef]
- Øksendal, Bernt, Agnes Sulem, and Tusheng Zhang. 2011. Optimal control of stochastic delay equations and time-advanced backward stochastic differential equations. *Advances in Applied Probability* 43: 572–96. [CrossRef]
- Pardoux, Étienne, and Shige Peng. 1990. Adapted solution of a backward stochastic differential equation. *Systems and Control Letters* 14: 55–61. [CrossRef]
- Peng, Shige, and Mingyu Xu. 2011. Numerical algorithms for backward stochastic differential equations with 1-d brownian motion: Convergence and simulations. *ESAIM: Mathematical Modelling and Numerical Analysis* 45: 335–60. [CrossRef]
- Peng, Shige, and Xiaoming Xu. 2009. Bsdes with random default time and their applications to default risk. *arXiv* arXiv:0910.2091.
- Peng, Shige, and Zhe Yang. 2009. Anticipated backward stochastic differential equations. *Annals of Probability* 37: 877–902. [CrossRef]
- Potter, Philip. 2005. *Stochastic Integration and Differential Equations*. Berlin/Heidelberg: Springer.
- Tang, Shanjian, and Xunjing Li. 1994. Necessary conditions for optimal control of stochastic systems with random jumps. *SIAM: Journal on Control and Optimization* 32: 1447–75. [CrossRef]
- Wang, Jingnan. 2020. Reflected anticipated backward stochastic differential equations with default risk, numerical algorithms and applications. Doctoral dissertation, University of Kaiserslautern, Kaiserslautern, Germany.
- Xu, Mingyu. 2011. Numerical algorithms and simulations for reflected backward stochastic differential equations with two continuous barriers. *Journal of Computational and Applied Mathematics* 236: 1137–54. [CrossRef]

