

A Service of

ZBW

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Zoia, Maria Grazia; Vacca, Gianmarco; Barbieri, Laura

Article

Modeling multivariate financial series and computing risk measures via Gram-Charlier-like expansions

Risks

Provided in Cooperation with: MDPI – Multidisciplinary Digital Publishing Institute, Basel

Suggested Citation: Zoia, Maria Grazia; Vacca, Gianmarco; Barbieri, Laura (2020) : Modeling multivariate financial series and computing risk measures via Gram-Charlier-like expansions, Risks, ISSN 2227-9091, MDPI, Basel, Vol. 8, Iss. 4, pp. 1-21, https://doi.org/10.3390/risks8040123

This Version is available at: https://hdl.handle.net/10419/258076

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.



WWW.ECONSTOR.EU

https://creativecommons.org/licenses/by/4.0/

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.









Modeling Multivariate Financial Series and Computing Risk Measures via Gram–Charlier-Like Expansions

Maria Grazia Zoia^{1,*}, Gianmarco Vacca¹, and Laura Barbieri²

- ¹ Department of Economic Policy, Università Cattolica del Sacro Cuore, 20123 Milan, Italy; gianmarco.vacca@unicatt.it
- ² Department of Economic and Social Sciences (DiSES), Università Cattolica del Sacro Cuore, 29122 Piacenza, Italy; laura.barbieri@unicatt.it
- * Correspondence: maria.zoia@unicatt.it

Received: 22 October 2020; Accepted: 12 November 2020; Published: 16 November 2020



Abstract: This paper develops an approach based on Gram–Charlier-like expansions for modeling financial series to take in due account features such as leptokurtosis. A Gram–Charlier-like expansion adjusts the moments of interest of a given distribution via its own orthogonal polynomials. This approach, formerly adopted for univariate series, is here extended to a multivariate context by means of spherical densities. Previous works proposed the Gram–Charlier of the multivariate Gaussian, obtained by using Hermite polynomials. This work shows how polynomial expansions of an entire class of spherical laws can be worked out with the aim of obtaining a wide set of leptokurtic multivariate distributions. A Gram–Charlier-like expansion is a distribution characterized by an additional parameter with respect to the parent spherical law. This parameter, which measures the increase in kurtosis due to the polynomial expansion, can be estimated so as to make the resulting distribution capable of describing the empirical kurtosis found in the data. An application of the Gram–Charlier-like expansions to a set of financial assets proves their effectiveness in modeling multivariate financial series and assessing risk measures, such as the value at risk and the expected shortfall.

Keywords: orthogonal polynomials; kurtosis; power raised hyperbolic-secant distributions; value at risk; expected shortfall

JEL Classification: C01; C51; C58

1. Introduction

There are plenty of examples in the literature on the non-normality of asset returns and its implications for pricing and measuring financial risk. Several papers (see, e.g., Boudt et al. 2007; Jondeau and Rockinger 2005, 2006a, 2006b, 2007; Jurczenko et al. 2004; Mandelbrot 1997) have analyzed the consequences arising from the misspecification of models related to the normality assumption and the importance of correctly accounting for stylized facts that are typically exhibited by financial series, such as leptokurtosis, volatility clustering, and asymmetry. They have also highlighted the convenience of modeling the joint portfolio distribution by assuming multivariate leptokurtic specifications. The attraction of referring to heavy tailed distributions for modeling and assessing risk measures is also well documented in the literature (Eun et al. 2004; Harmantzis et al. 2006; Marinelli et al. 2012; Rachev 2003).

Leptokurtic distributions can be easily obtained by reshaping a given law through its own orthogonal polynomials. The resulting distribution is called Gram–Charlier (GCE) or Gram–Charlier-Like Expansion

(GCLE), depending on what is being applied: the Gaussian or another law. The use of orthogonal polynomials to modify the moments of a given distributions was suggested by (Faliva et al. 2016; Jondeau and Rockinger 2001; Zoia 2010). Their work showed how a given parent density can be reshaped from the inside by using its own properly weighted orthogonal polynomials. The degree of the polynomial depends on the order of the moments to be adjusted, while the weights mirror the changes introduced in the moments of the parent distribution. The choice of the latter should depend on the features (skewness and kurtosis, for example) of the empirical series under investigation. In fact, a polynomially-modified distribution maintains some desirable properties, such as unimodality, once its moments do not exceed certain critical thresholds, which depend on some features of the parent law. In particular, when the fourth moment is concerned, the more leptokurtic the parent distribution is, the higher the upward push in this moment and, consequently, the higher the kurtosis of the polynomially-modified distribution (see Bagnato et al. 2015).

When the initial distribution is Gaussian, the intended orthogonal polynomials are the Hermite ones, and the polynomially-modified distribution is the classical GCE. Jondeau and Rockinger (2001) and Zoia (2010) showed that GCE are not adequate to fit well highly leptokurtic series, given that the attainable increase in the kurtosis obtained via GCE is moderate. To overcome this problem, Bagnato et al. (2015) proposed moving to other leptokurtic univariate distributions, such as the logistic and the hyperbolic-secant. The polynomially-modified versions of these distributions are termed GCLEs.

Some works in the literature (see, e.g., Berkowitz and Garner 1970; Weng 2015) proposed the extension of the Gram-Charlier expansions to the multivariate framework. In particular, Del Brio et al. (2009) and Bagnato et al. (2017) proposed to reshape the multivariate Gaussian, with its associated Hermite polynomials, to obtain a leptokurtic distribution to fit data exhibiting excess kurtosis. Unfortunately, the limitations involved in the polynomial expansion of the univariate Gaussian also apply to the multivariate case. For this reason, in this paper, we propose the polynomial expansion of spherical variables with the Mardia kurtosis index (Mardia 1970) higher than that of the multivariate Gaussian law. This approach allows us to obtain a class of multivariate leptokurtic distributions with a resulting kurtosis range that is capable of covering the kurtosis empirically found in multivariate financial data. The contribution of this paper is twofold. First, it shows how spherical variables can be built for the the power-raised hyperbolic-secant density family, which, besides the Gaussian, also includes leptokurtic distributions like the logistic and the hyperbolic-secant law (see Engel et al. 2010; Faliva and Zoia 2017; Vaughan 2002). Second, it explains how multivariate heavy-tailed distributions can be obtained through the polynomial expansions of these spherical laws. It is proven that the orthogonal polynomials associated with spherical variables can be computed in a fairly straightforward manner from those of their associated generating variates. Since the latter are one-dimensional variables, the problem becomes easily tractable.

The resulting GCLEs are leptokurtic distributions with closed form representations characterized by one additional parameter denoting the excess kurtosis with respect to the parent spherical distribution. This parameter can be easily estimated and set equal to the value that makes the empirical kurtosis equal to that of the GCLE at hand. Thus, one of the main advantages of using GCEs is that, unlike other alternative methods (see Kotz and Nadarajah 2004; Kozubowski et al. 2013; Madan 2020), they can be effectively tailored on the features, in particular the leptokurtosis, encountered in the data.

An application of the GCLEs, to a set of international financial assets, provides evidence of their good performance both in fitting data and assessing risk measures like the Value at Risk (VaR) and the Expected Shortfall (ES), both in- and out-of-sample. In this regard, we generalize the Δ VaR method for linear portfolios with elliptically distributed risk factors, as proposed by (Kamdem 2005; Nesmith et al. 2017), to portfolios where the risk factors are GCLE distributed.

The structure of the paper is as follows. In Section 2, the paper's new methodological approach is set out. Here, it is explained how to construct spherical laws for the class of the power-raised hyperbolic-secant densities and their GCLE. In Section 3, the effectiveness of GCLEs in modeling

financial series and in computing risk measures, such as the value at risk and the expected shortfall, is investigated. Section 4 draws some conclusions. Appendix A provides the proofs of the methodological results presented in Section 2.

2. Spherical Extensions of Gram–Charlier-Like Expansions

Zoia (2010) and Bagnato et al. (2015) proved that the kurtosis of a given univariate even density f(x) can be adjusted via a binomial of the form:

$$q(x,\beta) = 1 + \beta p_4(x) \tag{1}$$

where β is a positive coefficient and $p_4(x)$ is the fourth-degree orthogonal polynomial:

$$p_4(x) = \frac{1}{b_4}(x^4 - b_2x^2 + b_0) \tag{2}$$

with coefficients b_i , j = (0, 2, 4) specified in terms of the (even) moments m_j of f(x) as follows:

$$b_2 = \frac{m_6 - m_2 m_4}{m_4 - m_2^2}, \quad b_0 = \frac{m_6 m_2 - m_4^2}{m_4 - m_2^2}, \quad b_4 = m_8 - b_2 m_6 + b_0 m_4.$$
 (3)

Under the condition:

$$0 < \beta \le \frac{4b_4}{b_2^2 - 4b_0} \tag{4}$$

the binomial (1) is positive, and the Gram-Charlier-Like Expansion (GCLE):

$$\widetilde{f}(x,\beta) = q(x,\beta)f(x)$$
 (5)

is a density whose fourth moment, \tilde{m}_4 , is increased by a quantity equal to β , with respect to the same moment m_4 of its parent density f(x), that is:

$$\widetilde{m}_4 = m_4 + \beta \tag{6}$$

while its lower order moments remain unchanged (see Bagnato et al. 2015).

This approach, which has been successfully applied to univariate distributions, is here extended to spherical densities.

Spherical distributions can be viewed as multivariate generalization of univariate symmetrical densities and prove useful in several financial studies (see, e.g., Landsman and Valdez 2003; Szegö 2004).

As is well known, any spherically distributed random vector **x** admits a stochastic representation of the form:

$$\mathbf{x} = R\mathbf{U} \tag{7}$$

where $R = ||\mathbf{x}|| = (\mathbf{x}'\mathbf{x})^{1/2}$ is a non-negative random variable, called the generating variate, independent of **U**, which is uniformly distributed on the (n - 1)-dimensional unit hyper sphere.

In Equation (7), **U** produces elliptically contoured surfaces, while *R* determines the shape and, in particular, the kurtosis of the distributions, as shown in the next theorem, which introduces some results concerning these variables that play a crucial role in the forthcoming analysis.

Theorem 1. *The density of an n-dimensional spherical variable x is:*

$$g_n(\mathbf{x}) = \frac{k_n}{M(n)} g(\mathbf{x}'\mathbf{x})$$
(8)

Here:

$$k_n = \frac{\Gamma(n/2)}{2\pi^{n/2}} \tag{9}$$

where $\Gamma(\cdot)$ denotes the gamma function and $g(\cdot)$ is a non-negative Lebesgue measurable function, named the density generator, with finite Mellin transform (see Zayed 1996, p. 201):

$$M(n) = \int_0^\infty r^{n-1} g(r^2) dr < \infty, \quad n > 0,$$
 (10)

The density generator is related to the density function of R as follows:

$$h(r) = \frac{r^{n-1}}{k_n} g(r^2)$$
(11)

where $r = (\mathbf{x}'\mathbf{x})^{1/2}$.

Mardia's kurtosis index (Mardia 1970) of the spherical vector \mathbf{x} , K^{SP} , is given by:

$$K^{SP} = m_4 \tag{12}$$

where m_{2i} is the *j*-th moment of R^2 :

$$m_{2j} = \mathbb{E}\left[(R^2)^j \right] = \frac{M(n+2j)}{M(n)}, \ j = 0, 1, 2, \dots$$
 (13)

that tallies with the 2*j*-th norm-moment of the spherical density $g_n(x)$:

$$m_{2j} = \int_{\mathbb{R}^n} ||\mathbf{x}||^{2j} g_n(||\mathbf{x}||^2) d\mathbf{x} = \int_{\mathbb{R}^n} (\mathbf{x}'\mathbf{x})^j g_n(\mathbf{x}'\mathbf{x}) d\mathbf{x}$$
(14)

with $d\mathbf{x}$ standing for $\prod_{i=1}^{n} dx_i$.

Proof. See Appendix A. \Box

The next theorem shows how to obtain GCLEs of spherical laws and provides their Mardia kurtosis indexes. The latter highlight the rise of kurtosis, which is introduced in spherical laws by the polynomial expansion.

Theorem 2. A GCLE, based on the fourth-degree orthogonal polynomial, of an *n*-dimensional spherical variable x with density function $g_n(x)$ is defined as:

$$\tilde{g}_n(\boldsymbol{x},\boldsymbol{\beta}) = q(\boldsymbol{x},\boldsymbol{\beta})g_n(\boldsymbol{x}) \tag{15}$$

Here, $q(x, \beta)$ *is the binomial defined in* (1) *with* $p_4(x)$ *specified as:*

$$p_4\left((\mathbf{x}'\mathbf{x})^{1/2}\right) = \frac{1}{b_4}\left[(\mathbf{x}'\mathbf{x})^2 - b_2(\mathbf{x}'\mathbf{x}) + b_0\right]$$
(16)

with coefficients, b_2 and b_4 , determined as in (3) from the (even) norm-moments m_{2j} of the spherical distribution defined as in (14):

$$m_{2j} = \int_{\mathbb{R}^n} (\mathbf{x}'\mathbf{x})^j g_n(\mathbf{x}'\mathbf{x})$$
(17)

Mardia's kurtosis index of the GCLE as defined in (15) is:

$$K_{GCLE}^{SP} = K^{SP} + \beta = \frac{M(n+4)}{M(n)} + \beta$$
(18)

Proof. See Appendix A. \Box

Spherical distributions can be built from univariate laws. In the following, we focus on spherical distributions ensuing from the so-called Power-raised Hyperbolic Secant densities (PHS). This is a family of bell-shaped symmetric univariate distributions arising from the hyperbolic-secant law raised to a positive power λ , which in standard form can be specified as follows:

$$g_{\lambda}(y) = b \left[\operatorname{sech}(ay^{1/2}) \right]^{\lambda}, \quad \lambda > 0$$
(19)

where:

$$a = \left[\frac{1}{2}\psi^{(1)}\left(\frac{\lambda}{2}\right)\right]^{1/2}, \quad b = a\left[B\left(\frac{\lambda}{2}, \frac{1}{2}\right)\right]^{-1}$$
(20)

In (20), $\psi^{(1)}(\cdot)$ is the trigamma function and $B(\cdot, \cdot)$ is the beta function. It can be proven that this family embraces both the Gaussian law, for $\lambda \to \infty$, and the Laplace law, for $\lambda \to 0$, as limit distributions (see Faliva and Zoia 2017). The logistic and hyperbolic-secant distributions arise when $\lambda = 2$ and $\lambda = 1$, respectively. Thus, this family includes distributions whose kurtosis ranges from three (corresponding to the case of the Gaussian law) to six (corresponding to the case of the Laplace distribution).

The next theorem explains how to build Spherical laws from PHS densities (SPHS).

Theorem 3. An *n*-dimensional SPHS engendered by the power-raised hyperbolic-secant family has a density of the form:

$$g_{n,\lambda}(\mathbf{x}) = \frac{a^n k_n}{M(n,\lambda)} \left[\operatorname{sech} \left(a(\mathbf{x}'\mathbf{x})^{1/2} \right) \right]^{\lambda}, \quad \lambda > 0$$
⁽²¹⁾

where k_n is as in (9) and $M(n, \lambda)$ is the (finite) Mellin transform:

$$M(n,\lambda) = \int_0^\infty v^{n-1} \left[\operatorname{sech}(v) \right]^\lambda dv.$$
(22)

Mardia's kurtosis index of an SPHS variable is given by:

$$K^{SPHS} = \frac{M(n+4,\lambda)}{a^4 M(n,\lambda)}$$
(23)

Proof. See Appendix A. \Box

In what follows, we consider some SPHS laws that are characterized by different kurtosis levels, namely the Gaussian, the logistic, and the hyperbolic-secant law.

Corollary 1. The *n*-variate spherical hyperbolic-secant has a density specified as follows:

$$g_{n,\lambda=1}(\mathbf{x}) = \frac{2^{n-2}\pi^{n/2}}{\left(\frac{n}{2}\right)\frac{n}{2}\phi(n)} \operatorname{sech}\left(\frac{\pi}{2}(\mathbf{x}'\mathbf{x})^{1/2}\right)$$
(24)

Here, $(n)_k$ *is the Pochhammer symbol and:*

$$\phi(n) = \{\xi(n, 1/4) - \xi(n, 3/4)\}$$
(25)

with $\xi(\cdot, \cdot)$ denoting the Hurwitz zeta function.

The *n*-variate spherical logistic has a density specified as:

$$g_{n,\lambda=2}(\mathbf{x}) = c_n \operatorname{sech}^2\left(\frac{\pi}{2\sqrt{3}}(\mathbf{x}'\mathbf{x})^{1/2}\right)$$
(26)

where:

$$c_n = \begin{cases} \frac{\pi}{24\ln 2} & \text{if } n = 2\\ \frac{\pi^{n/2}}{8(\sqrt{3})^n (1 - 2^{2-n})\zeta(n-1)\left(\frac{n}{2}\right)_{\frac{n}{2}}} & \text{otherwise} \end{cases}$$
(27)

with $\zeta(\cdot)$ denoting the Riemann zeta function.

Finally, the n-variate spherical Gaussian distribution has a density specified as follows:

$$g_{n,\lambda\to\infty}(\mathbf{x}) = \frac{1}{(2\pi)^{n/2}} e^{-(\mathbf{x}'\mathbf{x})/2}$$
 (28)

Proof. See Appendix A. \Box

Spherical laws built from PHS distributions are, besides the multinormal, leptokurtic distributions. Even so, the Mardia kurtosis index that characterizes these distributions does not always resemble the value of the empirical kurtosis typically exhibited by multivariate financial series. As happens in the univariate case, the target of increasing the kurtosis index of SPHS laws can be accomplished via their polynomial expansions. The upcoming theorem is concerned with the problem of building GCLEs of SPHS laws and the kurtosis increase attainable with GCLE distributions. The corollary that follows provides closed-form expressions for the GCLE of the Gaussian, the logistic, and the hyperbolic-secant laws considered in Corollary 1.

Theorem 4. The density of a GCLE of the SPHS law as defined in (21) is:

$$\widetilde{g}_{n,\lambda}(\mathbf{x}) = q(\mathbf{x},\beta)g_{n,\lambda}(\mathbf{x})$$
(29)

where $q(\mathbf{x}, \beta)$ is as in (16) with coefficients determined as in (3) from (even) moments, m_{2i} defined as:

$$m_{2j} = \frac{M(n+2j,\lambda)}{a^{2j}M(n,\lambda)}$$
(30)

where M(.) is as in Equation (22), with a = 1 for the Gaussian case. Mardia's kurtosis index of the GCLE, as specified in (29), is given by:

$$K_{GCLE}^{SPHS} = K^{SPHS} + \beta = \frac{M(n+8,\lambda)}{a^8 M(n,\lambda)} + \beta$$
(31)

Proof. See Appendix A. \Box

Corollary 2. The GCLEs of the *n*-variate hyperbolic, logistic, and Gaussian density, henceforth denoted by $\tilde{g}_{n,\lambda=1}(\mathbf{x},\beta)$, $\tilde{g}_{n,\lambda=2}(\mathbf{x},\beta)$, and $\tilde{g}_{n,\lambda\to\infty}(\mathbf{x},\beta)$, respectively, are given by:

$$\tilde{g}_{n,\lambda=1}(\boldsymbol{x},\beta) = q(\boldsymbol{x},\beta) \frac{2^{n-2} \pi^{n/2}}{\phi(n)(\frac{n}{2})_{\frac{n}{2}}} \operatorname{sech}\left(\frac{\pi}{2} (\boldsymbol{x}'\boldsymbol{x})^{1/2}\right)$$
(32)

$$\tilde{g}_{n,\lambda=2}(\boldsymbol{x},\beta) = q(\boldsymbol{x},\beta)c_n sech^2\left(\frac{\pi}{2\sqrt{3}}(\boldsymbol{x}'\boldsymbol{x})^{1/2}\right)$$
(33)

$$\tilde{g}_{n,\lambda\to\infty}(\boldsymbol{x},\boldsymbol{\beta}) = q(\boldsymbol{x},\boldsymbol{\beta}) \frac{1}{(2\pi)^{n/2}} e^{-(\boldsymbol{x}'\boldsymbol{x})/2}$$
(34)

Here, $q(\mathbf{x}, \beta)$ *is a binomial with coefficients depending on moments specified as in* (30) *with* M(.) *defined as in* (A16), (A18) *and* (A22) *in Appendix A, for the hyperbolic-secant, the logistic, and the Gaussian, respectively.*

Proof. See Appendix A. \Box

3. Modeling Financial Series and Assessing Risk Measures via GCLEs

3.1. Data Description

The aim of this section is to evaluate the performance of the GCLEs in both fitting leptokurtic financial series and quantifying risk measures such as the Value at Risk (VaR) and the Expected Shortfall (ES). To this end, we considered a set of daily indexes: the Hang Seng Index (HSI), the 5-year U.S. Treasury Yield Index (FVX), and the CAC40index (FCHI), observed over a period that goes from 1 January 2009 to 31 December 2016, so as to cover the 2008 financial crisis and its aftermath; the data refer to three different geographical areas representing three different markets: the Chinese, the American, and the European one.

The dataset, including T = 1903 observations for each index (excluding missing cases), was split into two sub-samples of size $T_1 = 952$ and $T_2 = 951$, respectively. The former was employed to evaluate the goodness of fit of the GCLEs and their in-sample capability in assessing VaR and ES. The latter was used to evaluate the out-of-sample performance of said distributions in estimating the above risk measures.

For each series, the returns from data were computed as the log-ratio between daily closing prices at time t and t - 1. The preliminary statistics on the returns in the whole sample period are shown in Table 1.

Table 1. Number of observations (*T*), mean (μ), standard deviation (*sd*), skewness (*sk*) with Agostino's test significance, kurtosis (*ku*), and Jarque–Bera (JB) and Ljung–Box (Q²(20)) statistics of the three series. HSI, Hang Seng Index.

	Т	μ	sd	sk	ku	JB <i>p</i> -Value	$Q^2(20) p$ -Value
HSI	1903	0.0002	0.0134	-0.0432	5.155	< 0.001	< 0.001
FCHI	1903	0.0002	0.0145	-0.1263 *	6.482	< 0.001	< 0.001
FVX	1903	0.0001	0.0379	-0.0940*	5.421	< 0.001	< 0.001

* denotes statistical significance at 5% level, according to D'Agostino (1970). The JB test is based on Jarque and Bera (1987). The $Q^2(20)$ test is based on Ljung and Box (1978).

All the kurtosis indexes exceed three significantly, while the Jarque–Bera normality test statistics (Jarque and Bera 1987) are significant at the 1% level, thus providing evidence that the above returns are not normally distributed. Moreover, all the Ljung–Box $Q^2(20)$ statistics for the squared returns (Ljung and Box 1978) are significant at the 1% level. Finally, the D'Agostino test (D'Agostino 1970) signals skewness only at the 5% significance level for FCHI and FVX. As the returns are only mildly affected by skewness, the use of spherical distributions is suitable for their modeling.

3.2. Model Estimation and Fit

This sections explains how GCLEs can be fit to the empirical data described above. First of all, we modeled the three mentioned returns with a trivariate spherical law. More precisely, we considered a trivariate Gaussian, logistic, and hyperbolic-secant density.

According to Equations (24), (26) and (28) in Section 2, the spherical Gaussian, logistic, and hyperbolic-secant laws, denoted by g_N , g_L , and g_{HS} , are given by:

$$g_N(\mathbf{x}) = \frac{1}{(2\pi)^{3/2}} e^{-(\mathbf{x}'\mathbf{x})/2}$$
(35)

$$g_L(\mathbf{x}) = \frac{1}{8\sqrt{3}} \operatorname{sech}^2\left(\frac{\pi}{2\sqrt{3}} (\mathbf{x}'\mathbf{x})^{1/2}\right)$$
(36)

$$g_{HS}(\mathbf{x}) = \frac{1}{4\pi} \operatorname{sech}\left(\frac{\pi}{2} (\mathbf{x}'\mathbf{x})^{1/2}\right)$$
(37)

Taking into account Equations (32)–(34) in Corollary 2, the GCLE versions of these distributions, denoted by GCN, GCL, and GCHS respectively, are:

$$\tilde{g}_{GCN}(\mathbf{x}, \beta) = \left[1 + \frac{\beta}{120} \left((\mathbf{x}'\mathbf{x})^2 - 10(\mathbf{x}'\mathbf{x}) + 15 \right) \right] \frac{1}{(2\pi)^{3/2}} e^{-(\mathbf{x}'\mathbf{x})/2}$$
(38)

$$\tilde{g}_{GCL}(\mathbf{x},\beta) = \left[1 + \frac{\beta}{5129.5} \left((\mathbf{x}'\mathbf{x})^2 - 23.333(\mathbf{x}'\mathbf{x}) + 58.143 \right) \right] \frac{1}{8\sqrt{3}} \operatorname{sech}^2 \left(\frac{\pi}{2\sqrt{3}} (\mathbf{x}'\mathbf{x})^{1/2} \right)$$
(39)

$$\tilde{g}_{GCHS}(\mathbf{x}, \beta) = \left[1 + \frac{\beta}{14400} \left((\mathbf{x}'\mathbf{x})^2 - 30(\mathbf{x}'\mathbf{x}) + 89 \right) \right] \frac{1}{4\pi} \operatorname{sech} \left(\frac{\pi}{2} (\mathbf{x}'\mathbf{x})^{1/2} \right)$$
(40)

Here, β is the excess-kurtosis of the empirical trivariate series with respect to the (Mardia) kurtosis index (see Equation (12) in Appendix A) of the spherical Gaussian, logistic, and hyperbolic-secant laws, given in Equations (35)–(37).

By denoting with \tilde{K} the kurtosis of the empirical trivariate series and with K the (Mardia) kurtosis index of the spherical Gaussian, logistic, and hyperbolic-secant law that have been fit, in turn, to the empirical series, the excess kurtosis β can be estimated via the Method of Moments (MM) as follows:

$$\hat{\beta}_{\rm MM} = \tilde{K} - K \tag{41}$$

Alternatively, Maximum Likelihood (ML) estimation can be employed to estimate μ , Σ , and β from the data, numerically maximizing (via the optim function available in the R software, (R Core Team 2020)) the log-likelihood:

$$l(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \boldsymbol{\beta}) = \sum_{t=1}^{T} \log \left(\frac{\tilde{g}(\mathbf{z}_t, \boldsymbol{\beta})}{|\boldsymbol{\Sigma}|^{1/2}} \right)$$
(42)

where $\tilde{g}(.)$ is one of the trivariate densities in (38)–(40) and $\mathbf{z} = \mathbf{\Sigma}^{-1/2}(\mathbf{x} - \boldsymbol{\mu})$, thus obtaining the related $\hat{\beta}_{ML}$ estimate.

Table 2 reports the estimates of β , the AIC information criterion values for the GCLEs, together with those of the multivariate Student *t* (M*t*; Kotz and Nadarajah 2004) and the power exponential (MPE; Gómez et al. 1998) distributions, which have been fit to the series for comparison reasons. Furthermore, for the GCLEs, the levels of kurtosis of each parent distribution (K^{SPHS}) and of the associated expansions (K^{SPHS}_{GCLE}) are also reported. For the MM distribution, the latter clearly corresponds to the empirical Mardia's kurtosis index of the data.

Looking at the AIC results, when these criteria are applicable, we conclude that GCLEs fit better the empirical series than the other distributions, and this is particularly true for the GCHS.

Table 2. Estimates of β , the information criterion (AIC), and the estimated Mardia kurtosis indexes of the 3-variate series. For the Method of Moments (MM) estimates, K_{GCLE}^{SPHS} naturally corresponds to the empirical kurtosis in the data.

Series	Distribution	β	AIC	K ^{SPHS}	K_{GCLE}^{SPHS}
	GCN (ML)	11.565	-28,895.63	15.000	26.565
HSI	GCL (ML)	6.351	-29,978.95	20.335	26.686
	GCHS (ML)	5.603	-29,986.86	21.960	27.563
FCHI	GCN (MM)	11.432	-	15.000	26.432
	GCL (MM)	6.097	-	20.335	26.432
	GCHS (MM)	4.472	-	21.960	26.432
FVX	Mt	4.682 *	-29,611.20	-	-
	MPE	0.518 **	-29,919.72	-	-

* denotes the degrees of freedom of the Mt distribution. ** denotes the shape parameter of the MPE distribution.

In order to better investigate the goodness of fit of these polynomially adjusted densities in comparison with the popular M*t* and MPE distributions, the multivariate version of the Wilcoxon rank test (Liu and Singh 1993) was also employed. To implement the test, the following procedure was devised:

- An artificial population of one million data points was generated from each distribution, using the *β* estimates in Table 2. For the GCLE distributions, since no standard routine is available, the Hamiltonian Monte Carlo (HMC; Neal 2011) was employed;
- *B* = 1000 samples were extracted from each population. The MW test was performed via the R package DepthProc (Kosiorowski and Zawadzki 2014) on each of the samples coming from each population. This yields a *B*-dimensional vector containing the binary outcome of the tests (acceptance = 0/rejection = 1);
- A non-parametric bootstrap procedure was then performed on the above vector, to obtain a confidence interval on the proportion of rejections. The higher the proportion of rejections is, the worse the performance of the underlying model.

Table 3 shows the average *p*-value of the MW test across the *B* replicates, along with the proportion of rejections (p_R) and its 95% confidence interval (p_l , p_u), for each model. According to the test, a *p*-value higher than the significance threshold provides evidence that the two datasets come from the same underlying population. In our situation, one dataset represents a simulation from a given model (thus, it can be regarded as a proxy of the model itself), while the other represents the empirical data. This entails that the models with an average *p*-value above the significance threshold have a sufficiently adequate fit to the data. The results provide evidence that GCLEs perform well compared to M*t* and MPE, with the only exception of the GCN, since they are the only distributions with an average *p*-value above the 10% level. Looking at the results shown in the other columns of the table, we can see that GCLEs show also a much lower rate of rejection of the null hypothesis of the MW test across the *B* replications, except for the GCN. Furthermore, looking at both the average MW *p*-value and the rejection rate, we conclude that the GCHS distributions obtained via the method of moments estimation also adequately fit the data.

Distribution	Avg. <i>p</i> -Value	<i>p</i> _R	p_l	p_u
GCN (ML)	< 0.0001	1	1	1
GCL (ML)	0.5658	0.008	0.003	0.014
GCHS (ML)	0.5742	0.014	0.007	0.021
GCN (MM)	< 0.0001	1	1	1
GCL (MM)	0.1010	0.423	0.393	0.454
GCHS (MM)	0.4126	0.018	0.010	0.027
Mt	0.0025	0.993	0.987	0.998
MPE	< 0.0001	1	1	1

Table 3. Multivariate Wilcoxon test results. Average *p*-values, proportion of rejections in *B* replications (p_R) , and related non-parametric bootstrap confidence interval (p_l, p_u) .

3.3. Evaluation of VaR and ES via GCLEs

So far, we have evaluated the capability of the GCLEs to fit financial series. In the following, the validity of these densities in computing risk measures such as the value at risk and the expected shortfall is also investigated (see Chen 2018; McNeil et al. 2005), both in- and out-of-sample, for a linear portfolio:

$$P_z = \frac{1}{n} \sum_{i=1}^{n} Z_i$$
 (43)

To this end, the entire observation period, which ranges from 1 January 2009 to 31 December 2012 (1903 observations), was split into two sub-periods having approximately the same length: T_1 ,

going from 1 January 2009 to 31 December 2012 (952 observations), and T_2 , going from 1 January 2013 to 31 December 2016 (951 observations).

The observations of the period T_1 were used to obtain parameter estimates of the GCLEs, the M*t*, and the MPE law. Then, the latter were employed to estimate risk measures, such as the value at risk and the expected shortfall.

Given the spherical nature of the mentioned distributions, *VaR* and *ES* were estimated by using the approach proposed by Kamdem (2005). The value at risk at a given level α , VaR $_{\alpha}$, of a linear portfolio described by an *n*-dimensional elliptical random variable **z** with vector mean μ and covariance matrix **V**, was evaluated as:

$$VaR_{\alpha} = -\delta' \mu + q_{\alpha,n} \sqrt{\delta' \Sigma \delta}$$
(44)

Here, δ is a vector of weights, $\Sigma = \frac{n}{m_2} V$ with m_2 as defined in (17), and $q_{\alpha,n}$ is the unique positive solution of the following equation:

$$\alpha = \frac{\pi^{(n-1)/2}}{\Gamma(\frac{n-1}{2})} \int_{q_{\alpha,n}}^{\infty} \int_{v^2}^{\infty} (u - v^2)^{(n-3)/2} g_n(u) du dv.$$
(45)

In (45), $g_n(\cdot)$ is the density of the vector **z**, and $\Gamma(\cdot)$ denotes the gamma function. In the case of GCLEs of spherical variables and by setting equal weights $\delta_k = \frac{1}{n} \forall k$, Equation (44) simply becomes:

$$VaR_{\alpha} = \frac{q_{\alpha,n}}{\sqrt{m_2}} \tag{46}$$

where $q_{\alpha,n}$ is the solution of (45) obtained by replacing $g_n(u)$ with $\tilde{g}_n(\mathbf{x}, \beta)$ defined as in (15).

Following Kamdem (2005) and Nesmith et al. (2017), the expected shortfall at level α , ES_{α}, is given by:

$$\mathsf{ES}_{\alpha} = -\delta' \mu + \frac{\pi^{(n-1)/2}}{2\alpha\Gamma(\frac{n+1}{2})} \left(\int_{q_{\alpha,n}^2}^{\infty} (u - q_{\alpha,n}^2)^{(n-1)/2} g_n(u) du \right) |\delta' \Sigma \delta|^{1/2}$$
(47)

where the symbols are defined as in (44)–(46). In the case of GCLEs of spherical variables, by setting equal weights in δ and by replacing $g_n(u)$ in (47) with $\tilde{g}_n(u,\beta)$ as defined in (15), the above equation becomes:

$$\mathrm{ES}_{\alpha} = \frac{\pi^{(n-1)/2}}{2\sqrt{m_2}\alpha\Gamma(\frac{n+1}{2})} \left(\int_{q_{\alpha,n}^2}^{\infty} (u - q_{\alpha,n}^2)^{(n-1)/2} \tilde{g}_n(u,\beta) du \right)$$
(48)

Equation (48) was used to compute ES_{α} by using VaR_{α} estimated as in (44). In particular, GCLEs as defined in (38)–(40) were employed to compute ES_{α} as given in (48). The empirical counterparts of VaR and ES, $\text{VaR}_{\alpha,\text{emp}}$ and $\text{ES}_{\alpha,\text{emp}}$, respectively, of the linear portfolio P_z are given by:

$$\operatorname{VaR}_{\alpha,\operatorname{emp}} = -\inf_{z} \{ \hat{F}_{P_{z}} > \alpha \}$$

$$\tag{49}$$

$$\mathrm{ES}_{\alpha,\mathrm{emp}} = -\frac{\sum_{t=1}^{T} P_{zt} I(P_{zt} < -\mathrm{VaR}_{\alpha,\mathrm{emp}})}{\sum_{t=1}^{T} I(P_{zt} < -\mathrm{VaR}_{\alpha,\mathrm{emp}})}$$
(50)

where \hat{F}_{P_z} is the empirical cumulative distribution function of the portfolio defined as in Equation (43).

Table 4 compares the estimates of VaR_{α}, computed as in (46) by using GCN, GCL, and GCHS, with the corresponding VaR_{α ,emp} in the first sub-sample *T*₁. In this table, the percentile bootstrap intervals, $[V_l^{1p}, V_u^{1p}]$, for VaR^{1p}_{α ,emp} at the 95% confidence level are also reported, along with the absolute difference between VaR_{α} and VaR_{α ,emp}. These intervals were obtained by extracting 1000 bootstrap samples from the whitened data of the first period. To ensure preservation of the structure of the data, the maximum entropy bootstrap of the method of (Vinod 2006) was employed, via the R package

meboot (Vinod and López-de Lacalle 2009). A distribution whose VaR_{α} falls inside the 95% confidence interval of $VaR_{\alpha,emp}$ can be regarded as adequate for this risk measure.

Distribution	α	VaR _α	VaR ^{1p} _{emp}	V_l^{1p}	V_u^{1p}	$ VaR_{\alpha} - VaR_{emp}^{1p} $
	0.01	1.6713	1.5403	1.4304	1.6807	0.1310
CCN(MI)	0.025	1.2970	1.2073	1.1795	1.3111	0.0898
GCN (ML)	0.05	0.9044	0.9511	0.9243	1.0405	0.0466
	0.1	0.6307	0.6780	0.6510	0.7404	0.0473
	0.01	1.4689	1.5403	1.4304	1.6807	0.0714
CCL (MI)	0.025	1.1635	1.2073	1.1795	1.3111	0.0437
GCL (ML)	0.05	0.9322	0.9511	0.9243	1.0405	0.0189
	0.1	0.6939	0.6780	0.6510	0.7404	0.0159
	0.01	1.4950	1.5403	1.4304	1.6807	0.0453
CCHS (MI)	0.025	1.1788	1.2073	1.1795	1.3111	0.0285
GCI IS (IVIL)	0.05	0.9369	0.9511	0.9243	1.0405	0.0142
	0.1	0.6883	0.6780	0.6510	0.7404	0.0102
	0.01	1.6674	1.5403	1.4304	1.6807	0.1270
CCN (MM)	0.025	1.2930	1.2073	1.1795	1.3111	0.0857
	0.05	0.9057	0.9511	0.9243	1.0405	0.0454
	0.1	0.6326	0.6780	0.6510	0.7404	0.0454
	0.01	1.4705	1.5403	1.4304	1.6807	0.0698
CCL (MM)	0.025	1.1630	1.2073	1.1795	1.3111	0.0442
GCL (MINI)	0.05	0.9310	0.9511	0.9243	1.0405	0.0200
	0.1	0.6928	0.6780	0.6510	0.7404	0.0147
	0.01	1.4934	1.5403	1.4304	1.6807	0.0469
CCHS (MM)	0.025	1.1796	1.2073	1.1795	1.3111	0.0278
GCI IS (IVIIVI)	0.05	0.9383	0.9511	0.9243	1.0405	0.0128
	0.1	0.6896	0.6780	0.6510	0.7404	0.0115
	0.01	1.4675	1.5403	1.4304	1.6807	0.0728
Mt	0.025	1.1538	1.2073	1.1795	1.3111	0.0534
1411	0.05	0.9225	0.9511	0.9243	1.0405	0.0285
	0.1	0.6876	0.6780	0.6510	0.7404	0.0096
	0.01	1.4332	1.5403	1.4304	1.6807	0.1071
MPF	0.025	1.1661	1.2073	1.1795	1.3111	0.0412
1411 12	0.05	0.9493	0.9511	0.9243	1.0405	0.0018
	0.1	0.7145	0.6780	0.6510	0.7404	0.0365

Table 4. VaR_{α} estimated via the Mt, MPE, GCN, GCL, and GCHS, the empirical value at risk VaR^{1p}_{α,emp} in the period *T*₁, the 95% percentile bootstrap interval $[V_l^{1p}, V_u^{1p}]$, and the absolute difference between the estimated and the empirical VaR. Italic values denote a VaR_{α} falling outside the percentile interval.

Looking at the results shown in the table, we see that GCLEs manage to fall inside the confidence intervals across all α 's. In particular, the GCN is adequate for the two highest risk levels, while the GCL and the GCHS are adequate for the two lowest risk levels. Neither the M*t* nor the MPE follow a particular pattern and tend to underestimate the risk for $\alpha = 0.025$ and $\alpha = 0.05$. The only distribution that is able to estimate VaR_{α ,emp} satisfactorily across all levels of α is the GCHS estimated via the method of moments. Finally, looking at the results reported in the last column of the table, we can conclude that the GCHS distributions provide the closest fit to VaR_{α ,emp} for $\alpha = 0.01$ and $\alpha = 0.025$, while the MPE the closest fit for $\alpha = 0.05$ and the M*t* the closest fit for $\alpha = 0.1$, immediately followed by the GCHS.

A similar analysis was carried out for the expected shortfall.

Table 5 shows ES_{α} and $\text{ES}_{\alpha,\text{emp}}$ for the period T_1 together with the percentile bootstrap intervals, $\left[\text{E}_l^{1p}, \text{E}_u^{1p}\right]$, at the 95% confidence level. Looking at the results reported in the table, we see that the ES

is correctly estimated by GCLEs across all levels of α with the exception of the GCL, which slightly underestimates the expected VaR-exceeding losses for $\alpha = 0.025$; the same does not occur for both the Mt and the MPE densities, especially for $\alpha = 0.05$. Looking at the results provided in the last column of the table, we can conclude that GCHS distributions offer the best estimates of ES_{α ,emp} for all levels of α , except for $\alpha = 0.1$, which is better estimated by GCN.

In particular, the last two tables highlight how the most leptokurtic distribution, the GCHS, has a great advantage in estimating losses associated with high risk levels, thanks to a tail heaviness that is more pronounced relatively to the other GCLEs

Table 5. ES_{α} estimated via the M*t*, MPE, GCN, GCL, and GCHS, the empirical value at risk $\text{ES}_{\alpha,\text{emp}}^{1p}$ in the period *T*₁, the 95% percentile bootstrap interval $\left[E_l^{1p}, E_u^{1p}\right]$, and the absolute difference between the estimated and the empirical ES. Italic values denote a ES_{α} falling outside the percentile interval.

Distribution	α	ES _α	$\mathrm{ES}^{1p}_{\alpha,\mathrm{emp}}$	E_l^{1p}	E_u^{1p}	$ \mathrm{ES}_{\alpha} - \mathrm{ES}_{\alpha,\mathrm{emp}}^{1p} $
	0.01	1.9248	1.8420	1.7843	2.1157	0.0828
CCN (MI)	0.025	1.6532	1.5324	1.4977	1.7013	0.1208
GCN (ML)	0.05	1.3638	1.3032	1.2746	1.4267	0.0605
	0.1	1.054	1.0534	1.0249	1.1439	0.0006
	0.01	1.818	1.8420	1.7843	2.1157	0.0239
CCL (MI)	0.025	1.5029	1.5324	1.4977	1.7013	0.0295
GCL (ML)	0.05	1.2689	1.3032	1.2746	1.4267	0.0343
	0.1	1.0346	1.0534	1.0249	1.1439	0.0187
	0.01	1.8463	1.8420	1.7843	2.1157	0.0043
CCHS (MI)	0.025	1.5264	1.5324	1.4977	1.7013	0.0060
GCI IS (ML)	0.05	1.2854	1.3032	1.2746	1.4267	0.0178
	0.1	1.0423	1.0534	1.0249	1.1439	0.0111
	0.01	1.9215	1.8420	1.7843	2.1157	0.0795
CCN (MM)	0.025	1.6493	1.5324	1.4977	1.7013	0.1169
GCIN (IVIIVI)	0.05	1.3612	1.3032	1.2746	1.4267	0.0580
	0.1	1.0537	1.0534	1.0249	1.1439	0.0003
	0.01	1.8251	1.8420	1.7843	2.1157	0.0169
CCL (MM)	0.025	1.5058	1.5324	1.4977	1.7013	0.0266
GCL (WIWI)	0.05	1.2700	1.3032	1.2746	1.4267	0.0333
	0.1	1.0346	1.0534	1.0249	1.1439	0.0188
	0.01	1.8385	1.8420	1.7843	2.1157	0.0035
CCHS (MM)	0.025	1.5232	1.5324	1.4977	1.7013	0.0091
GCI IS (IVIIVI)	0.05	1.2843	1.3032	1.2746	1.4267	0.0189
	0.1	1.0424	1.0534	1.0249	1.1439	0.0109
	0.01	1.8543	1.8420	1.7843	2.1157	0.0123
Mt	0.025	1.5133	1.5324	1.4977	1.7013	0.0191
1411	0.05	1.2691	1.3032	1.2746	1.4267	0.0342
	0.1	1.0307	1.0534	1.0249	1.1439	0.0226
	0.01	1.7007	1.8420	1.7843	2.1157	0.1413
MPE	0.025	1.4487	1.5324	1.4977	1.7013	0.0837
	0.05	1.2477	1.3032	1.2746	1.4267	0.0555
	0.1	1.034	1.0534	1.0249	1.1439	0.0194

The out-of sample performance of the CGLE in estimating the value at risk was evaluated by using Kupiec's Likelihood Ratio test (LR; Kupiec 1995) and the Binomial Test (BT; Lee and Su 2012). Both the LR and BT null hypotheses state that the percentage of portfolio losses that in the second part of the sample exceed VaR_{α} is equal to α . Since VaR_{α} is estimated via GCLEs in the first period of the sample, this analysis should provide evidence of the out-of-sample stability of the GCLEs in assessing this risk measure.

The *p*-values of both the LR (*p*-value LR) and binomial test (*p*-value BT) are shown in Table 6 together with two loss measures: the Average Binary Loss Function (ABLF) and the Average Quadratic Loss Function (AQLF). The former is the average of the Binary Loss (BL) function, which gives a penalty of one when each return r_t exceeds the VaR, without concern for the magnitude of the excess:

$$BL = \begin{cases} 1 & \text{if } r_t \ge \text{VaR} \\ 0 & \text{otherwise} \end{cases}$$
(51)

If a VaR model truly provides the level of coverage defined by its confidence level α , then the ABLF should be as close as possible to the latter.

The AQLF (Lopez 1999) is the average of the Quadratic Loss (QL) function:

$$QL = \begin{cases} 1 + (r_t - VaR)^2 & \text{if } r_t \ge \text{VaR} \\ 0 & \text{otherwise} \end{cases}$$
(52)

and pays more attention to the magnitude of the violations, as large violations are more penalized than the small ones.

Table 6 shows the results obtained by performing the LR and the BN test. They confirm the good performance offered by GCLEs, with the exception of the GCN. Looking at the loss functions, we can conclude that the GCHS outperforms all the other distributions for $\alpha = 0.025$. In particular, the GCL and the M*t* are equivalent for $\alpha = 0.01$; the GCN (ML) and the GCHS (MM) provide the best fitting for $\alpha = 0.05$ (the former for the ABLF, the latter for the AQLF); and the GCN is the only distribution not underestimating the ABLF for $\alpha = 0.1$, which is better approximated by the MPE according to the AQLF.

Finally, the out-of-sample GCLE performance in estimating the ES was evaluated by performing the Z_1 and Z_2 tests of Acerbi and Szekely (2014). The null hypothesis of both tests assumes that the estimated ES_{α} tallies with the ES calculated from the data. This entails, in the present application, that ES_{α} should provide a good estimate of the empirical expected shortfall computed in the second part of the sample. Under the alternative, this does not happen, and ES_{α} systematically underestimates the effective loss mean, ES_{emp} , thus implying financial damage.

The first test is based on the following statistic:

$$Z_1 = \frac{1}{N_T} \sum_{t=1}^T \left(\frac{L_t I_t > VaR_\alpha}{ES_\alpha} \right) - 1$$
(53)

where $N_T = \sum_{t=1}^{T} I_t$ is the number of losses L_t , which in the second part of the sample, break the threshold VaR_{α}, and I_t is an indicator variable that is equal to one if $L_t > VaR_{\alpha}$ and zero otherwise.

As observed by Acerbi and Szekely (2014), this test is completely insensitive to an excessive number of exceptions as it is simply an average taken over the exceptions themselves. Under the null, the realized value Z_1 is expected to be zero, and it signals a problem when it displays values significantly greater than zero.

The second test statistic is:

$$Z_2 = \frac{1}{T} \sum_{t=1}^{T} \frac{L_t I_{L_t > VaR_\alpha}}{\alpha ES_\alpha} - 1$$
(54)

where *T* denotes the sample size of the second period, and the other symbols are defined as above. As for the previous test, under the null, the realized value Z_2 is expected to be zero, and it signals a problem when it assumes values significantly greater than zero.

Distribution	α	ABLF	AQLF	<i>p</i> -Value LR	<i>p</i> -Value BT
	0.01	0.0032	0.0032	0.0132	0.0318
CCN (MI)	0.025	0.0147	0.016	0.0281	0.0472
GCN (ML)	0.05	0.0494	0.0575	0.9347	0.9999
	0.1	0.1136	0.1351	0.1713	0.1601
	0.01	0.0105	0.0109	0.8741	0.8695
CCL (MI)	0.025	0.0242	0.0269	0.8714	0.9999
GCL (ML)	0.05	0.0442	0.0515	0.3999	0.4567
	0.1	0.0915	0.1089	0.3750	0.4174
	0.01	0.0074	0.0077	0.3907	0.5146
CCUS (MI)	0.025	0.0242	0.0267	0.8714	0.9999
GCI IS (ML)	0.05	0.0442	0.0513	0.3999	0.4567
	0.1	0.0915	0.1092	0.3750	0.4174
	0.01	0.0032	0.0032	0.0131	0.0318
	0.025	0.0147	0.0161	0.0281	0.0472
GCIN (IVIIVI)	0.05	0.0484	0.0565	0.8167	0.8818
	0.1	0.1136	0.135	0.1713	0.1601
	0.01	0.0105	0.0109	0.8741	0.8695
	0.025	0.0242	0.0269	0.8714	0.9999
GCL (IVIIVI)	0.05	0.0442	0.0515	0.3999	0.4567
	0.1	0.0915	0.1089	0.3750	0.4174
	0.01	0.0084	0.0087	0.6129	0.7453
	0.025	0.0231	0.0256	0.7089	0.8351
GCH5 (MINI)	0.05	0.0431	0.0502	0.3187	0.3716
	0.1	0.0915	0.1091	0.3750	0.4174
	0.01	0.0105	0.0109	0.8741	0.8695
M4	0.025	0.0242	0.0270	0.8714	0.9999
IVIL	0.05	0.0452	0.0528	0.4917	0.5517
	0.1	0.0925	0.1103	0.4376	0.4822
	0.01	0.0105	0.0110	0.8741	0.8695
MPE	0.025	0.0242	0.0268	0.8714	0.9999
IVII E	0.05	0.0431	0.0499	0.3187	0.3716
	0.1	0.0862	0.1024	0.1480	0.1763

Table 6. The Average Quadratic Loss Function (AQLF) and the Average Binary Loss Function (ABLF) computed in the out-of-sample period T_2 and the *p*-values of the likelihood ratio and binomial tests.

To perform the test and obtain the associated *p*-values, the following parametric bootstrap approach was used:

- 1. A population of one million observations was generated from each distribution under investigation (M*t*, MPE, or one of the GCLEs considered in the paper), with parameters estimated by using the data of the first sample period. For the class of the GCLEs, HMC was employed to this aim;
- 2. B = 5000 samples, of a size equal to the dimension of the out-of-sample dataset, were drawn from each of these populations;
- 3. the bootstrap distributions of the statistics Z_1 and Z_2 were obtained and used to calculate the *p*-values, p_{Boot} , of the statistics z_1 and z_2 computed by using the observed sample data:

$$p_{Boot} = \frac{\sum_{b=1}^{B} I_{z_{i,b} > z_i}}{B}, \quad i = 1, 2.$$
(55)

The results are reported in Table 7. They allow concluding that the GCLEs adequately estimate the ES, with the exception of the GCN. According to the Z_1 statistic, the Mt distribution underestimates the expected losses for $\alpha = 0.01$, while the MPE underestimates the ES according to the Z_2 statistic for

 α = 0.1. As in the case of the in-sample assessment, distributions that are appreciably more leptokurtic than the GCN show a better tail sensitivity and are thus particularly adequate to evaluate risk measures at high levels of α .

Distribution	α	Z_1	<i>p</i> -Value Z ₁	Z_2	<i>p</i> -Value Z ₂
	0.01	-0.0836	0.0042	-0.7109	0.0022
CCN(MI)	0.025	-0.0575	0.0366	-0.4450	0.0032
GCIN (IVIL)	0.05	-0.1065	0.0010	-0.1169	0.1860
	0.1	-0.0971	0.0026	0.0254	0.6784
	0.01	-0.1068	0.0506	-0.0609	0.5502
CCL (MI)	0.025	-0.0454	0.2248	-0.0765	0.4508
GCL (ML)	0.05	-0.0113	0.5132	-0.1267	0.2134
	0.1	-0.0130	0.4572	-0.0971	0.1634
	0.01	-0.0873	0.0748	-0.3282	0.1578
CCHS (ML)	0.025	-0.0601	0.1156	-0.0907	0.3646
GCI IS (ML)	0.05	-0.0240	0.2976	-0.1379	0.1826
	0.1	-0.0203	0.3330	-0.1037	0.1566
	0.01	-0.0820	0.0056	-0.7104	0.0028
CCN (MM)	0.025	-0.0553	0.0524	-0.4437	0.0036
GCIV (IVIIVI)	0.05	-0.0998	0.0014	-0.1292	0.1562
	0.1	-0.0968	0.0018	0.0257	0.6428
	0.01	-0.1104	0.0286	-0.0646	0.4832
CCL (MM)	0.025	-0.0472	0.1860	-0.0783	0.3570
GCL (IVIIVI)	0.05	-0.0121	0.4472	-0.1274	0.1428
	0.1	-0.0129	0.3952	-0.0970	0.1330
	0.01	-0.0964	0.0394	-0.2399	0.2256
CCHS (MM)	0.025	-0.0505	0.1614	-0.1214	0.2264
GCI IS (IVIIVI)	0.05	-0.0171	0.3980	-0.1526	0.0706
	0.1	-0.0204	0.2472	-0.1039	0.0786
	0.01	-0.1244	0.0178	-0.0793	0.3936
Mt	0.025	-0.0519	0.1524	-0.0829	0.2794
1011	0.05	-0.0174	0.3370	-0.1115	0.1642
	0.1	-0.0130	0.3208	-0.0867	0.1504
	0.01	-0.0453	0.1736	0.0040	0.5074
MPF	0.025	-0.0096	0.3998	-0.0419	0.4108
1911 L	0.05	0.0116	0.6566	-0.1277	0.1296
	0.1	0.0063	0.5998	-0.1323	0.0458

Table 7. Acerbi and Szekely (2014) tests results. Z₁ and Z₂ statistics and related bootstrap *p*-values.

4. Conclusions

In this paper, we extend the Gram–Charlier-Like Expansion (GCLE) based approach to modify the kurtosis of multivariate distributions. We obtain a class of multivariate leptokurtic distributions, called GCLEs, which prove suitable to model the excess kurtosis typically found in financial data. GCLEs are tail sensitive densities and potentially fit well the tails of the empirical distributions of multivariate financial returns. As such, they can be conveniently used to compute risks measures like the Value at Risk (VaR) and the Expected Shortfall (ES). An application of GCLEs to a portfolio of a set of financial assets provides evidence of their effectiveness in computing the mentioned risk measures both in- and out-of-sample. Furthermore, a comparison of GCLEs with other commonly employed spherical distributions such as the Mt and the MPE show that the former can yield a comparable, if not better performance with respect to the latter, both in fitting the data and in estimating risk measures. Generally, we find no systematic reason to prefer the MM estimation to ML, barring computational convenience. The paper opens up a potentially fruitful line of research providing a class of multivariate distributions that are of interest in applications involving the modeling of heavy-tailed distributions. The results thus far obtained suggest that GCLEs can be an interesting tool for risk management. This paper investigated their potential in modeling leptokurtic data that do not need any significant adjustment for skewness. In fact, the GCLEs being spherical laws, they are not suitable to account for skewness. In the context of the polynomial expansion based approach for modeling data, skewness could be dealt with by using as starting distributions other laws than the spherical ones. Thus, supplementary research could include the consideration of other moments (e.g., skewness and volatility dynamics) and the investigation of the GCLEs' performance in other financial applications such as asset pricing or risk forecasting.

Author Contributions: Conceptualization, M.G.Z.; methodology, M.G.Z.; software, L.B. and G.V.; investigation, G.V.; data curation, L.B. and G.V.; writing, original draft preparation, M.G.Z.; writing, review and editing, M.Z. and G.V.; visualization, G.V. and L.B.; supervision, M.G.Z. All authors read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

GCLE	Gram–Charlier-Like Expansion
GCE	Gram–Charlier Expansion
SPHS	Spherical Power-raised Hyperbolic Secant
GCN	Gram–Charlier Normal
GCL	Gram–Charlier Logistic
GCHS	Gram–Charlier Hyperbolic Secant
Mt	Multivariate <i>t</i>
MPE	Multivariate Power Exponential
ML	Maximum Likelihood
MM	Method of Moments
MW	Multivariate Wilcoxon
ABLF	Average Binary Loss Function
AQLF	Average Quadratic Loss Function
LR	Likelihood Ratio
BT	Binomial Test
VaR	Value at Risk
ES	Expected Shortfall
HMC	Hamiltonian Monte Carlo

Appendix A. Proof of Relevant Theorems

Proof of Theorem 1. For the proof of (8)–(11), see Fang et al. (2002). The proofs of (12) and (13) can be found in Gómez et al. (2003). As far as (13) is concerned, let us consider the following identity given in Fang et al. (2002), ibidem, on page 35:

$$\int_{\mathbb{R}^n} g(||\mathbf{x}||^2) d\mathbf{x} = \frac{1}{2k_n} \int_0^\infty r^{n/2-1} g(r) dr = \frac{M(n)}{k_n}$$
(A1)

where $r = ||\mathbf{x}|| = (\mathbf{x}'\mathbf{x})^{1/2}$ and M(n) is as defined in (10).

Equation (A1) entails:

$$\int_{\mathbb{R}^n} ||\mathbf{x}||^{2j} g(||\mathbf{x}||^2) d\mathbf{x} = \frac{1}{2k_n} \int_0^\infty r^{(2j+n)/2 - 1} g(r) dr = \frac{M(n+2j)}{k_n}$$
(A2)

which, in turn, taking into account (8), implies what follows:

$$\int_{\mathbb{R}^n} (\mathbf{x}'\mathbf{x})^j g_n(\mathbf{x}) d\mathbf{x} = \frac{1}{M(n)} \int_{\mathbb{R}^n} ||\mathbf{x}||^{2j} g(||\mathbf{x}||^2) d\mathbf{x} = \frac{M(n+2j)}{M(n)} = m_{2j}.$$
 (A3)

As for (12), it must be considered that spherical variables are the standardized form of elliptical variables and that the Mardia kurtosis index, *K*, of elliptical variables is defined as follows (see Gómez et al. 2003):

$$K = n^2 \frac{m_4}{m_2^2}$$
(A4)

Equation (A4) simplifies into (12) in the case of spherical variables, as simple computations prove. To obtain this, let us observe that a vector \mathbf{y} of elliptical variables has a density of the form:

$$\mathbf{y} \approx |\mathbf{\Sigma}|^{-1/2} g \left[(\mathbf{y} - \boldsymbol{\mu})' \mathbf{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu}) \right]$$
(A5)

where μ is the mean vector and Σ is an $n \times n$ matrix related to the variance-covariance matrix of **y** by the the relationship:

$$\mathbb{V}(\mathbf{y}) = \frac{n}{\mathbb{E}(R^2)} \mathbf{\Sigma}$$
(A6)

Then, according to (A6), $\mathbb{V}(\mathbf{x}) = \mathbf{\Sigma} = \mathbf{I}$, as occurs in the spherical case, if and only if $n = \mathbb{E}(R^2) = m_2$. Substituting this result into (A4) yields (12). \Box

Proof of Theorem 2. Simple computations prove that the density $g_n(\mathbf{x})$ can be also expressed as follows:

$$g_n(\mathbf{x}) = k_n(\mathbf{x}'\mathbf{x})^{(1-n)/2}h((\mathbf{x}'\mathbf{x})^{1/2})$$
(A7)

where $h(\cdot)$ is the density of the generating variate *R* (see (11)).

Now, adjusting $h(\cdot)$ with a binomial specified as in (1) with $p_4(.)$ defined as in (16) gives:

$$\tilde{h}((\mathbf{x}'\mathbf{x})^{1/2}) = q(\mathbf{x},\beta)h((\mathbf{x}'\mathbf{x})^{1/2})$$
(A8)

where $\tilde{h}((\mathbf{x}'\mathbf{x})^{1/2})$ is the density of a generating variate, \tilde{R} , whose fourth moment, \tilde{m}_4 , is increased by a quantity β with respect to the same moment, m_4 , of the generating variate, R, of the parent density $h((\mathbf{x}'\mathbf{x})^{1/2})$, that is (see Nicolussi and Zoia 2020):

$$E(\tilde{R}^4) = E(R^4) + \beta = m_4 + \beta \tag{A9}$$

The latter, according to (13), can be also expressed as follows:

$$m_4 = \frac{M(n+4)}{M(n)} + \beta \tag{A10}$$

Simple computations prove that the density of the spherical variable having $\tilde{h}((x'x)^{1/2})$ as the generating variate:

$$\tilde{g}_n(\mathbf{x}) = k_n(\mathbf{x}'\mathbf{x})^{(1-n)/2} \tilde{h}((\mathbf{x}'\mathbf{x})^{1/2})$$
(A11)

tallies with that of the GCLE, $g_n(\mathbf{x})$, given in (15). Moreover, in light of (12), Mardia's kurtosis index of $\tilde{g}_n(\mathbf{x})$ is equal to the fourth moment of \tilde{R} , which proves (18).

Proof of Theorem 3. With:

$$g_{\lambda}(r^2) = [\operatorname{sech}(ar)]^{\lambda}, \quad \lambda, r > 0$$
(A12)

playing the role of the density generator, Equation (10) after the change of variable v = ar becomes:

$$M(n) = \int_0^\infty r^{n-1} \operatorname{sech}(ar)^\lambda dr = \frac{1}{a^n} M(n,\lambda)$$
(A13)

where $M(n, \lambda)$ is defined as in (22). This together with (A12) leads to (21) in light of (8). Now, note that in light of (13) and (A13), the 2*j*-th moment of the generating variate, *R*, of an *SPHS* variable is given by:

$$m_{2j} = \frac{M(n+2j,\lambda)}{a^{2j}M(n,\lambda)}, \qquad j = 1, 2, \dots$$
 (A14)

This, together with (12), yields (23) \Box

Proof of Corollary 1. When $\lambda = 1$, the parameters *a* and *b* of Equation (20) take the values:

$$a = \frac{\pi}{2}, \quad b = \frac{1}{2}$$
 (A15)

and accordingly, the density $g_{\lambda}(y)$ of Equation (19) becomes a univariate hyperbolic-secant. The Mellin transform (22), necessary to obtain its spherical extension, is (see Oberhettinger 1974, p.61):

$$M(n,1) = \int_0^\infty r^{n/2-1} \operatorname{sech}\left(\frac{\pi}{2}r^{1/2}\right) dr = \frac{\phi(n)\Gamma(n)}{(4)^{n-1}} \left(\frac{2}{\pi}\right)^n$$
(A16)

This, taking into account Equation (21), leads to (24).

When $\lambda = 2$, the parameters *a* and *b* of Equation (20) take the values:

$$a = \frac{\pi}{2\sqrt{3}}, \quad b = \frac{\pi}{4\sqrt{3}} \tag{A17}$$

and accordingly, the density $g_{\lambda}(y)$ of Equation (19) becomes a univariate logistic. The Mellin transform (22), necessary to obtain its spherical extension, is (see Gradshteyn and Ryzhik 2014, p. 352):

$$M(n,2) = \int_0^\infty r^{n/2-1} \operatorname{sech}^2\left(\frac{\pi}{2\sqrt{3}}r^{1/2}\right) dr =$$
(A18)

$$= \begin{cases} \frac{24}{\pi^2} \ln 2 & \text{if } n = 2\\ 8(1 - 2^{2-n})\Gamma(n)\zeta(n-1)\left(\frac{\sqrt{3}}{\pi}\right)^n & \text{otherwise} \end{cases}$$
(A19)

and accordingly, Equation (21) specifies into (26).

Eventually, as $\lambda \to \infty$, the following holds for the density generator of the *PHS* density (see Faliva and Zoia 2013).

$$g_{\lambda \to \infty}(r) = e^{-r/2} \tag{A20}$$

Furthermore, in light of the following integral transform (see Gradshteyn and Ryzhik 2014, p. 317):

$$\int_{0}^{\infty} v^{\tau - 1} e^{-sv/2} dv = \frac{1}{s^{\tau}} \Gamma(\tau), \quad s > 0$$
 (A21)

the following Mellin transform:

$$M(n,\infty) = \int_0^\infty r^{n-1} e^{-\frac{r^2}{2}} dr = 2^{n/2-1} \Gamma\left(\frac{n}{2}\right)$$
(A22)

holds. Under these premises, Equation (28) follows from (8). \Box

Proof of Theorem 4. The proof of (29) ensues from (15) by taking $g_n(x) = g_{n,\lambda}(x)$. Equation (30) comes from (A14). Finally, Equation (31) ensues from (18) bearing in mind (23).

Proof of Corollary 2. Equations (32)–(34) are obtained by replacing in (29) $g_n(\mathbf{x})$ with the density of a spherical hyperbolic-secant, logistic, and Gaussian law, respectively. The coefficients of the binomial $q(x, \beta)$ appearing in (32)–(34) are obtained as in (A14) by using (A16), (A18) and (A22) for the hyperbolic-secant, the logistic, and the Gaussian, respectively. \Box

References

Acerbi, Carlo, and Balazs Szekely. 2014. Back-testing expected shortfall. Risk 27: 76-81.

- Bagnato, Luca, Antonio Punzo, and Maria G. Zoia. 2017. The multivariate leptokurtic-normal distribution and its application in model-based clustering. *Canadian Journal of Statistics* 45: 95–119. [CrossRef]
- Bagnato, Luca, Valerio Potì, and Maria G. Zoia. 2015. The role of orthogonal polynomials in adjusting hyperbolic secant and logistic distributions to analyse financial asset returns. *Statistical Papers* 56: 1205–34. [CrossRef]
- Berkowitz, S., and F. J. Garner. 1970. The calculation of multidimensional hermite polynomials and gram-charlier coefficients. *Mathematics of Computation* 24: 537–55. [CrossRef]
- Boudt, Kris, Brian G. Peterson, and Christophe Croux. 2007. Estimation and decomposition of downside risk for portfolios with non-normal returns. *Journal of Risk* 11: 79–103. [CrossRef]
- Chen, James Ming. 2018. On exactitude in financial regulation: Value-at-risk, expected shortfall, and expectiles. *Risks* 6: 61. [CrossRef]
- D'Agostino, Ralph B. 1970. Transformation to normality of the null distribution of g1. *Biometrika* 57: 679–81. [CrossRef]
- Del Brio, Esther B., Trino-Manuel Niguez, and Javier Perote. 2009. Gram–charlier densities: A multivariate approach. *Quantitative Finance* 9: 855–68. [CrossRef]
- Engel, Eduardo M. R. A., Ronald D. Fischer, and Alexander Galetovic. 2010. The economics of infrastructure finance: Public-private partnerships versus public provision. *EIB Papers* 15: 40–69.
- Eun, Cheol S., Bruce Resnick, and Sanjiv Sabherwal. 2004. *International Financial Management*. New York: McGraw-Hill/Irwin.
- Faliva, Mario, and Maria G. Zoia. 2013. The family of power-raised hyperbolic-secant distributions: moments, kurtosis, limit laws. *Contributi di Ricerca in Econometria e Matematica* 14: 1–16.
- Faliva, Mario, and Maria G. Zoia. 2017. A distribution family bridging the gaussian and the laplace laws, gram–charlier expansions, kurtosis behaviour, and entropy features. *Entropy* 19: 149. [CrossRef]
- Faliva, Mario, Valerio Potì, and Maria G. Zoia. 2016. Orthogonal polynomials for tailoring density functions to excess kurtosis, asymmetry and dependence. *Communications in Statistics-Theory and Methods* 45: 49–62. [CrossRef]
- Fang, Hong-Bin, Kai-Tai Fang, and Samuel Kotz. 2002. The meta-elliptical distributions with given marginals. *Journal of Multivariate Analysis* 82: 1–16. [CrossRef]
- Gómez, Eusebio, Miguel A. Gómez-Villegas, and J. Miguel Marín. 1998. A multivariate generalization of the power exponential family of distributions. *Communications in Statistics Theory and Methods* 27: 589–600. [CrossRef]
- Gómez, Eusebio, Miguel A. Gómez-Villegas, and J. Miguel Marín. 2003. A survey on continuous elliptical vector distributions. *Revista Matemática Complutense* 16: 345–61. [CrossRef]
- Gradshteyn, Izrail S., and Iosif M. Ryzhik. 2014. Table of Integrals, Series, and Products. Cambridge: Academic Press.
- Harmantzis, Fotios C., Linyan Miao, and Yifan Chien. 2006. Empirical study of value-at-risk and expected shortfall models with heavy tails. *The Journal of Risk Finance* 7: 117–35. [CrossRef]
- Jarque, Carlos M., and Anil K. Bera. 1987. A test for normality of observations and regression residuals. International Statistical Review 55: 163–72. [CrossRef]
- Jondeau, Eric, and Michael Rockinger. 2001. Gram-charlier densities. *Journal of Economic Dynamics and Control* 25: 1457–83. [CrossRef]
- Jondeau, Eric, and Michael Rockinger. 2005. Conditional asset allocation under non-normality: How costly is the mean-variance criterion? *EFA 2005 Moscow Meetings Paper*. Available online: http://dx.doi.org/10.2139/ssrn.674424 (accessed on 1 October 2020).

- Jondeau, Eric, and Michael Rockinger. 2006a. The impact of news on higher moments. *Swiss Finance Institute Research Paper*. Available online: http://dx.doi.org/10.2139/ssrn.947082 (accessed on 1 October 2020). .
- Jondeau, Eric, and Michael Rockinger. 2006b. Optimal portfolio allocation under higher moments. *European Financial Management* 12: 29–55. [CrossRef]
- Jondeau, Eric, and Michael Rockinger. 2007. The economic value of distributional timing. *Swiss Finance Institute Research Paper*. Available online: http://dx.doi.org/10.2139/ssrn.957514 (accessed on 1 October 2020).
- Jurczenko, Emmanuel, Bertrand Maillet, and Bogdan Negréa. 2004. A note on skewness and kurtosis adjusted option pricing models under the martingale restriction. *Quantitative Finance* 4: 479–88.
- Kamdem, Jules S. 2005. Value-at-risk and expected shortfall for linear portfolios with elliptically distributed risk factors. *International Journal of Theoretical and Applied Finance* 8: 537–51. [CrossRef]
- Kosiorowski, Daniel, and Zygmunt Zawadzki. 2014. Depthproc: An R package for robust exploration of multidimensional economic phenomena. *arXiv* arXiv:1408.4542.
- Kotz, Samuel, and Saraleed Nadarajah. 2004. *Multivariate t-Distributions and Their Applications*. Cambridge: Cambridge University Press.
- Kozubowski, Tomasz J., Krzysztof Podgórski, and Igor Rychlik. 2013. Multivariate generalized laplace distribution and related random fields. *Journal of Multivariate Analysis* 113: 59–72. [CrossRef]
- Kupiec, Paul. 1995. Techniques for verifying the accuracy of risk measurement models. *The Journal of Derivatives* 3: 73–84. [CrossRef]
- Landsman, Zinoviy M., and Emiliano A. Valdez. 2003. Tail conditional expectations for elliptical distributions. North American Actuarial Journal 7: 55–71. [CrossRef]
- Lee, Cheng-Few, and Jung-Bin Su. 2012. Alternative statistical distributions for estimating value-at-risk: theory and evidence. *Review of Quantitative Finance and Accounting* 39: 309–31. [CrossRef]
- Liu, Regina Y., and Kesar Singh. 1993. A quality index based on data depth and multivariate rank tests. *Journal of the American Statistical Association* 88: 252–60.
- Ljung, Greta M., and George E.P. Box. 1978. On a measure of lack of fit in time series models. *Biometrika* 65: 297–303. [CrossRef]
- Lopez, Jose A. 1999. *Methods for Evaluating Value-at-Risk Estimates*. San Francisco: Economic Review-Federal Reserve Bank of San Francisco, vol. 3.
- Madan, Dilip B. 2020. Multivariate distributions for financial returns. *International Journal of Theoretical and Applied Finance*. [CrossRef]
- Mandelbrot, Benoit B. 1997. The variation of certain speculative prices. In *Fractals and Scaling in Finance*. New York: Springer, pp. 371–418.
- Mardia, Kanti V. 1970. Measures of multivariate skewness and kurtosis with applications. *Biometrika* 57: 519–30. [CrossRef]
- Marinelli, Carlo, Stefano d'Addona, and Svetlozar T. Rachev. 2012. Multivariate heavy-tailed models for value-at-risk estimation. *International Journal of Theoretical and Applied Finance* 15: 1250029. [CrossRef]
- McNeil, Alexander J., Rüdiger Frey, and Paul Embrechts. 2005. *Quantitative Risk Management: Concepts, Techniques and Tools*. Princeton: Princeton University Press, ISBN 978-0-691-12255-7.
- Neal, Radford M. 2011. MCMC using hamiltonian dynamics. In *Handbook of Markov ChainMonte Carlo*. Boca Raton: Chapman & Hall/CRC, chp. 5, pp. 113–62.
- Nesmith, Travis D., Dong H. Oh, and Dobrislav Dobrev. 2017. Accurate evaluation of expected shortfall for linear portfolios with elliptically distributed risk factors. *Journal of Risk and Financial Management* 10: 5.
- Nicolussi, Federica, and Maria G. Zoia. 2020. Gram–charlier-like expansions of the convoluted hyperbolic-secant density. *Journal of Statistical Theory and Practice* 14: 15. [CrossRef]
- Oberhettinger, Fritz. 1974. Tables of Mellin Transforms. New York: Springer.
- R Core Team. 2020. *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Rachev, Svetlozar T. 2003. *Handbook of Heavy Tailed Distributions in Finance: Handbooks in Finance, Book 1.* New York: Elsevier.
- Szegö, Giorgio. 2004. Risk Measures for the 21st Century. New York: Wiley.
- Vaughan, David C. 2002. The generalized secant hyperbolic distribution and its properties. *Communications in Statistics-Theory and Methods* 31: 219–38. [CrossRef]

- Vinod, Hrishikesh D., and Javier López-de Lacalle. 2009. Maximum entropy bootstrap for time series: The meboot r package. *Journal of Statistical Software* 29: 1–19. [CrossRef]
- Vinod, Hrishikesh D. 2006. Maximum entropy ensembles for time series inference in economics. *Journal of Asian Economics* 17: 955–78. [CrossRef]
- Weng, Ruby C. 2015. Expansions for multivariate densities. *Journal of Statistical Planning and Inference* 167: 174–81. [CrossRef]
- Zayed, Ahmed I. 1996. *Handbook of Function and Generalized Function Transformations*. Boca Raton: Chapman & Hall/CRC.
- Zoia, Maria G. 2010. Tailoring the gaussian law for excess kurtosis and skewness by hermite polynomials. *Communications in Statistics-Theory and Methods* 39: 52–64. [CrossRef]

Sample Availability: The data used for this paper is available online, on the Yahoo®Finance repository. It has been downloaded via the R statistical software, using the tseries package.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).