

A Service of

**ZBW** 

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Martinovic, John

# **Article — Published Version** A note on the integrality gap of cutting and skiving stock instances

4OR

# **Provided in Cooperation with:** Springer Nature

*Suggested Citation:* Martinovic, John (2020) : A note on the integrality gap of cutting and skiving stock instances, 4OR, ISSN 1614-2411, Springer, Berlin, Heidelberg, Vol. 20, Iss. 1, pp. 85-104, [https://doi.org/10.1007/s10288-020-00469-4](https://doi.org/10.1007/s10288-020-00469-4%0A)

This Version is available at: <https://hdl.handle.net/10419/288456>

#### **Standard-Nutzungsbedingungen:**

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.



<https://creativecommons.org/licenses/by/4.0/>

#### **Terms of use:**

*Documents in EconStor may be saved and copied for your personal and scholarly purposes.*

*You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.*

*If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.*



# WWW.ECONSTOR.EU

**RESEARCH PAPER**



# **A note on the integrality gap of cutting and skiving stock instances**

**Why 4***/***3 is an upper bound for the divisible case?**

**John Martinovic[1](http://orcid.org/0000-0002-5428-9351)**

Received: 31 March 2020 / Revised: 23 November 2020 / Accepted: 3 December 2020 / Published online: 23 December 2020 © The Author(s) 2020

# **Abstract**

In this paper, we consider the (additive integrality) gap of the cutting stock problem (CSP) and the skiving stock problem (SSP). Formally, the gap is defined as the difference between the optimal values of the ILP and its LP relaxation. For both, the CSP and the SSP, this gap is known to be bounded by 2 if, for a given instance, the bin size is an integer multiple of any item size, hereinafter referred to as the divisible case. In recent years, some improvements of this upper bound have been proposed. More precisely, the constants 3/2 and 7/5 have been obtained for the SSP and the CSP, respectively, the latter of which has never been published in English language. In this article, we introduce two reduction strategies to significantly restrict the number of representative instances which have to be dealt with. Based on these observations, we derive the new and improved upper bound 4/3 for both problems under consideration.

**Keywords** Cutting and packing · Cutting stock problem · Skiving stock problem · Additive integrality gap · Divisible case

**Mathematics Subject Classification** 90C10 · 90C27

# **1 General introduction**

Let a capacity (bin size)  $L \in \mathbb{N}$  and  $m \in \mathbb{N}$  items, characterized by their sizes  $l_i \in \mathbb{N}$ and quantities  $b_i \in \mathbb{N}, i \in I := \{1, \ldots, m\}$ , be given. More compactly, we will refer to these input data by a tuple  $E = (m, l, L, b)$  with  $l := (l_1, \ldots, l_m)^\top \in \mathbb{N}^m$  and  $b := (b_1, \ldots, b_m)^\top \in \mathbb{N}^m$ , termed as an *instance*. In this paper, we consider the following two combinatorial optimization problems:

B John Martinovic john.martinovic@tu-dresden.de

<sup>1</sup> Institute of Numerical Mathematics, Technische Universität Dresden, 01062 Dresden, Germany

- *Cutting stock problem (CSP)*: Find the minimum number of bins (of capacity *L*) that is able to accommodate all items without exceeding the capacity of any bin.
- *Skiving stock problem (SSP)*: Find the maximum number of bins (of capacity *L*) that can be filled using the given items, so that the total load of any bin at least reaches the capacity.

Note that, usually, the CSP and the SSP are introduced from the perspective of cutting large items into smaller ones or combining small objects to obtain larger ones. However, to overcome the difficulty that the input parameters of an instance would partly have different meanings for the CSP and the SSP, we decided to choose the bin packing terminology throughout this article, so that both problems can suitably be addressed by the same vocabulary.

Obviously, the above optimization tasks are united by an economic component (i.e., lower costs in a broad sense) and the idea of sustainability (i.e., restricting the waste of resources). Nevertheless, from a historical point of view, there is a quite remarkable difference. While the CSP already started to attract scientific interest several decades ago, see (Gilmore and Gomor[y](#page-20-0) [1961](#page-20-0); Kantorovic[h](#page-20-1) [1939\)](#page-20-1) for early publications, the SSP is a rather young field of research which was introduced as a natural counterpart of the CSP in a specific application (Johnson et al[.](#page-20-2) [1997;](#page-20-2) Za[k](#page-20-3) [2003](#page-20-3)). In recent years, according to the constantly growing importance of the CSP, see particularly (Delorme et al[.](#page-19-0) [2015](#page-19-0), Figure 1), also a larger body of work has been established with respect to the SSP. For a more detailed introduction to both optimization problems and possible areas of application, here we recommend the survey articles (Delorme et al[.](#page-20-4) [2016](#page-20-4); Valério de Carvalh[o](#page-20-5) [2002](#page-20-5)) for the CSP, the papers (Martinovic et al[.](#page-20-6) [2020](#page-20-6); Martinovic and Scheithaue[r](#page-20-7) [2016a\)](#page-20-7) for the SSP, as well as the book Scheithaue[r](#page-20-8) [\(2018](#page-20-8)) for a general overview on cutting and packing. Observe that although the CSP and the SSP form an obviously related pair of minimization and maximization problems, they are not dual formulations in the sense of mathematical optimization. Consequently, both problems have to be considered as theoretically independent. However, it is worth noting that the CSP and the SSP also appear side-by-side in holistic cutting-andskiving scenarios within different fields of industry (Chen et al[.](#page-19-1) [2019;](#page-19-1) Johnson et al[.](#page-20-2) [1997\)](#page-20-2).

Over time, many different *integer linear programming (ILP)* formulations have been presented for the two problems. Starting with assignment models Kantorovic[h](#page-20-1) [\(1939\)](#page-20-1) or pattern-based approaches (Gilmore and Gomor[y](#page-20-0) [1961](#page-20-0); Za[k](#page-20-3) [2003\)](#page-20-3), nowadays more and more research deals with pseudo-polynomial alternative frameworks like onecut models (Dyckhof[f](#page-20-9) [1981](#page-20-9); Martinovic and Scheithaue[r](#page-20-7) [2016a](#page-20-7)) or flow-based formulations (Delorme and Ior[i](#page-19-2) [2020;](#page-19-2) Martinovic et al[.](#page-20-6) [2020](#page-20-6); Valério de Carvalh[o](#page-20-5) [2002\)](#page-20-5), the latter of which actually showed the most competitive performances so far. Besides structural parameters (like the numbers of variables and constraints or the sparsity of the system matrices), the strength of the corresponding LP relaxation is one of the most important indicators for an efficient solution procedure. More precisely, research has shown, and it became widely accepted, that the quality of the bound provided by the LP relaxation of an ILP model is a crucial factor in the size of the branch-and-bound search trees. In the next section, this aspect shall be introduced more thoroughly.

### <span id="page-3-0"></span>**2 The additive integrality gap: preliminaries and literature review**

Let  $E = (m, l, L, b)$  denote an instance (of the CSP or the SSP). To appropriately quantify the tightness of the LP bound, the *additive integrality gap* (or briefly *gap*), i.e., the difference between the optimal values of the LP relaxation and the original ILP formulation, is considered in this paper. In what follows, we will introduce this concept with respect to the pattern-based formulations presented by Gilmore and Gomor[y](#page-20-0) [\(1961\)](#page-20-0) and Za[k](#page-20-3) [\(2003\)](#page-20-3). Due to well known equivalence results, see (Martinovic et al[.](#page-20-10) [2018,](#page-20-10) Theorem 10) or Delorme and Ior[i](#page-19-2) [\(2020](#page-19-2)) for the CSP and (Martinovic and Scheithaue[r](#page-20-7) [2016a,](#page-20-7) Section 3) for the SSP, the gap is independent of the fact whether the pattern-based model or alternative formulations like flow-based or onecut approaches are considered.

**Definition 1** Let  $E = (m, l, L, b)$  denote an instance (of the CSP or the SSP). Then, any vector  $a \in \mathbb{Z}_+^m$  with

- $-l^{\perp}a \leq L$  is called a *cutting pattern*,
- $-$  *l*<sup>⊥</sup> $a \ge L$  is called a *packing pattern*,
- $l^{\dagger} a = L$  is called an *exact pattern*.

We will refer to the respective sets by  $P_{\leq} := P_{\leq}(E), P_{\geq} := P_{\geq}(E)$ , and  $P_{\equiv} :=$  $P=(E)$ . Moreover, in order to address a specific pattern, we will use the notation  $a^j = (a_1^j, \ldots, a_m^j)^\top \in \mathbb{Z}_+^m$ , where *j* can belong to one of the index sets  $J \leq := J \leq (E)$ ,  $J_{\geq} := J_{\geq}(E)$ , or  $J_{\equiv} := J_{\equiv}(E)$ .

Based on these definitions, we can formulate the

#### **Pattern-based Model of the CSP**

$$
z^{CSP} = \sum_{j \in J_{\leq}} x_j \to \min
$$
  
s.t. 
$$
\sum_{j \in J_{\leq}} x_j \cdot a_i^j \geq b_i, \qquad i \in I,
$$

$$
x_j \in \mathbb{Z}_+, \qquad j \in J_{\leq},
$$

and the

#### **Pattern-based Model of the SSP**

$$
z^{SSP} = \sum_{j \in J_{\ge}} x_j \to \max
$$
  
s.t. 
$$
\sum_{j \in J_{\ge}} x_j \cdot a_i^j \le b_i, \qquad i \in I,
$$

$$
x_j \in \mathbb{Z}_+, \qquad j \in J_{\ge},
$$

where  $x_i$  counts how many bins are filled according to the (cutting or packing) pattern  $a^j$ , while the respective sets of constraints manage that at least (or at most)  $b_i$  items of type  $i \in I$  are used. In both cases, the LP relaxation can be obtained by replacing the condition  $x_i \in \mathbb{Z}_+$  with  $x_i \geq 0$ .

**Definition 2** Let  $E = (m, l, L, b)$  denote an instance (of the CSP or the SSP). Then, the differences

$$
\Delta^{CSP}(E) := z^{CSP,\star}(E) - z_{LP}^{CSP,\star}(E),
$$
  

$$
\Delta^{SSP}(E) := z_{LP}^{SSP,\star}(E) - z^{SSP,\star}(E),
$$

are called the *(additive integrality) gap* of *E*, depending on whether the CSP or SSP is referred to. Here, the index "*L P*" stands for the continuous relaxation, whereas the superscript " $\star$ " indicates the optimal value of the respective formulation.

Besides only considering one single instance, frequently the gap of a whole class *T* of instances is of interest, too. In these situations, the gap of  $T$  shall be understood in the following way:

$$
\Delta^{CSP}(T) := \sup_{E \in T} \Delta^{CSP}(E) \quad \text{or} \quad \Delta^{SSP}(T) := \sup_{E \in T} \Delta^{SSP}(E).
$$

*Remark 1* It is worth mentioning that there is no "dominance" between both gaps. By way of example, consider the instances

 $E = (3, (21, 14, 6), 42, (1, 2, 6))$  with  $\Delta^{CSP}(E) = 41/42 < 1$  and  $\Delta^{SSP}(E) =$  $43/42 > 1$ , or  $-E = (9, (62, 40, 37, 32, 29, 16, 9, 6, 4), 81, (1, 1, 1, 1, 1, 1, 1, 2, 1))$  with  $\Delta^{CSP}(E) = 1$  and  $\Delta^{SSP}(E) = 20/21 < 1$ , see (Kartak et al[.](#page-20-11) [2015,](#page-20-11) Table 2).

For the purpose of a clearer presentation, let us briefly mention the patterns used in the optimal solutions corresponding to the first instance from the above list. For both, the CSP and the SSP, the optimal LP value results to 85/42, and it is obtained by taking the exact patterns

$$
\frac{1}{2} \times (2, 0, 0)^{\top}, \quad \frac{2}{3} \times (0, 3, 0)^{\top}, \quad \frac{6}{7} \times (0, 0, 7)^{\top}.
$$

As for the integer optimization problems, in the CSP scenario three bins are required to pack all the items (e.g., by using  $(1, 1, 1)^\top$ ,  $(0, 1, 4)^\top$ , and  $(0, 0, 1)^\top$  precisely once), while in the SSP case it is not possible to pack more than one bin. Hence, there is no obvious relationship between these two concepts, even though the corresponding optimization problems possess a similar structure.

As regards the CSP, a significant body of work on the additive integrality gap has been employed in recent decades. At the beginning of these investigations, it was conjectured that the gap of the CSP can be bounded from above by the constant 1. A first counterexample to this claim was presented by Marcott[e](#page-20-12) [\(1986\)](#page-20-12), but it required very large (thus practically irrelevant) input data. Some years later, Fieldhous[e](#page-20-13) [\(1990\)](#page-20-13) and Nic[a](#page-20-14) [\(1994a](#page-20-14)) constructed further counterexamples of moderate sizes, so that finally

 $\Delta^{CSP}(E)$  < 2, the so called *modified integer round-up property (MIRUP)*, was conjectured for all instances *E*, see Scheithauer and Tern[o](#page-20-15) [\(1995\)](#page-20-15). With respect to the SSP, a similar claim (the *modified integer round-down property (MIRDP)* Δ*SSP*(*E*) < 2) was introduced by Za[k](#page-20-3) [\(2003\)](#page-20-3). Even nowadays, it is still an open question whether these inequalities hold or not. Moreover, there are only very few discrete optimization problems, whose additive integrality gap is known to be bounded by a small constant (being independent of the instance); see Nemhauser and Par[k](#page-20-16) [\(1991](#page-20-16)) for an example in edge coloring. Due to these reasons, particularly in the last years, research on the gap (of both, the CSP and the SSP) was further intensified, mainly with respect to

- The construction of instances having large gaps (Caprara et al[.](#page-19-3) [2015](#page-19-3); Kartak and Ripatt[i](#page-20-17) [2019](#page-20-17); Rietz and Demp[e](#page-20-18) [2008](#page-20-18); Rietz and Scheithaue[r](#page-20-19) [2002](#page-20-19)),
- Upper bounds for the maximum gap (Martinovic and Scheithaue[r](#page-20-20) [2016b;](#page-20-20) Riet[z](#page-20-21) [2003b;](#page-20-21) Rietz et al[.](#page-20-22) [2002](#page-20-22)),
- The investigation of special cases (Martinovic and Scheithaue[r](#page-20-23) [2017,](#page-20-23) [2018a](#page-20-24)).

One of the most important special cases for which the MIRUP and the MIRDP could successfully be proved is given by the *divisible case*[1](#page-5-0).

**Definition 3** Let  $E = (m, l, L, b)$  denote an instance (of the CSP or the SSP). Then, *E* belongs to the divisible case ( $E \in \mathcal{DC}$  for short) if  $L/l_i \in \mathbb{N}$  holds for all  $i \in I$ .

Note that (in a slight abuse of our initial assumption  $l \in \mathbb{N}^m$ ) for the sake of simplicity, we will always represent an instance  $E \in \mathcal{DC}$  by its normalized form with  $L = 1$  and  $l_i = 1/q_i \in \{1/q : q \in \mathbb{N}\}\$  for all  $i \in I$ . Besides this convenient structure, one main advantage of the divisible case is given by the following result about the optimal LP value:

<span id="page-5-1"></span>**Lemma 1** *Let*  $E = (m, l, 1, b) \in \mathcal{DC}$ *. Then, we have* 

$$
z_{LP}^{CSP,\star}(E) = z_{LP}^{SSP,\star}(E) = l^{\top}b.
$$

*Proof* Let  $l_1 = 1/q_1$  for some  $q_1 \in \mathbb{N}$ . Then, the exact pattern  $a^1 := (q_1, 0, \ldots, 0)^\top \in$  $P=(E)$  can be used  $x_1 = b_1/q_1$  times in the LP relaxation. In the same manner, we can proceed with the further items and obtain the objective value  $\sum_{i \in I} b_i / q_i = l^{\dagger} b$ . This value is optimal, for the CSP as well as for the SSP, since we did not waste any material.

Hence, only one optimization problem needs to be solved when investigating the gap of the divisible case.

From a theoretical point of view, the following key results have already been obtained in the literature:

- W[e](#page-20-25) have  $\Delta^{CSP}(E)$  < 2, see e.g. (Marcotte [1983](#page-20-25), Theorem 2.16), and  $\Delta^{SSP}(E)$  < 2, see e.g. (Ma[r](#page-20-20)tinovic and Scheithauer [2016b](#page-20-20), Theorem 9), for all  $E \in \mathcal{DC}$ .
- Meanwhile, these upper bounds could be improved to  $\Delta^{CSP}(E) < 1.4$ , see (Riet[z](#page-20-21) [2003b,](#page-20-21) Satz 1), and  $\Delta^{SSP}(E) < 3/2$ , see Ma[r](#page-20-23)tinovic and Scheithauer [\(2017](#page-20-23)).

<span id="page-5-0"></span><sup>&</sup>lt;sup>1</sup> An exemplary instance is given in the first part of the previous remark.

Observe that the first result has never been published in English language, and, unfortunately, its documentation in the German dissertation Riet[z](#page-20-21) [\(2003b\)](#page-20-21) was not established thoroughly enough to understand and reconstruct all the arguments. Hence, we will not make active use of this inequality here.

- The largest known gaps for the divisible case are  $\Delta^{CSP}(E) = 137/132$  provided by *E* = (3, (44, 33, 12), 132, (2, 3, 6)), see (Scheithaue[r](#page-20-8) [2018,](#page-20-8) Section 4.11.6), and  $\Delta^{SSP}(E) = 22/21$  approximated by an infinite sequence of instances, see (Martinovic and Scheithaue[r](#page-20-20) [2016b](#page-20-20), Corollary 16). Actually, it is even conjectured that 22/21 represents the best possible upper bound for the divisible case of the SSP, see (Martinovic and Scheithaue[r](#page-20-26) [2018b,](#page-20-26) Conjecture 1).
- Upper bounds for the gap of the divisible case can directly be used to formulate upper bounds for the gap of arbitrary instances, see (Martinovic and Scheithaue[r](#page-20-27) [2019,](#page-20-27) Theorem 4) for an example. Hence, any improvement that is achieved for this special case can be transferred to more general instances, too.

Note that, besides being of theoretical importance, the divisible case sometimes also appears in practically relevant scenarios, especially when there is a high degree of standardization such as in bin-packing based (multiprocessor) scheduling applications (Bar-Noy et al[.](#page-19-4) [2007;](#page-19-4) Coffman et al[.](#page-19-5) [1987\)](#page-19-5).

Given the aforementioned observations, the main contributions of the present article can be summarized as follows:

- We introduce two new reduction strategies that allow to only focus on a considerably restricted set of (somewhat homogeneous) instances ( $\rightarrow$  Sect. [3\)](#page-6-0).
- We derive the improved upper bound  $\Delta^{CSP}(E) < 4/3$  for all  $E \in \mathcal{DC} \rightarrow$  Sect. [4\)](#page-10-0).
- We transfer the previous result to the SSP, i.e., we show  $\Delta^{SSP}(E) < 4/3$  for all  $E \in \mathcal{DC} \; (\rightarrow \text{Sect. 5}).$

On the one hand, we significantly improve the currently best upper bounds for the gap of the divisible case for both, the CSP and the SSP. Moreover, we highlight the fact that our underlying theoretical approach is very general, so that a much more extensive analysis of the applied arguments could potentially lead to further improvements of these upper bounds.

#### <span id="page-6-0"></span>**3 Reduction strategies**

The following definition will be required to decompose the set of all instances of the divisible case.

**Definition 4** Let  $\delta \in \mathbb{N}$  with  $\delta \geq 2$  be given. For any  $n \in \mathbb{N}$ , we define the sets

$$
\mathcal{DC}^{-}(n,\delta) := \left\{ E = (m,l,1,b) \in \mathcal{DC} : l^{\top}b \in \left[n-1, n-\frac{1}{\delta}\right] \right\},\
$$

$$
\mathcal{DC}^{+}(n,\delta) := \left\{ E = (m,l,1,b) \in \mathcal{DC} : l^{\top}b \in \left[n+\frac{1}{\delta}, n+1\right] \right\}.
$$

*Remark 2* In fact, the integrality of the parameter  $\delta$  is not required for the definition here, and also most of the general arguments used during this article could be modified to become applicable to  $\delta \in \mathbb{R}$ , too. However, since integer values are sufficient for our main contributions and slightly simplify the corresponding formulas and calculations, we will restrict ourselves to the integrality assumption, but not without pointing out at the corresponding places, which changes would be necessary in case of real numbers or which challenges (e.g., in the proofs) would result from that.

For fixed  $\delta \geq 2$ , the sets  $\mathcal{DC}^-(n, \delta)$ ,  $n \in \mathbb{N}$ , will be used for the CSP, whereas the sets  $DC^{+}(n, \delta)$ ,  $n \in \mathbb{N}$ , will later be important for the SSP case. Observe that, due to Lemma [1,](#page-5-1) the optimal LP value of an instance  $E \in \mathcal{DC}^-(n, \delta)$  is known to satisfy

$$
z_{LP}^{CSP,\star}(E) \in \left[ n-1, n-\frac{1}{\delta} \right],
$$

while  $E \in \mathcal{DC}^+(n, \delta)$  directly implies

$$
z_{LP}^{SSP,\star}(E) \in \left[ n + \frac{1}{\delta}, n + 1 \right].
$$

In the following, we first concentrate on the CSP case. To this end, let  $\delta \geq 2$  be fixed to some appropriate value (which is specified later).

**Lemma 2** *If for any n* ∈ *N and any instance*  $\widetilde{E}$  ∈  $\mathcal{DC}^-(n, \delta)$ *, the items of*  $\widetilde{E}$  *can be assigned to (at most) n bins (cutting patterns), then we have*

<span id="page-7-0"></span>
$$
\Delta^{CSP}(E) < 1 + \frac{1}{\delta}
$$

*for all*  $E \in \mathcal{DC}$ *.* 

*Proof* For any instance  $E = (m, l, 1, b) \in \mathcal{DC}$  there is some  $n := n(E) \in \mathbb{N}$  with  $l^{\perp} b \in [n-1, n)$ :

 $-$  If *E* ∈ *DC*<sup>−</sup>(*n*, δ), then we have *z*<sup>CSP,★</sup>(*E*) < *n* by hypothesis and, finally,

$$
\Delta^{CSP}(E) = z^{CSP,\star}(E) - z_{LP}^{CSP,\star}(E) \le n - l^{\top}b \le 1 < 1 + \frac{1}{\delta}.
$$

 $-$  If *E* ∉ *DC*<sup>−</sup>(*n*, δ), then we first have

$$
z_{LP}^{CSP,\star}(E) = l^{\top}b \in \left(n - \frac{1}{\delta}, n\right).
$$

On the other hand, given that MIRUP holds for the entire divisible case, we obtain  $z^{CSP,\star}(E) \in \{n, n+1\}$ . Both observations lead to

$$
\Delta^{CSP}(E) = z^{CSP,\star}(E) - z_{LP}^{CSP,\star}(E) < (n+1) - \left(n - \frac{1}{\delta}\right) = 1 + \frac{1}{\delta}.
$$

 $\mathcal{D}$  Springer

**Remark 3** In an analogous manner, proving that the items of any instance  $\tilde{E} \in \mathbb{R}^{C+}(u, s)$ ,  $u \in \mathbb{N}$  can be used to huild (et locat) using (positive national) is sufficient  $\mathcal{DC}^+(n, \delta), n \in \mathbb{N}$ , can be used to build (at least) *n* bins (packing patterns) is sufficient to show

$$
\Delta^{SSP}(E) < 1 + \frac{1}{\delta}
$$

for all  $E \in \mathcal{DC}$ . Indeed, let us consider an instance  $E \in \mathcal{DC}$  (of the SSP) with  $l^{\top}b \in (n, n + 1]$  for some  $n \in \mathbb{N}$ , then one of the two cases appears:

 $-$  For *E* ∈ *DC*<sup>+</sup>(*n*, δ), we have *z*<sup>SSP,★</sup>(*E*) > *n* by hypothesis, leading to

$$
\Delta^{SSP}(E) = z_{LP}^{SSP,\star}(E) - z^{SSP,\star}(E) \le l^{\top}b - n \le (n+1) - n = 1 < 1 + \frac{1}{\delta}.
$$

 $-$  For *E* ∉ *DC*<sup>+</sup>(*n*, δ), we have *z*<sup>*SSP*,\*(*E*) ∈ {*n* − 1, *n*} by MIRDP (which holds for</sup> the entire divisible case), leading to

$$
\Delta^{SSP}(E) = z_{LP}^{SSP,\star}(E) - z^{SSP,\star}(E) \le l^{\top}b - (n-1) < \left(n + \frac{1}{\delta}\right) - (n-1) = 1 + \frac{1}{\delta}.
$$

We refer the reader to Sect. [5](#page-16-0) for more details concerning the SSP case.

Obviously, for fixed  $\delta$ , the sets  $\mathcal{DC}^-(n, \delta)$ ,  $n \in \mathbb{N}$ , contain an infinite number of instances to be checked with respect to Lemma [2.](#page-7-0) Moreover, these instances can be considered to be very heterogeneous since neither the sizes  $l_i$  nor the quantities  $b_i$ , *i* ∈ *I*, of an instance  $E \text{ } \in \mathcal{DC}^-(n, \delta)$  are restricted. To overcome these issues, we will introduce two reduction strategies leading to

- (1) an upper bound for the quantities  $b_i$ ,  $i \in I$ ,
- (2) a lower bound for the sizes  $l_i$ ,  $i \in I$ ,

so that coping with only finitely many cases (each of which offering some structural information) will be sufficient to prove the result for all possible instances of the divisible case.

Before explaining these steps in more details, the following definition is required:

**Definition 5** An instance  $E = (m, l, 1, b) \in \mathcal{DC}$  is called *irreducible*, if two or more items (not necessarily having different sizes) cannot be used to build a unit fraction. More formally, an irreducible instance is characterized by

$$
a \le b, \sum_{i \in I} a_i \ge 2 \Longrightarrow \forall k \in \mathbb{N} : \sum_{i \in I} a_i l_i \ne \frac{1}{k}
$$

for any vector  $a \in \mathbb{Z}_{+}^{m}$ . Otherwise, *E* is called *reducible*.

<span id="page-9-1"></span>*Remark 4* In particular, any irreducible instance  $E \in \mathcal{DC}$  has the following properties:

- − *E* does not possess any exact pattern  $a \text{ ∈ } P_=(E)$  with  $a \text{ ≤ } b$  (in a componentwise sense), since this would contradict to the above definition for  $k = 1$ .
- Let *<sup>t</sup>*(*q*) denote the smallest prime divisor of *<sup>q</sup>* <sup>∈</sup> <sup>N</sup> with *<sup>q</sup>* <sup>≥</sup> 2. Then, any item of size  $1/q$  can appear at most  $t(q) - 1$  times in *E*. (Otherwise,  $t(q)$  items of size 1/*q* would sum up to a unit fraction).
- In addition to the previous observation, item combinations like  $1/3 + 1/6 = 1/2$ or  $1/4 + 1/12 = 1/3$  cannot appear in *E*.

In order to implement the first reduction mentioned above, we now proceed as follows:

**Reduction 1** *If*  $z^{CSP, \star}(E) \leq n$  *can be shown for all irreducible instances*  $E \in$  $DC^{-}(n, \delta)$ *, then it also holds for any reducible instance in*  $DC^{-}(n, \delta)$ *.* 

*Proof* Consider a reducible instance  $E \in \mathcal{DC}^-(n, \delta)$ . Then, we can find a subset (containing at least two items) summing up to some value  $1/k$ ,  $k \in \mathbb{N}$ . By replacing this subset of items with one artificial item of size  $1/k$ , we obtain an instance  $E'$  with fewer items but the same optimal LP value. After a finite number of such steps, we end up with an irreducible instance whose items can be packed into at most *n* bins (cutting patterns) by hypothesis. In this feasible packing, any artificial item of size 1/*k* can be replaced by the corresponding subset of original items (of *E*) which was used to build the item of size  $1/k$ . Consequently, also the items of E can be packed into at most *n* bins and we are done.

While this method can mainly be used to restrict the quantities  $b_i$ ,  $i \in I$ , of an instance, the second reduction strategy is useful to only focus on the "large items" of *E*.

**Definition 6** Let  $\delta \geq 2$ ,  $n \in \mathbb{N}$ , and an instance  $E \in \mathcal{DC}$  be given. Then, the set of *large items* (of *E*) is defined by

<span id="page-9-0"></span>
$$
\Lambda_E := \Lambda_E(n,\delta) = \left\{ i \in I : l_i \ge \frac{1}{\delta \cdot (n-1) - 1} \right\}.
$$
 (1)

All other items will be termed as *small*.

*Remark 5* When dealing with real-valued parameters  $\delta$ , the definition has to be adjusted, so that an item  $i \in I$  is called large, whenever it satisfies

$$
l_i \geq \frac{1}{\lceil \delta \cdot (n-1) - 1 \rceil}.
$$

Then, we obtain the following observation.

<span id="page-10-2"></span>**Reduction 2** *Let*  $E$  ∈  $DC^{-}(n, \delta)$  *be irreducible. If it is possible to assign all large items of E to at most n bins (cutting patterns), then we have*  $z^{CSP,\star}(E) \le n$ , *i.e., all items actually fit into those n bins.*[2](#page-10-1)

*Proof* Let us assume that, after having assigned the large items of *E* to at most *n* bins, it is not possible to put one of the remaining small items, say one item of size  $l^* := l_i$ for some  $i \in I \setminus \Lambda_E$ , into the existing bins without exceeding the bin size. This can only happen, if for any bin  $B_j$ ,  $j \in \{1, ..., n\}$ , the total load  $C(j)$  (of the items already allocated to  $B_j$ ) is greater than  $1 - l^*$ . Consequently, we have

$$
n - \frac{1}{\delta} \ge l^{\top} b \ge \sum_{j=1}^{n} C(j) + l^{\star} > n \cdot (1 - l^{\star}) + l^{\star}.
$$

Rearranging the terms leads to

$$
n \cdot (1 - l^*) + l^* < n - \frac{1}{\delta} \Longleftrightarrow (n - 1)l^* \\
> \frac{1}{\delta} \Longleftrightarrow l^* > \frac{1}{\delta \cdot (n - 1)} \Longleftrightarrow l^* \ge \frac{1}{\delta \cdot (n - 1) - 1},
$$

where the last equivalence is true since any item size is a unit fraction. However, this would imply that the item of size  $l^*$  is a large item which already has been feasibly assigned to a bin by hypothesis. Hence, we obtain a contradiction and the statement is proved.  $\Box$ 

This second reduction strategy allows to only consider the large items of an instance  $E \in \mathcal{DC}^{-}(n, \delta)$ , where the term *large* is specified by Condition [\(1\)](#page-9-0). Moreover, it implicitly states that, after having distributed these objects, the small items can be assigned in an arbitrary manner, as long as the capacities are respected. By way of example, one appropriate strategy to assign the small items is based on the *best-fit decreasing heuristic*, as described in (Martinovic and Scheithaue[r](#page-20-23) [2017,](#page-20-23) Algorithm 1). In that algorithm, the (remaining) items are sorted with respect to non-increasing sizes and processed separately. More precisely, in a given iteration, the currently largest object is taken and it is assigned to the least filled bin (where ties are broken by choosing the lowest-indexed bin).

#### <span id="page-10-0"></span>**4 The CSP case: improved upper bounds for the gap**

As a first contribution, we consider the case  $\delta = 2$  which would lead to the upper bound  $\Delta^{CSP}(E)$  < 3/2 for  $E \in \mathcal{DC}$  according to Lemma [2.](#page-7-0) The reasons for presenting this auxiliary result are twofold:

<span id="page-10-1"></span><sup>2</sup> Actually,  $z^{CSP,\star}(E) \le n$  for an instance  $E \in \mathcal{DC}^-(n, \delta)$  already shows the optimality of the value *n*. However, as the feasibility of the assignment is sufficient for our purposes, we do not specifically stress this fact.

- Remember that the proof for the currently best upper bound  $\Delta^{CSP}(E)$  < 1.4 has never been published in English language, and so it is hardly known in the scientific community. Hence, instead of using a result that cannot easily be verified, it is more convenient to briefly prove an upper bound of nearly the same quality.
- Later, we can apply this preliminary observation to appropriately deal with some subcases appearing in our main theorem.

<span id="page-11-1"></span>**Theorem 1** *Let*  $n \in \mathbb{N}$  *be given, and let us consider an irreducible instance*  $E \in$ *DC*−(*n*, 2)*. Then, these items can be assigned to at most n bins (cutting patterns). In particular, we have*  $\Delta^{CSP}(E) < 3/2$  *for all*  $E \in \mathcal{DC}$ *.* 

*Proof* It is sufficient to consider the large items of E, meaning that we focus on those items with

$$
l_i \ge \frac{1}{\delta \cdot (n-1) - 1} = \frac{1}{2n - 3}.
$$

Since *E* is irreducible, for any fixed  $k = 1, \ldots, n - 1$  we have at most one item of size  $1/(2k)$  and at most  $2k - 2$  items of size  $1/(2k - 1)$ . Due to

$$
\frac{1}{2k} + \frac{2k - 2}{2k - 1} = 1 + \frac{1}{2k} - \frac{1}{2k - 1} = 1 - \frac{1}{2k(2k - 1)} < 1,
$$

these items would fit into one bin. Hence, the large items of *E* can be assigned to at most *n* bins<sup>[3](#page-11-0)</sup> which concludes the proof thanks to Reduction [2.](#page-10-2)

Note that also this proof is constructive since it contains a precise method to obtain a feasible solution using *n* bins. More precisely, the large items of *E* are grouped according to the instructions in the previous proof, whereas the small items can exemplarily be distributed by the best-fit decreasing heuristic from (Martinovic and Scheithaue[r](#page-20-23) [2017,](#page-20-23) Algorithm 1).

Now we intend to show an analogous result for  $\delta = 3$  which would lead to the improved upper bound  $\Delta^{CSP}(E) < 4/3$  for all  $E \in \mathcal{DC}$ . To this end, let us consider an irreducible instance  $E = (m, l, 1, b) \in \mathcal{DC}^{-}(n, 3)$  for some  $n \in \mathbb{N}$ . Note that for  $n = 1$  it is obvious that all items can be packed into one bin. Consequently, we can assume  $n \geq 2$ . Moreover, due to Reduction [2,](#page-10-2) it is sufficient to consider the large items of *E*, meaning that we only have to show that all items with

$$
l_i \ge \frac{1}{\delta \cdot (n-1) - 1} = \frac{1}{3n - 4},
$$

can be assigned to (at most) *n* bins (cutting patterns). To prove the latter, Theorem [1](#page-11-1) can be applied. More formally, for  $\delta = 3$ , let  $T_E(n) := T_E(n, \delta)$  denote the total size of all large items appearing in *E*, i.e.,

$$
T_E(n) := \sum_{i \in \Lambda_E(n,\delta)} l_i b_i = \sum_{i \in \Lambda_E(n,3)} l_i b_i,
$$

<span id="page-11-0"></span><sup>&</sup>lt;sup>3</sup> Actually, we have shown a bit more, namely that already  $n - 1$  bins are sufficient.

and let us define

$$
T(n) := \max \{ T_E(n) : E \in \mathcal{DC}^-(n, 3) \text{ is irreducible} \}.
$$

Then it suffices<sup>4</sup> to verify that

<span id="page-12-1"></span>
$$
T(n) \le n - \frac{1}{2} \tag{2}
$$

holds for all  $n \ge 2$ . To prove this claim, observe that for  $N := 3n - 4$  we have

$$
T(n) \le \underbrace{\sum_{q=2}^{N} \frac{t(q) - 1}{q}}_{=: U(N)} - R(N),
$$

where  $U(N) = U(3n - 4)$  sums up at most  $t(q) - 1$  items of size  $1/q$  for any large item  $q = 2, \ldots, N = 3n - 4$  (again with  $t(q)$  representing the smallest prime divisor of *q*), and  $R(N) > 0$  is a correction term that possibly subtracts some item sizes that we obtain from forbidden item combinations, see Remark [4.](#page-9-1) Hence, our main strategy in the following proofs consists of showing

<span id="page-12-2"></span>
$$
U(3n - 4) - R(3n - 4) \le n - \frac{1}{2}
$$
 (3)

<span id="page-12-3"></span>for an appropriately defined term  $R(3n - 4) \ge 0$ , which directly implies Condition [\(2\)](#page-12-1).

**Theorem 2** *Condition [\(2\)](#page-12-1) is true for all*  $n \in \{2, 3, ..., 24\}$ *.* 

*Proof* At first note that the values  $U(3n - 4)$  given in [\(3\)](#page-12-2) can be easily computed, see Table [1.](#page-13-0)

Based on these data and the choice  $R(N) = R(3n - 4) = 0$ , we can directly see that [\(3\)](#page-12-2) is true for  $n \in \{2, 3, 4\}$  ∪  $\{10, 11, ..., 24\}$ . The remaining cases can be dealt with separately:

– For *n* = 5 we have 3*n* − 4 = 11 and *U*(11) ≈ 4.5968 > 5 − 1/2. However, since  $1/6+1/3 = 1/2$  holds, we can at least subtract  $R(11) = 1/6$  from  $U(11)$ , so that we end up with

$$
T(5) \le U(11) - \frac{1}{6} \le 5 - \frac{1}{2}.
$$

<span id="page-12-0"></span><sup>&</sup>lt;sup>4</sup> Indeed, then the large items of *E* either form an instance of  $\mathcal{DC}(n, 2)$  (if  $T_E(n) \ge n - 1$  holds), so that they can be assigned to at most *n* bins by the previous theorem. Or we have  $T_E(n) < n - 1$  also meaning that at most *n* bins are sufficient to accommodate all large items as a direct consequence of MIRUP for the divisible case.

<span id="page-13-0"></span>

The last column of the table indicates whether Condition [\(3\)](#page-12-2) is satisfied  $(\checkmark)$  or not  $(\times)$ 

– For *n* = 6 we have 3*n* − 4 = 14 and *U*(14) ≈ 5.6746 > 6 − 1/2. However, since additionally  $2/5+1/10 = 1/2$  holds, we can at least subtract  $R(14) = 1/6+1/10$ from  $U(14)$ , so that we end up with

$$
T(6) \le U(14) - \frac{1}{6} - \frac{1}{10} \le 6 - \frac{1}{2}.
$$

– For *n* = 7 we have 3*n* − 4 = 17 and *U*(17) ≈ 6.8116 > 7 − 1/2. However, since additionally  $1/4 + 1/12 = 1/3$  holds, we obtain

$$
T(7) \le U(17) - \frac{1}{6} - \frac{1}{10} - \frac{1}{12} \le 7 - \frac{1}{2}.
$$

– For *n* = 8 we have 3*n* − 4 = 20 and *U*(20) ≈ 7.8646 > 8 − 1/2. However, since additionally  $3/7 + 1/14 = 1/2$  holds, we obtain

$$
T(8) \le U(20) - \frac{1}{6} - \frac{1}{10} - \frac{1}{12} - \frac{1}{14} \le 8 - \frac{1}{2}.
$$

<sup>2</sup> Springer

<span id="page-14-1"></span> $\Box$ 

– For *n* = 9 we have 3*n* − 4 = 23 and *U*(23) ≈ 8.9618 > 9 − 1/2. However, since additionally  $1/9 + 1/18 = 1/6$  holds, we obtain

$$
T(9) \le U(23) - \frac{1}{6} - \frac{1}{10} - \frac{1}{12} - \frac{1}{14} - \frac{1}{18} \le 9 - \frac{1}{2}.
$$

Obviously, we cannot compute the true values  $T(n)$  for all  $n \in \mathbb{N}$ , and neither the upper bound  $U(3n - 4)$  can always be used since, depending on *n*, we would possibly have to find more and more forbidden item combinations (which can contribute to  $R(3n - 4)$ ). The next lemma helps us to establish an upper bound for  $T(n)$  based on some well known properties of prime numbers.

**Lemma 3** *For*  $N \ge 8$  *(with*  $N = 3n - 4$  *for some*  $n \in \mathbb{N}$ *) we have* 

$$
T(n) < \frac{N}{\ln(N)} \cdot \left( 1 + \frac{1}{\ln(N)} + \frac{2}{(\ln(N))^2} + \frac{7.59}{(\ln(N))^3} \right) + \frac{1}{30}
$$

**Proof** Let  $N > 8$  be fixed. At first we define the set

$$
U_p(N) := \{k \in \{2, \ldots, N\} : t(k) = p\}
$$

for every prime  $p \in \mathbb{P}$  with  $2 \leq p \leq N$ . Note that  $U_p(N)$  contains all integers from  $\{2, \ldots, N\}$  having p as the smallest prime divisor. The main calculation of this proof requires some inequalities which we are going to mention beforehand:

(A) Let  $N := 3n - 4 \ge 8$  (meaning that we have  $n \ge 4$ ) and  $R(N) = 1/6$ , then we obtain

$$
T(n) \le U(N) - \frac{1}{6} = \sum_{q=2}^{N} \frac{t(q) - 1}{q} - \frac{1}{6},
$$

since only one of the items 1/3 and 1/6 can be part of an irreducible instance. (B) Since every  $q \in U_p(N)$  has p as the smallest prime divisor, we can state

$$
\sum_{q \in U_p(N)} \frac{1}{\frac{q}{p}} < \sum_{k=0}^{\infty} \frac{1}{p^k} = \frac{1}{1 - \frac{1}{p}} = 1 + \frac{1}{p-1}
$$

due to the well known convergence properties of the geometric series. (C) Due to  $N \geq 8$ , we have

<span id="page-14-0"></span>
$$
\sum_{p \in \mathbb{P}, p \le N} \frac{1}{p-1} - \sum_{q=2}^{N} \frac{1}{q} \le 1 - \frac{1}{3} - \frac{1}{5} - \frac{1}{7} - \frac{1}{8} < \frac{1}{5}.\tag{4}
$$

For  $N = 8$  this can be verified by a direct calculation. Beyond that, we can see the correctness of this inequality in an inductive way. More precisely, the difference on the left hand side definitely decreases until we reach the next prime number  $N = p$  because the first sum stays constant while the second sum gathers more and more unit fractions. Hence, for the induction, we only need to focus on those steps referring to prime numbers. Having arrived at  $N = p \in \mathbb{P}$ , the first sum receives an additional term  $1/(N-1)$ . However, since (for  $N > 8$ ) we do not find two consecutive prime numbers, the second sum at least increased by  $1/(N-1) + 1/N$  compared to the previous prime number (where we noticed the last increase of the first sum), meaning that the overall difference stays below 1/5.

(D) Let  $\pi(N)$  count the number of primes in  $\{2, \ldots, N\}$ , then we can state

$$
\pi(N) \le \frac{N}{\ln(N)} \cdot \left(1 + \frac{1}{\ln(N)} + \frac{2}{(\ln(N))^2} + \frac{7.59}{(\ln(N))^3}\right)
$$

by means of (Dusar[t](#page-20-28) [2018,](#page-20-28) Theorem 5.1). This inequality even holds for  $N \ge 2$ . Based on these ingredients, we can proceed as follows

$$
T(n) \stackrel{\text{(A)}}{\leq} U(N) - \frac{1}{6} = \sum_{q=2}^{N} \frac{t(q)}{q} - \sum_{q=2}^{N} \frac{1}{q} - \frac{1}{6} = \sum_{p \in \mathbb{P}, p \leq N} \sum_{q \in U_p(N)} \frac{1}{\frac{q}{p}} - \sum_{q=2}^{N} \frac{1}{q} - \frac{1}{6}
$$
  
\n
$$
\stackrel{\text{(B)}}{\leq} \sum_{p \in \mathbb{P}, p \leq N} \left(1 + \frac{1}{p-1}\right) - \sum_{q=2}^{N} \frac{1}{q} - \frac{1}{6}
$$
  
\n
$$
= \sum_{p \in \mathbb{P}, p \leq N} 1 + \sum_{p \in \mathbb{P}, p \leq N} \frac{1}{p-1} - \sum_{q=2}^{N} \frac{1}{q} - \frac{1}{6}
$$
  
\n
$$
\stackrel{\text{(C)}}{\leq} \pi(N) + \frac{1}{5} - \frac{1}{6} \stackrel{\text{(D)}}{\leq} \frac{N}{\ln(N)} \cdot \left(1 + \frac{1}{\ln(N)} + \frac{2}{(\ln(N))^2} + \frac{7.59}{(\ln(N))^3}\right) + \frac{1}{30},
$$

and the proof is complete.

Note that  $N \geq 8$  is required in the proof to obtain the additional term  $-1/8$  when subtracting the two sums in [\(4\)](#page-14-0).

For the sake of simplicity, let us use

<span id="page-15-0"></span>
$$
\widetilde{U}(N) := \frac{N}{\ln(N)} \cdot \left(1 + \frac{1}{\ln(N)} + \frac{2}{(\ln(N))^2} + \frac{7.59}{(\ln(N))^3}\right) + \frac{1}{30}
$$

as an abbreviation. Then it can be verified that:

**Theorem 3** *Condition* [\(2\)](#page-12-1) *is true for all n*  $\geq$  25*.* 

*Proof* Let  $n \ge 25$  and  $N := 3n - 4$  be fixed, then we have  $T(n) < \widetilde{U}(N) \le n - \frac{1}{2}$ i.e., Condition [\(2\)](#page-12-1) is true. For all the details, we refer the interested reader to the "Appendix". Putting Theorems [2](#page-12-3) and [3](#page-15-0) together, we can conclude that:

**Theorem 4** *We have*  $\Delta^{CSP}(E) < 4/3$  *for all*  $E \in \mathcal{DC}$ *.* 

Consequently, we have found a new and improved upper bound for the gap of divisible case instances of the CSP.

## <span id="page-16-0"></span>**5 A transfer to the SSP case**

Let us now consider an instance  $E = (m, l, 1, b) \in \mathcal{DC}$  of the skiving stock problem. In analogy to the CSP case we can focus on irreducible instances.

**Lemma 4** *Let*  $E \in \mathcal{DC}^+(n, 3)$  *be irreducible, then the items of*  $E$  *can be assigned to n bins (packing patterns).*

*Proof* Let us consider the large items of *E*, i.e., those items satisfying

$$
l_i \geq \frac{1}{3n-4}.
$$

In the proofs of the previous section, we have shown that (for given *n* and  $\delta = 3$ ) all large items (of any irreducible instance  $E \in \mathcal{DC}$ ) can always be assigned to *n* cutting<br>natterns. Hence, the same is two for the large items contained in the considered instance patterns. Hence, the same is true for the large items contained in the considered instance *E*, so that they possess a total size of at most *n*. Thus, some items of the SSP instance are still to be distributed. Note that all of them are small items, i.e., they have a size

$$
l_i \leq \frac{1}{3n-3}.
$$

Let us assign the remaining items based on the best-fit decreasing heuristic presented in (Martinovic and Scheithaue[r](#page-20-23) [2017,](#page-20-23) Algorithm 1), meaning that the current item is allocated to a bin  $B_j$ ,  $j \in \{1, \ldots, n\}$ , whose total item load  $C(j)$  is the smallest. Then, it is clear that a bin  $B_i$  with  $C(j) \geq 1$  will receive an additional item only in those cases when all bins actually satisfy  $C(j) \geq 1$ .

For the sake of contradiction, let us assume that after having assigned the last item of *E*, there is some  $k \in \{1, \ldots, n\}$  with  $C(k) < 1$ , i.e., we did not end up with a feasible solution for the SSP. This would mean that no item was assigned to a bin which already represented a packing pattern, i.e., any filled bin satisfies

$$
1 \le C(j) < 1 + \frac{1}{3n - 3},
$$

since only small items are distributed in this phase. This would lead to

$$
n + \frac{1}{3} \le l^{\top} b = \sum_{j=1}^{n} C(j) = C(k) + \sum_{j \neq k} C(j) < 1 + (n-1) \cdot \left( 1 + \frac{1}{3n - 3} \right)
$$

$$
= n + \frac{n-1}{3n-3} = n + \frac{1}{3}
$$

giving the contradiction. Hence, we have constructed *n* bins (for the SSP) and the proof is complete.

Given this observation, a direct consequence is the following:

# **Theorem 5** *We have*  $\Delta^{SSP}(E) < \frac{4}{3}$  *for any*  $E \in \mathcal{DC}$ *.*

Hence, we have also improved the best upper bound for the SSP case to 4/3. Note that a much more detailed analysis of the arguments applied in the CSP case could also directly influence the quality of the upper bound of the SSP.

*Remark 6* In fact, the strategy from the previous proof is much more general. In the CSP case, we have not only proved that the large items of a given instance  $E \in \mathcal{DC}^-(n, \delta)$ would fit into at most *n* cutting patterns; the proofs of Theorems [2](#page-12-3) and [3](#page-15-0) actually show that all large items of any(!) irreducible instance  $E \in \mathcal{DC}$  can be assigned to at most *n* bins<sup>3</sup>. In the proof of the previous lemma, we used this observation to easily transfer the upper bound from the CSP to the SSP. Effectively, this means that we can apply the same proof also for some larger values of  $\delta$ , as long as we can verify that all large items  $i \in A_E(n, \delta)$  of any irreducible instance  $E \in \mathcal{DC}$  can feasibly be assigned to (at most) *n* bins without exceeding their capacities.

Finally, it is important to note again that improved upper bounds for the divisible case directly influence the upper bounds for more general instances. By way of example, given the results of this section, the constant term in (Martinovic and Scheithaue[r](#page-20-27) [2019,](#page-20-27) Theorem 4) can now be reduced from 3/2 to 4/3 for any arbitrary instance *E* of the SSP.

# **6 Conclusions**

In this article, we investigated the additive integrality gap of both, the CSP and the SSP, from a theoretical point of view. In particular, we focussed on the well known divisible case, where the bin size is assumed to be an integer multiple of any item size  $l_i$ ,  $i \in I$ . For such instances we first developed two reduction strategies to considerably limit the number of instances that has to be considered. More precisely, one of these reductions aims at restricting the quantities  $b_i$ , while the second one implements a lower bound on the item sizes  $l_i$ ,  $i \in I$ . Based on these two observations, we were able to state the improved upper bound 4/3 for the CSP and the SSP. Moreover, our approach potentially offers the possibility of further improvements if a much more extensive analysis is conducted. Another aspect of future research deals with narrowing the interval between the currently best upper bounds and the largest known gaps provided by concrete instances, as mentioned at the end of Sect. [2.](#page-3-0)

<span id="page-17-0"></span><sup>5</sup> Indeed, in Theorems [2](#page-12-3) and [3](#page-15-0) we did not use the fact that the considered instance has a total item size  $l \mid b \in [n-1, n-1/\delta]$  to prove that there is a feasible assignment to at most *n* bins.

**Acknowledgements** The author would like to thank two anonymous referees for their very valuable comments during the review process. In particular, the ideas presented by one of these experts contributed to easier proofs of Lemma [3](#page-14-1) and Theorem [3.](#page-15-0)

**Funding** Open Access funding enabled and organized by Projekt DEAL.

#### **Compliance with ethical standards**

**Conflict of interest** The author declares that he has no conflict of interest.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit [http://creativecommons.org/licenses/by/4.0/.](http://creativecommons.org/licenses/by/4.0/)

## **A A Proof of Theorem 3**

Let  $n \ge 25$  and  $N := 3n - 4$ . Remember that we have to show  $\widetilde{U}(N) \le n - \frac{1}{2}$  or equivalently

<span id="page-18-0"></span>
$$
\frac{N}{\ln(N)} \cdot \left(1 + \frac{1}{\ln(N)} + \frac{2}{(\ln(N))^2} + \frac{7.59}{(\ln(N))^3}\right) + \frac{1}{30} \le n - \frac{1}{2}
$$
(5)

- First of all, note that for *n* = 25 (i.e., *N* = 71) we have *U*(71) ≈ 24.0625, so that [\(5\)](#page-18-0) holds by a direct calculation.
- Now let us interpret both sides of [\(5\)](#page-18-0) as a function of  $n \in \mathbb{R}$ , i.e., we set

$$
f_1(3n-4) := \frac{3n-4}{\ln(3n-4)} \cdot \left(1 + \frac{1}{\ln(3n-4)} + \frac{2}{(\ln(3n-4))^2} + \frac{7.59}{(\ln(3n-4))^3}\right) + \frac{1}{30},
$$
  

$$
f_2(n) := n - \frac{1}{2}.
$$

Given the previous observation  $\tilde{U}(71) \leq 25 - \frac{1}{2}$ , it suffices to show that

$$
\frac{d}{dn}f_1(3n-4) < \frac{d}{dn}f_2(n) \Longleftrightarrow \frac{d}{dn}f_1(3n-4) < 1 \tag{6}
$$

holds for all  $n > 25$ . By a direct calculation, it can be verified that

$$
\frac{d}{dn}f_1(3n-4) = \frac{3 \cdot ((\ln(3n-4))^4 + 1.59 \cdot \ln(3n-4) - 30.36)}{(\ln(3n-4))^5}
$$

 $\textcircled{2}$  Springer

holds for any *n* > 4/3. Consequently, since  $n \ge 25$  (and thus  $\ln(3n - 4) > 0$ ) is true, we have

$$
\frac{d}{dn} f_1(3n-4) < 1
$$
\n
$$
\iff 3 \cdot \left( (\ln(3n-4))^4 + 1.59 \cdot \ln(3n-4) - 30.36 \right) < (\ln(3n-4))^5.
$$

Setting  $x := \ln(3n - 4)$  it remains to show

$$
x^5 - 3x^4 - 4.77x + 91.08 > 0,
$$

where  $x > 4$  can be assumed due to  $x \geq \ln(3 \cdot 25 - 4) \approx 4.26$ . Now, we end up with

<span id="page-19-6"></span>
$$
x^5 - 3x^4 - 4.77x + 91.08 > x^5 - 3x^4 - 5x = x(x^4 - 3x^3 - 5) > 0 \Longleftrightarrow x^4 - 3x^3 > 5
$$
\n(7)

because  $x > 4$  is true. But the final condition in [\(7\)](#page-19-6) is satisfied since we can state

$$
x^4 - 3x^3 = x^3(x - 3) > 4^3 \cdot (4 - 3) > 5
$$

for  $x > 4$ . This finally shows

$$
\frac{\mathrm{d}}{\mathrm{d}n}f_1(3n-4) < 1
$$

for all  $n \ge 25$ , so that the left hand side of [\(5\)](#page-18-0) grows more slowly than the right hand side. Together with  $\widetilde{U}(71) \leq 25 - \frac{1}{2}$  this shows [\(5\)](#page-18-0) for any  $n \geq 25$  and we are done.

**Remark 7** Note that this proof strategy cannot simply be overtaken for the case of real-valued parameters  $\delta$  since we would be confronted with the non-differentiable term  $\lceil \delta(n-1) - 1 \rceil$ , so that the chain rule cannot be applied to  $f_1$ .

## **References**

- <span id="page-19-4"></span>Bar-Noy A, Ladner RE, Tami T (2007) Windows scheduling as a restricted version of bin packing. ACM Trans Algorithms 3(3), Article 28
- <span id="page-19-3"></span>Caprara A, Dell'Amico M, Diaz-Diaz JC, Iori M, Rizzi R (2015) Friendly bin packing instances without integer round-up property. Math Program 150:5–17
- <span id="page-19-1"></span>Chen Y, Song X, Ouelhadj D, Cui Y (2019) A heuristic for the skiving and cutting stock problem in paper and plastic film industries. Int Trans Op Res 26(1):157–179

<span id="page-19-5"></span>Coffman EG, Garey MR, Johnson DS (1987) Bin packing with divisible item sizes. J Complex 3(4):406–428

<span id="page-19-2"></span>Delorme M, Iori M (2020) Enhanced pseudo-polynomial formulations for bin packing and cutting stock problems. INFORMS J Comput 32(1):101–119

<span id="page-19-0"></span>Delorme M, Iori M, Martello S (2015) Bin packing and cutting stock problems: mathematical models and exact algorithms. Research report OR-15-1, University of Bologna

- <span id="page-20-4"></span>Delorme M, Iori M, Martello S (2016) Bin packing and cutting stock problems: mathematical models and exact algorithms. Eur J Op Res 255:1–20
- <span id="page-20-28"></span>Dusart P (2018) Explicit estimates of some functions over primes. Ramanujan J 45:227–251
- <span id="page-20-9"></span>Dyckhoff H (1981) A new linear approach to the cutting stock problem. Op Res 29(6):1092–1104 Fieldhouse M (1990) The duality gap in trim problems. SICUP Bull 5:4–5
- <span id="page-20-13"></span><span id="page-20-0"></span>Gilmore PC, Gomory RE (1961) A linear programming approach to the cutting-stock problem (part I). Op Res 9:849–859
- <span id="page-20-2"></span>Johnson MP, Rennick C, Zak EJ (1997) Skiving addition to the cutting stock problem in the paper industry. SIAM Rev 39(3):472–483
- <span id="page-20-1"></span>Kantorovich LV (1939) Mathematical methods of organizing and planning production. Manag Sci 6:366– 422 (1939 Russian, 1960 English)
- <span id="page-20-17"></span>Kartak V, Ripatti A (2019) Large proper gaps in bin packing and dual bin packing problems. J Glob Optim 74:467–476
- <span id="page-20-11"></span>Kartak V, Ripatti A, Scheithauer G, Kurz S (2015) Minimal proper non-IRUP instances of the onedimensional cutting stock problem. Discrete Appl Math 187:120–129
- <span id="page-20-25"></span>Marcotte O (1983) Topics in combinatorial packing and covering. Technical report No. 568, Cornell University
- <span id="page-20-12"></span>Marcotte O (1986) An instance of the cutting stock problem for which the rounding property does not hold. Op Res Lett 4(5):239–243
- <span id="page-20-6"></span>Martinovic J, Delorme M, Iori M, Scheithauer G, Strasdat N (2020) Improved flow-based formulations for the skiving stock problem. Comput Op Res 113, Article 104770
- <span id="page-20-7"></span>Martinovic J, Scheithauer G (2016) Integer linear programming models for the skiving stock problem. Eur J Op Res 251(2):356–368
- <span id="page-20-20"></span>Martinovic J, Scheithauer G (2016) Integer rounding and modified integer rounding for the skiving stock problem. Discrete Optim 21:118–130
- <span id="page-20-23"></span>Martinovic J, Scheithauer G (2017) An upper bound of  $\Delta(E) < 3/2$  for skiving stock instances of the divisible case. Discrete Appl Math 229:161–167
- <span id="page-20-24"></span>Martinovic J, Scheithauer G (2018) Characterizing IRDP-instances of the skiving stock problem by means of polyhedral theory. Optimization 67(10):1797–1817
- <span id="page-20-26"></span>Martinovic J, Scheithauer G (2018) Combinatorial investigations on the maximum gap for skiving stock instances of the divisible case. Ann Op Res 271(2):811–829
- <span id="page-20-27"></span>Martinovic J, Scheithauer G (2019) New theoretical investigations on the gap of the skiving stock problem. Pesquisa Operacional 39(1):1–35
- <span id="page-20-10"></span>Martinovic J, Scheithauer G, Valério de Carvalho JM (2018) A comparative study of the arcflow model and the one-cut model for one-dimensional cutting stock problems. Eur J Op Res 266(2):458–471
- <span id="page-20-16"></span>Nemhauser GL, Park S (1991) A polyhedral approach to edge coloring. Op Res Lett 10:315–322
- <span id="page-20-14"></span>Nica V (1994) A general counterexample to the integer round-up property. Working paper, Department of Economic Cybernetics, Academy of Economic Studies, Bucharest
- <span id="page-20-21"></span>Rietz J (2003) Untersuchungen zu MIRUP für Vektorpackprobleme. PhD thesis, TU Bergakademie Freiberg
- <span id="page-20-18"></span>Rietz J, Dempe S (2008) Large gaps in one-dimensional cutting stock problems. Discrete Appl Math 156(10):1929–1935
- <span id="page-20-19"></span>Rietz J, Scheithauer G (2002) Families of non-IRUP instances of the one-dimensional cutting stock problem. Discrete Appl Math 121:229–245
- <span id="page-20-22"></span>Rietz J, Scheithauer G, Terno J (2002) Tighter bounds for the gap and non-IRUP constructions in the one-dimensional cutting stock problem. Optimization 51(6):927–963
- <span id="page-20-8"></span>Scheithauer G (2018) Introduction to cutting and packing optimization-problems, modeling approaches, solution methods. International series in operations research and management science 263, vol 1. Springer, Berlin
- <span id="page-20-15"></span>Scheithauer G, Terno J (1995) The modified integer round-up property of the one-dimensional cutting stock problem. Eur J Op Res 84:562–571
- <span id="page-20-5"></span>Valério de Carvalho JM (2002) LP models for bin packing and cutting stock problems. Eur J Op Res 141(2):253–273
- <span id="page-20-3"></span>Zak EJ (2003) The skiving stock problem as a counterpart of the cutting stock problem. Int Trans Op Res 10:637–650

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.