

A Service of

PRIII

Leibniz-Informationszentrum Wirtschaft Leibniz Information Centre for Economics

Biørn, Erik

Working Paper The efficiency of panel data estimators: GLS versus estimators which do not depend on variance components

Memorandum, No. 2001,28

Provided in Cooperation with: Department of Economics, University of Oslo

Suggested Citation: Biørn, Erik (2001) : The efficiency of panel data estimators: GLS versus estimators which do not depend on variance components, Memorandum, No. 2001,28, University of Oslo, Department of Economics, Oslo

This Version is available at: <https://hdl.handle.net/10419/63055>

Standard-Nutzungsbedingungen:

Die Dokumente auf EconStor dürfen zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden.

Sie dürfen die Dokumente nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, öffentlich zugänglich machen, vertreiben oder anderweitig nutzen.

Sofern die Verfasser die Dokumente unter Open-Content-Lizenzen (insbesondere CC-Lizenzen) zur Verfügung gestellt haben sollten, gelten abweichend von diesen Nutzungsbedingungen die in der dort genannten Lizenz gewährten Nutzungsrechte.

Terms of use:

Documents in EconStor may be saved and copied for your personal and scholarly purposes.

You are not to copy documents for public or commercial purposes, to exhibit the documents publicly, to make them publicly available on the internet, or to distribute or otherwise use the documents in public.

If the documents have been made available under an Open Content Licence (especially Creative Commons Licences), you may exercise further usage rights as specified in the indicated licence.

WWW.ECONSTOR.EU

MEMORANDUM

No 28/2001

The Efficiency of Panel Data Estimators: GLS Versus Estimators Which Do Not Depend on Variance Components

> *By Erik Biørn*

ISSN: 0801-1117

Department of Economics University of Oslo

List of the latest 10 Memoranda:

A complete list of this memo-series is available in a PDF® format at: http://www.oekonomi.uio.no/memo/

THE EFFICIENCY OF PANEL DATA ESTIMATORS: GLS VERSUS ESTIMATORS WHICH DO NOT DEPEND ON VARIANCE COMPONENTS

by

ERIK BIØRN

ABSTRACT

For a balanced panel data regression model with random effects, we discuss the efficiency of the GLS estimator relative to the OLS, the between, and the within estimators. Focus is on how the efficiency responds to changes in (a) the relative variance components and (b) the composition of the regressor covariance matrix which into between and within variation. Both one-way and two-way models are considered. For the one-way, one regressor model, we show that (i) OLS has maximal inefficiency relative to GLS when the within and between individual variation in the regressor account for the same part of the total variation, (ii) the between estimator is always less efficient than the OLS estimator. For the two-way, one regressor model, the between individual (between period) estimator is more efficient than OLS if the between period (between individual) share of the total variation in the regressor and/or the time specific (individual specific) disturbance variance component are sufficiently large. Illustrations relating to marginal budget shares in household consumption are given.

Keywords: Panel Data. Variance Components. Efficiency. Generalized Least Squares. Within estimation. Between estimation

JEL classification: C13, C23

1 Introduction

According to a familiar textbook result in econometrics, the Generalized Least Squares (GLS) estimator of a regression coefficient vector in the case with a non-scalar disturbance covariance matrix is Minimum Variance Linear Unbiased [Greene (2000, section 11.3). When applied to a panel data model with random effects, this extension of Gauss-Markov's theorem implies that the GLS is more efficient than the Within (W), the Between (B), and the Ordinary Least Squares (OLS) estimators. Strict GLS, however, is an `impracticable' method, as it presumes knowledge of the (relative) disturbance variances, which is rarely available, and the efficiency of *Feasible* GLS relative to W, B, and $0\,\mathrm{LS}$ when the variance components are estimated from residuals from a finite (and often small) number of units and/or periods, is not in general known. Some results are available, though [Swamy and Arora (1972), Maddala and Mount (1973), Taylor (1980), and Baltagi (1981)], *inter alia*, based on Monte Carlo simulations when considering alternative ways of estimating the variance components, but they cannot be easily summarized.

In this paper, we investigate, for a balanced panel data model, the most important determinants of the efficiency of strict GLS relative to estimators which do not presume the disturbance variances to be known, or estimated. To the author's knowledge, this issue has not been discussed in the literature, apart from certain special cases. Focus is on how the efficiency of GLS and other panel data estimators responds to changes in (a) the relative variances of the disturbance components, (b) the number of individuals and periods, and (c) the composition of the regressor covariance matrix in between and within variation. Both one-way and two-way models of the latent heterogeneity are considered.

We examine on the one hand the inefficiency of W, B, and OLS relative to GLS, on the other hand the relative efficiency of W, B, and OLS. Since strict GLS may be unobtainable, choosing between W, B, and OLS is of considerable practical interest, not least from the point of view of robustness. First, the consistency of W, unlike the consistency of OLS, is robust to correlation between the latent random effects and the covariate vector [see Hsiao (1986, section 3.4)]. Second, even if OLS often outperforms B in terms of finite sample efficiency, the consistency of the between individual (between period) estimator, unlike the OLS, is robust to errors of measurement in the regressors when the number of periods (individuals) goes to infinity [see Bigrn $(1996, \text{section } 10.2.3)$].

The paper is organized as follows. In Section 2, we define four estimators for the oneway random effects model and derive general efficiency results for a regression model of arbitrary dimension. More detailed results are given for the one regressor (or orthogonal regressors) case. We find that OLS is maximally inefficient relative to GLS when half of the variation in the regressor is within individual and the other half is between individual variation. For this model, we show that the between individual estimator is less efficient than OLS, since the latter exploits both between and within variation in the data.

In Section 3 we extend the analysis to the two-way model and present five estimators. We show, for the one regressor case, that OLS is most inefficient relative to GLS (i) in the presence of between period variation in the regressor when less than half of the variation in the regressor is between individual variation, and (ii) in the presence of between individual variation in the regressor when less than half of the variation in the regressor is between period variation. While for the one-way model the between estimator is uniformly less efficient than OLS, this is not generally true for the two-way model. In the one regressor case, we find for the latter that the *between individual (between period)* estimator is more efficient than OLS if the *between period (between individual)* share of the total variation in the regressor and/or the *time specific (individual specific)* part of the disturbance variance exceeds a certain level. The ranking of the strict OLS and the twoway fixed effects OLS (double within) estimator is also discussed. Empirical illustrations relating to marginal budget shares in household consumption are given. Concluding remarks follow in Section 4.

2 Results for the one-way model

2.1Model and estimators

Assume that we have a panel data set in which N (\geq 2) individuals are observed in T (\geq 2) periods, and consider the model

(1)
$$
y_{it} = k + \boldsymbol{x}_{it}\boldsymbol{\beta} + \alpha_i + u_{it},
$$

$$
\alpha_i \sim \mathsf{IID}(0, \sigma_{\alpha}^2), \quad u_{it} \sim \mathsf{IID}(0, \sigma^2), \quad i = 1, ..., N,
$$

$$
\alpha_i, u_{it}, \boldsymbol{x}_{it} \text{ are independent for all } i, t, \quad t = 1, ..., T,
$$

where i indicates individual, t indicates period, x_{it} is a (row) vector of regressors, β its (column) vector of coefficients, α_i is an individual specific random effect, and u_{it} is a disturbance. Let $\bm{y}_i = (y_{i1}, \dots, y_{iT})\,, \ \bm{y} = (\bm{y}_1, \dots, \bm{y}_N) \,, \ \bm{\Lambda}_i = (\bm{x}_{i1}, \dots, \bm{x}_{iT})\,, \bm{\Lambda}\, =$ $(\bm{\Lambda}_1,\ldots,\bm{\Lambda}_N)$, etc, $\bm{\alpha}=(\alpha_1,\ldots,\alpha_N)$, and \bm{e}_m be the (m x 1) vector of ones. Compactly, the model can then be written

(2)
$$
y = e_{NT}k + X\beta + \epsilon, \quad \epsilon = e_T \otimes \alpha + u,
$$

$$
E(\epsilon) = 0, \quad V(\epsilon) = E(\epsilon \epsilon') = \Omega,
$$

where

(3)
$$
\mathbf{\Omega} = \mathbf{I}_N \otimes (\sigma_\alpha^2 \mathbf{e}_T \mathbf{e}_T' + \sigma^2 \mathbf{I}_T) = \mathbf{I}_N \otimes [\sigma^2 \mathbf{K}_T + (\sigma^2 + T \sigma_\alpha^2) \mathbf{J}_T],
$$

 \otimes is the Kronecker product operator, $\bm{J}_m = (\bm{e}_m \bm{e}_m)/m$ and $\bm{\Lambda}_m = \bm{I}_m - \bm{J}_m, \; m = 0$ $1, 2, \ldots$. The latter matrices are idempotent and have orthogonal columns.

We use the following notation for the within individual, the between individual, and the total covariation in arbitrary matrices of panel data, Z and Q , constructed in the same way as \boldsymbol{X} above:

$$
\mathbf{W}_{ZQ} = \mathbf{Z}'(\mathbf{I}_N \otimes \mathbf{K}_T)\mathbf{Q} = \sum_{i=1}^N \sum_{t=1}^T (z_{it} - \bar{z}_{i\cdot})'(\mathbf{q}_{it} - \bar{\mathbf{q}}_{i\cdot}),
$$

\n
$$
\mathbf{B}_{ZQ} = \mathbf{Z}'(\mathbf{K}_N \otimes \mathbf{J}_T)\mathbf{Q} = T \sum_{i=1}^N (\bar{z}_{i\cdot} - \bar{z})'(\bar{\mathbf{q}}_{i\cdot} - \bar{\mathbf{q}}),
$$

\n
$$
\mathbf{T}_{ZQ} = \mathbf{Z}'(\mathbf{I}_{NT} - \mathbf{J}_{NT})\mathbf{Q} = \sum_{i=1}^N \sum_{t=1}^T (z_{it} - \bar{z})'(\mathbf{q}_{it} - \bar{\mathbf{q}}) = \mathbf{W}_{ZQ} + \mathbf{B}_{ZQ},
$$

where \bar{z}_i , $= T^{-1} \sum_{t=1}^l z_{it}$, $\bar{z} = (NT)^{-1} \sum_{i=1}^N \sum_{t=1}^l z_{it}$, etc. The columns of W_{ZO} and \bm{B}_{ZQ} are orthogonal since \bm{K}_T and \bm{J}_T have this property. Four estimators of $\bm{\beta},$ familiar from the panel data literature [see, e.g., Hsiao (1986, chapter 3)], are considered:

$$
(4) \qquad \qquad \widehat{\boldsymbol{\beta}}_W = \boldsymbol{W}_{XX}^{-1} \boldsymbol{W}_{XY},
$$

$$
\widehat{\boldsymbol{\beta}}_B = \boldsymbol{B}_{XX}^{-1} \boldsymbol{B}_{XY},
$$

(6)
$$
\hat{\beta}_{OLS} = T_{XX}^{-1} T_{XY} = (W_{XX} + B_{XX})^{-1} (W_{XY} + B_{XY}),
$$

(*i*) $\mathbf{p}_{GLS} = (\mathbf{W}_{XX} + \mathbf{v}_B \mathbf{B}_{XX})$ ($\mathbf{W}_{XY} + \mathbf{v}_B \mathbf{B}_{XY}$),

where

(8)
$$
\theta_B = \frac{\sigma^2}{\sigma^2 + T\sigma_\alpha^2}.
$$

Here $\rho_W,$ the within individual estimator, is the Gauss-Markov estimator if the individual specific effects α_i are treated as fixed and unknown, ρ_{GLS} is the Gauss-Markov estimator if they are treated as random, ρ_{OLS} is the Gauss-Markov estimator when no heterogeneity occurs, and $\rho_B,$ the between individual estimator, is the OLS estimator constructed from individual specific means of the observations [see Hsiao (1986, section 3.3.2)]. The full within estimator will only exist when no regressor is time invariant, since otherwise W_{XX} has zero rows and columns.

Let λ_W and λ_B be non-negative scalar constants and define the more general estimator

(9)
$$
\hat{\boldsymbol{\beta}} = \hat{\boldsymbol{\beta}}(\lambda_W, \lambda_B) = [\lambda_W \boldsymbol{W}_{XX} + \lambda_B \boldsymbol{B}_{XX}]^{-1} [\lambda_W \boldsymbol{W}_{XY} + \lambda_B \boldsymbol{B}_{XY}].
$$

Obviously,

$$
\begin{array}{rcl}\n\widehat{\beta}_W & = & \widehat{\beta}(1,0), \\
\widehat{\beta}_B & = & \widehat{\beta}(0,1), \\
\widehat{\beta}_{OLS} & = & \widehat{\beta}(1,1), \\
\widehat{\beta}_{GLS} & = & \widehat{\beta}(1,\theta_B).\n\end{array}
$$

Inserting for y from (2) in (9) , we obtain

(10)
$$
\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta} = [\lambda_W \boldsymbol{W}_{XX} + \lambda_B \boldsymbol{B}_{XX}]^{-1} [\lambda_W \boldsymbol{W}_{X\epsilon} + \lambda_B \boldsymbol{B}_{X\epsilon}].
$$

2.2General efficiency results

From (2) and (3) we obtain

$$
E(\boldsymbol{W}_{X\epsilon}\boldsymbol{W}_{X\epsilon}') = E([X'(I_N \otimes \boldsymbol{K}_T)\epsilon][X'(I_N \otimes \boldsymbol{K}_T)\epsilon]') = X'(I_N \otimes \boldsymbol{K}_T)\Omega(I_N \otimes \boldsymbol{K}_T)X,
$$

\n
$$
E(\boldsymbol{B}_{X\epsilon}\boldsymbol{B}_{X\epsilon}') = E([X'(K_N \otimes \boldsymbol{J}_T)\epsilon][X'(K_N \otimes \boldsymbol{J}_T)\epsilon]') = X'(K_N \otimes \boldsymbol{J}_T)\Omega(K_N \otimes \boldsymbol{J}_T)X,
$$

leading to

$$
E(\boldsymbol{W}_{X\epsilon}\boldsymbol{W}_{X\epsilon}') = \sigma^2 \boldsymbol{X}'(\boldsymbol{I}_N \otimes \boldsymbol{K}_T) \boldsymbol{X} = \sigma^2 \boldsymbol{W}_{XX},
$$

\n
$$
E(\boldsymbol{B}_{X\epsilon}\boldsymbol{B}_{X\epsilon}') = (\sigma^2 + T\sigma^2_{\alpha}) \boldsymbol{X}'(\boldsymbol{K}_N \otimes \boldsymbol{J}_T) \boldsymbol{X} = (\sigma^2 + T\sigma^2_{\alpha}) \boldsymbol{B}_{XX}.
$$

Combining these expressions with (10) it follows that the covariance matrix of \bm{p} (conditional on X), is

(11)
$$
\mathsf{V}(\hat{\boldsymbol{\beta}}) = \mathsf{E}[(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})']
$$

$$
= [\lambda_W \mathbf{W}_{XX} + \lambda_B \mathbf{B}_{XX}]^{-1} [\lambda_W^2 \sigma^2 \mathbf{W}_{XX} + \lambda_B^2 (\sigma^2 + T \sigma_{\alpha}^2) \mathbf{B}_{XX}]
$$

$$
\times [\lambda_W \mathbf{W}_{XX} + \lambda_B \mathbf{B}_{XX}]^{-1}.
$$

This expression can be used to rank unbiased estimators with different (λ_W, λ_B) constellations. Sometimes, one estimator, p_1 , is uniformly superior to another, p_2 , if $\mathbf{v}(\mathbf{\mu}_2)$ – $\mathbf{v}(\mathbf{\mu}_1)$ is positive definite for any $\mathbf{\Lambda}$. In particular, we have

(12)
$$
\mathsf{V}(\hat{\boldsymbol{\beta}}_W) = \left[\frac{\boldsymbol{W}_{XX}}{\sigma^2}\right]^{-1},
$$

(13)
$$
\mathsf{V}(\hat{\boldsymbol{\beta}}_B) = \left[\frac{\boldsymbol{B}_{XX}}{\sigma^2 + T\sigma_{\alpha}^2}\right]^{-1},
$$

(14)
\n
$$
\mathsf{V}(\hat{\boldsymbol{\beta}}_{OLS}) = \mathbf{T}_{XX}^{-1} [\sigma^2 \mathbf{W}_{XX} + (\sigma^2 + T\sigma_{\alpha}^2) \mathbf{B}_{XX}] \mathbf{T}_{XX}^{-1}
$$
\n
$$
= \left[\frac{\mathbf{T}_{XX}}{\sigma^2} \right]^{-1} + T\sigma_{\alpha}^2 \mathbf{T}_{XX}^{-1} \mathbf{B}_{XX} \mathbf{T}_{XX}^{-1},
$$
\n(15)
\n
$$
\mathsf{V}(\hat{\boldsymbol{\beta}}_{GLS}) = \left[\frac{\mathbf{W}_{XX}}{\sigma^2} + \frac{\mathbf{B}_{XX}}{\sigma^2 + T\sigma_{\alpha}^2} \right]^{-1}.
$$

 $\sigma^2 + T \sigma_{\alpha}^2$

Using Magnus and Neudecker (1988, chapter 1, Theorem 24), it follows that

$$
V(\hat{\beta}_W) - V(\hat{\beta}_{GLS}) \text{ is pos. def.} \iff \frac{B_{XX}}{\sigma^2 + T\sigma^2_{\alpha}} \text{ is pos. def.},
$$

$$
V(\hat{\beta}_B) - V(\hat{\beta}_{GLS}) \text{ is pos. def.} \iff \frac{W_{XX}}{\sigma^2} \text{ is pos. def.}
$$

I ms implies that ρ_{GLS} is strictly more emclent than both ρ_W and ρ_B when both \bm{D}_{XX} and W_{XX} are positive definite and σ^2 and σ^2_{α} are positive and finite and T is finite. r urthermore, we can write ${\bf v}({\boldsymbol \rho}_{OLS}) - {\bf v}({\boldsymbol \rho}_{GLS})$ as a matrix product which can be shown [for instance by using results in Horn and Johnson (1985, section 7.6)] to be positive denite, in agreement with Gauss-Markov's theorem.

2.3GLS versus OLS: The one regressor case

Consider the relative efficiency of OLS and GLS in the one regressor case, $K = 1$. The results we derive below are not strictly confined to this case, however; they are also valid, for each regression coefficient, if all regressions are orthogonal so that \equiv Λ Λ Γ \equiv Λ Λ Γ \boldsymbol{W}_{XX} are diagonal matrices. Let, for $K=1,$ $b=\boldsymbol{B}_{XX}/\boldsymbol{T}_{XX},$ $i.e.,$ the share of the total variation in the regressor (or a typical regressor, under orthogonality) which is between individual variation. It then follows from (8), (14), and (15) that

$$
\begin{array}{rcl}\ne_{OLS} & = & \frac{\mathsf{V}(\hat{\beta}_{OLS})}{\mathsf{V}(\hat{\beta}_{GLS})} \\
 & = & \left[\sigma^2 (1-b) + (\sigma^2 + T \sigma_\alpha^2) b \right] \left[\frac{1-b}{\sigma^2} + \frac{b}{\sigma^2 + T \sigma_\alpha^2} \right] \\
 & = & (1-b)^2 + b(1-b) \left(\theta_B + \frac{1}{\theta_B} \right) + b^2,\n\end{array}
$$

i.e.,

(16)
$$
e_{OLS} = \left[1 + b\left(\theta_B + \frac{1}{\theta_B} - 1\right)\right](1 - b) + b^2.
$$

Since

$$
\frac{\partial e_{OLS}}{\partial \theta_B} = b(1 - b) \left(1 - \frac{1}{\theta_B^2} \right),
$$

$$
\frac{\partial e_{OLS}}{\partial b} = (1 - \theta_B) \left(\frac{1}{\theta_B} - 1 \right) (1 - 2b),
$$

the efficiency function (16) has the following properties: (i) It is convex in θ_B when $0 < b < 1$ and attains its minimum, one, for $\theta_B = 1$, *i.e.*, $\sigma_{\alpha}^2 = 0$ (which is obvious since OLS and GLS coincide in this case). (ii) It is concave in b when $\theta_B < 1$ and attains its maximum, $(1/4)(1 + \theta_B)(1/\theta_B + 1)$, for $\theta = \frac{1}{2}$. This means that β_{OLS} , for given T and σ_{α} (> 0) is maximally inemicient relative to ρ_{GLS} when half of the variation in the regressor is within individual and the other half is between individual variation. When $b = 0$ or $b = 1$, $e_{OLS} = 1$ for any θ_B .

2.4The ranking of the within, between, and OLS estimators

we see from (12) = (14) that the ranking of μ_B , ρ_W , and ρ_{OLS} depends in the general case on σ , I σ_{α} , \boldsymbol{B} XX , and W XX . The one regressor (or orthogonal regressors) case is most transparent, and we consider this case specifically. From (8) and $(12) - (14)$ we

obtain the following variance ratios when $K = 1$:

$$
\frac{\mathsf{V}(\beta_W)}{\mathsf{V}(\hat{\beta}_B)} = \frac{b\theta_B}{1-b} \qquad > (<) 1 \iff 1-b < (>) \frac{\theta_B}{1+\theta_B},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_W)}{\mathsf{V}(\hat{\beta}_{OLS})} = \frac{1}{(1-b)\left[1+b\left(\frac{1}{\theta_B}-1\right)\right]} > (<) 1 \iff 1-b < (>) \frac{\theta_B}{1-\theta_B},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_B)}{\mathsf{V}(\hat{\beta}_{OLS})} = \frac{1}{b\theta_B\left[1+b\left(\frac{1}{\theta_B}-1\right)\right]} > 1 \iff 0 < b < 1.
$$

If $0 \lt vB \lt 1$ and $0 \lt 0 \lt 1$, then μ_B is always less emicient than μ_{OLS}, μ_{OLS} is less emerent than ρ_{GLS} , and $\mathbf{v}(\rho_B)$ $\mathbf{v}(\rho_{OLS})$ is monotomeany declining in v . The ranking or ρ_W depends on $v.$

We find, by inserting from (8) , that the relative efficiency of the four estimators depends on the share of the variation in the regressor which is within individual variation, $1-b$, as follows [these inequalities generalize results in Malinvaud (1978, chapter 8.4(ii))]:

$$
V(\hat{\beta}_B) > V(\hat{\beta}_{OLS}) \ge V(\hat{\beta}_W) > V(\hat{\beta}_{GLS}) \quad \text{if} \quad \frac{\sigma^2}{T\sigma_{\alpha}^2} \le 1 - b < 1,
$$

\n
$$
V(\hat{\beta}_B) \ge V(\hat{\beta}_W) > V(\hat{\beta}_{OLS}) > V(\hat{\beta}_{GLS}) \quad \text{if} \quad \frac{\sigma^2}{2\sigma^2 + T\sigma_{\alpha}^2} \le 1 - b < \frac{\sigma^2}{T\sigma_{\alpha}^2},
$$

\n
$$
V(\hat{\beta}_W) > V(\hat{\beta}_B) > V(\hat{\beta}_{OLS}) > V(\hat{\beta}_{GLS}) \quad \text{if} \quad 0 < 1 - b < \frac{\sigma^2}{2\sigma^2 + T\sigma_{\alpha}^2}.
$$

The larger T or $\sigma_{\alpha}^2/\sigma^2$ for given b is, the more likely is the first inequality to be satisfied (ii, at the limit, $I \rightarrow \infty$, then $\theta_B \rightarrow \infty$, so that ρ_W and ρ_{GLS} coincide). The larger $v,$ for given T and $\sigma_{\alpha}^>/\sigma^-,$ is, the more likely is the last inequality to be satisfied (ii, at the finit, $v \to 1$, then ρ_B , ρ_{OLS} and ρ_{GLS} coincide and ρ_W is undefined). From the point of view of robustness, the first inequality in (17) is particularly interesting: If $1 - \theta > \sigma^2/(1 \sigma^2)$, then β_W is not only more efficient than β_{OLS} , it is also robust to violation of the assumption that α_i and \boldsymbol{x}_{it} are independent and neither depends on variance components. If $1 - b < \sigma^2/(T \sigma_\alpha^2)$, there will be a trade-off between efficiency and robustness for these two estimators.

3 Results for the two-way model

We now extend the analysis to the two-way model.

3.1Model and estimators

Consider the model

(18)
$$
y_{it} = k + \boldsymbol{x}_{it}\boldsymbol{\beta} + \alpha_i + \gamma_t + u_{it},
$$

$$
\alpha_i \sim \text{IID}(0, \sigma_{\alpha}^2), \quad \gamma_t \sim \text{IID}(0, \sigma_{\gamma}^2), \quad u_{it} \sim \text{IID}(0, \sigma^2), \qquad i = 1, ..., N,
$$

$$
\alpha_i, \gamma_t, u_{it}, \boldsymbol{x}_{it} \text{ are independent for all } i, t,
$$

where γ_t is a period specific random effect and the other symbols have the same interpretation as in Section 2. Compactly, the model can be written as

(19)
$$
y = e_{NT}k + X\beta + \epsilon, \quad \epsilon = (\alpha \otimes e_T) + (e_N \otimes \gamma) + u,
$$

$$
E(\epsilon) = 0, \quad V(\epsilon) = \Omega,
$$

where $\gamma = (\gamma_1, \ldots, \gamma_T)$ and

(20)
$$
\Omega = \sigma_{\alpha}^{2} \mathbf{I}_{N} \otimes (\mathbf{e}_{T} \mathbf{e}_{T}^{\prime}) + \sigma_{\gamma}^{2} (\mathbf{e}_{N} \mathbf{e}_{N}^{\prime}) \otimes \mathbf{I}_{T} + \sigma^{2} (\mathbf{I}_{N} \otimes \mathbf{I}_{T})
$$

\n
$$
= \sigma^{2} (\mathbf{K}_{N} \otimes \mathbf{K}_{T}) + (\sigma^{2} + T \sigma_{\alpha}^{2}) (\mathbf{K}_{N} \otimes \mathbf{J}_{T}) + (\sigma^{2} + N \sigma_{\gamma}^{2}) (\mathbf{J}_{N} \otimes \mathbf{K}_{T})
$$

\n
$$
+ (\sigma^{2} + T \sigma_{\alpha}^{2} + N \sigma_{\gamma}^{2}) (\mathbf{J}_{N} \otimes \mathbf{J}_{T}),
$$

see Fuller and Battese (1974, section 2).

We define \bm{B}_{ZQ} and \bm{T}_{ZQ} as in Section 2 and use the following notation for the residual $(i.e., double within)$ and the between period covariation for arbitrary matrices of panel data, \boldsymbol{Z} and \boldsymbol{Q} :

$$
\mathbf{R}_{ZQ} = \mathbf{Z}'(\mathbf{K}_N \otimes \mathbf{K}_T)\mathbf{Q} = \sum_{i=1}^N \sum_{t=1}^T (z_{it} - \bar{z}_{i \cdot} - \bar{z}_{\cdot t} + \bar{z})'(q_{it} - \bar{q}_{i \cdot} - \bar{q}_{\cdot t} + \bar{q}),
$$

\n
$$
\mathbf{C}_{ZQ} = \mathbf{Z}'(\mathbf{J}_N \otimes \mathbf{K}_T)\mathbf{Q} = N \sum_{t=1}^T (\bar{z}_{\cdot t} - \bar{z})'(\bar{q}_{\cdot t} - \bar{q}),
$$

where $\bar{z}_{\cdot t} = N^{-1} \sum_{i=1}^{N} z_{it},$ $T_{ZQ} = R_{ZQ} + B_{ZQ} + C_{ZQ},$ and the columns of $R_{ZQ},$ $B_{ZQ},$ and C_{ZQ} are orthogonal. Now, five estimators of β are considered:

$$
(21) \qquad \widehat{\boldsymbol{\beta}}_R = \boldsymbol{R}_{XX}^{-1} \boldsymbol{R}_{XY},
$$

$$
(22) \qquad \widehat{\boldsymbol{\beta}}_B = \boldsymbol{B}_{XX}^{-1} \boldsymbol{B}_{XY},
$$

$$
(23) \qquad \beta_C = \mathbf{C}_{XX}^{-1} \mathbf{C}_{XY},
$$

(24)
$$
\hat{\boldsymbol{\beta}}_{OLS} = \boldsymbol{T}_{XX}^{-1} \boldsymbol{T}_{XY} = (\boldsymbol{R}_{XX} + \boldsymbol{B}_{XX} + \boldsymbol{C}_{XX})^{-1} (\boldsymbol{R}_{XY} + \boldsymbol{B}_{XY} + \boldsymbol{C}_{XY}),
$$

(25)
$$
\widehat{\boldsymbol{\beta}}_{GLS} = (\boldsymbol{R}_{XX} + \theta_B \boldsymbol{B}_{XX} + \theta_C \boldsymbol{C}_{XX})^{-1} (\boldsymbol{R}_{XY} + \theta_B \boldsymbol{B}_{XY} + \theta_C \boldsymbol{C}_{XY}),
$$

where

(26)
$$
\theta_B = \frac{\sigma^2}{\sigma^2 + T\sigma_\alpha^2}, \qquad \theta_C = \frac{\sigma^2}{\sigma^2 + N\sigma_\gamma^2}.
$$

Here ρ_R , the residual (double within) estimator, is the Gauss-Markov estimator if the individual specific and period specific enects α_i and γ_t are all treated as fixed, ρ_{GLS} is the

Gauss-Markov estimator if they are all treated as random [see Fuller and Battese (1974, section 5) and matyas (1990, section 4.2.2)], ρ_{OLS} is the Gauss-Markov estimator when no neterogeneity occurs, ρ_B , the between individual estimator is, as in Section 2, the OLS estimator constructed from individual specific means, and ρ_C^+ , the between period estimator, is the symmetric estimator constructed from period specic means. The full residual estimator will only exist when no regressor is individual or time invariant, since otherwise \mathbf{R}_{XX} has zero rows and columns.

Let λ_R , λ_B , and λ_C be non-negative scalar constants and consider the more general estimator

(27)
$$
\hat{\boldsymbol{\beta}} = \hat{\boldsymbol{\beta}}(\lambda_R, \lambda_B, \lambda_C) \n= [\lambda_R \boldsymbol{R}_{XX} + \lambda_B \boldsymbol{B}_{XX} + \lambda_C \boldsymbol{C}_{XX}]^{-1} [\lambda_R \boldsymbol{R}_{XY} + \lambda_B \boldsymbol{B}_{XY} + \lambda_C \boldsymbol{C}_{XY}],
$$

which also generalizes (9) since latter corresponds to $\lambda_R = \lambda_C = \lambda_W$. Obviously,

$$
\begin{array}{rcl}\n\widehat{\beta}_R & = & \widehat{\beta}(1,0,0), \\
\widehat{\beta}_B & = & \widehat{\beta}(0,1,0), \\
\widehat{\beta}_C & = & \widehat{\beta}(0,0,1), \\
\widehat{\beta}_{OLS} & = & \widehat{\beta}(1,1,1), \\
\widehat{\beta}_{GLS} & = & \widehat{\beta}(1,\theta_B,\theta_C).\n\end{array}
$$

Inserting for y from (19) in (27), we obtain

(28)
$$
\hat{\boldsymbol{\beta}} - \boldsymbol{\beta} = [\lambda_R \boldsymbol{R}_{XX} + \lambda_B \boldsymbol{B}_{XX} + \lambda_C \boldsymbol{C}_{XX}]^{-1} [\lambda_R \boldsymbol{R}_{X\epsilon} + \lambda_B \boldsymbol{B}_{X\epsilon} + \lambda_C \boldsymbol{C}_{X\epsilon}].
$$

3.2General efficiency results

From (19) and (20) we obtain

$$
\mathsf{E}(\mathbf{R}_{X\epsilon}\mathbf{R}_{X\epsilon}')=\mathsf{E}([\mathbf{X}'(\mathbf{K}_{N} \otimes \mathbf{K}_{T})\epsilon][\mathbf{X}'(\mathbf{K}_{N} \otimes \mathbf{K}_{T})\epsilon]')=\mathbf{X}'(\mathbf{K}_{N} \otimes \mathbf{K}_{T})\Omega(\mathbf{K}_{N} \otimes \mathbf{K}_{T})\mathbf{X},
$$

\n
$$
\mathsf{E}(\mathbf{B}_{X\epsilon}\mathbf{B}_{X\epsilon}')=\mathsf{E}([\mathbf{X}'(\mathbf{K}_{N} \otimes \mathbf{J}_{T})\epsilon][\mathbf{X}'(\mathbf{K}_{N} \otimes \mathbf{J}_{T})\epsilon]')=\mathbf{X}'(\mathbf{K}_{N} \otimes \mathbf{J}_{T})\Omega(\mathbf{K}_{N} \otimes \mathbf{J}_{T})\mathbf{X},
$$

\n
$$
\mathsf{E}(\mathbf{C}_{X\epsilon}\mathbf{C}_{X\epsilon}')=\mathsf{E}([\mathbf{X}'(\mathbf{J}_{N} \otimes \mathbf{K}_{T})\epsilon][\mathbf{X}'(\mathbf{J}_{N} \otimes \mathbf{K}_{T})\epsilon]')=\mathbf{X}'(\mathbf{J}_{N} \otimes \mathbf{K}_{T})\Omega(\mathbf{J}_{N} \otimes \mathbf{K}_{T})\mathbf{X},
$$

leading to

$$
\mathsf{E}(\mathbf{R}_{X\epsilon}\mathbf{R}'_{X\epsilon}) = \sigma^2 \mathbf{X}'(\mathbf{K}_N \otimes \mathbf{K}_T)\mathbf{X} = \sigma^2 \mathbf{R}_{XX},
$$
\n
$$
\mathsf{E}(\mathbf{B}_{X\epsilon}\mathbf{B}'_{X\epsilon}) = (\sigma^2 + T\sigma^2_{\alpha})\mathbf{X}'(\mathbf{K}_N \otimes \mathbf{J}_T)\mathbf{X} = (\sigma^2 + T\sigma^2_{\alpha})\mathbf{B}_{XX},
$$
\n
$$
\mathsf{E}(\mathbf{C}_{X\epsilon}\mathbf{C}'_{X\epsilon}) = (\sigma^2 + N\sigma^2_{\gamma})\mathbf{X}'(\mathbf{J}_N \otimes \mathbf{K}_T)\mathbf{X} = (\sigma^2 + N\sigma^2_{\gamma})\mathbf{C}_{XX}.
$$

Combining these expressions with (28) it follows that

(29)
$$
V(\hat{\boldsymbol{\beta}}) = E[(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})(\hat{\boldsymbol{\beta}} - \boldsymbol{\beta})']
$$

\n
$$
= [\lambda_R \mathbf{R}_{XX} + \lambda_B \mathbf{B}_{XX} + \lambda_C \mathbf{C}_{XX}]^{-1}
$$

\n
$$
\times [\lambda_R^2 \sigma^2 \mathbf{R}_{XX} + \lambda_B^2 (\sigma^2 + T \sigma_\alpha^2) \mathbf{B}_{XX} + \lambda_C^2 (\sigma^2 + N \sigma_\gamma^2) \mathbf{C}_{XX}]
$$

\n
$$
\times [\lambda_R \mathbf{R}_{XX} + \lambda_B \mathbf{B}_{XX} + \lambda_C \mathbf{C}_{XX}]^{-1}.
$$

This expression can be used to rank unbiased estimators with different $(\lambda_R, \lambda_B, \lambda_C)$ constenations. Sometimes, one estimator, p_1 , is uniformly superior to another, p_2 , if $\mathbf{v}(\boldsymbol{\mu}_2)$ – $\mathbf{v}(\boldsymbol{\mu}_1)$ is positive definite for any $\boldsymbol{\Lambda}$. In particular, we have

(30)
$$
\mathsf{V}(\widehat{\boldsymbol{\beta}}_R) = \left[\frac{\boldsymbol{R}_{XX}}{\sigma^2}\right]^{-1},
$$

(31)
$$
\mathsf{V}(\hat{\boldsymbol{\beta}}_B) = \left[\frac{\boldsymbol{B}_{XX}}{\sigma^2 + T\sigma_{\alpha}^2}\right]^{-1},
$$

(32)
$$
\mathsf{V}(\hat{\boldsymbol{\beta}}_C) = \left[\frac{\boldsymbol{C}_{XX}}{\sigma^2 + N\sigma_{\gamma}^2}\right]^{-1},
$$

(33)
$$
\mathsf{V}(\hat{\boldsymbol{\beta}}_{OLS}) = \mathbf{T}_{XX}^{-1} [\sigma^2 \mathbf{R}_{XX} + (\sigma^2 + T\sigma_{\alpha}^2) \mathbf{B}_{XX} + (\sigma^2 + N\sigma_{\gamma}^2) \mathbf{C}_{XX}] \mathbf{T}_{XX}^{-1}
$$

$$
= \left[\frac{\mathbf{T}_{XX}}{\sigma^2} \right]^{-1} + \mathbf{T}_{XX}^{-1} [T\sigma_{\alpha}^2 \mathbf{B}_{XX} + N\sigma_{\gamma}^2 \mathbf{C}_{XX}] \mathbf{T}_{XX}^{-1},
$$

(34)
$$
\mathsf{V}(\hat{\boldsymbol{\beta}}_{GLS}) = \left[\frac{\boldsymbol{R}_{XX}}{\sigma^2} + \frac{\boldsymbol{B}_{XX}}{\sigma^2 + T\sigma_{\alpha}^2} + \frac{\boldsymbol{C}_{XX}}{\sigma^2 + N\sigma_{\gamma}^2} \right]^{-1}.
$$

Using Magnus and Neudecker (1988, chapter 1, Theorem 24), it follows that

$$
V(\hat{\beta}_R) - V(\hat{\beta}_{GLS}) \text{ is pos. def.} \iff \frac{B_{XX}}{\sigma^2 + T\sigma_{\alpha}^2} + \frac{C_{XX}}{\sigma^2 + N\sigma_{\gamma}^2} \text{ is pos. def.},
$$

\n
$$
V(\hat{\beta}_B) - V(\hat{\beta}_{GLS}) \text{ is pos. def.} \iff \frac{R_{XX}}{\sigma^2} + \frac{C_{XX}}{\sigma^2 + N\sigma_{\gamma}^2} \text{ is pos. def.},
$$

\n
$$
V(\hat{\beta}_C) - V(\hat{\beta}_{GLS}) \text{ is pos. def.} \iff \frac{R_{XX}}{\sigma^2} + \frac{B_{XX}}{\sigma^2 + T\sigma_{\alpha}^2}, \text{ is pos. def.}
$$

This implies that ρ_{GLS} is strictly more emclent than both ρ_R , ρ_B , and ρ_C when both $\bm{D}_{XX}, \bm{\cup}_{XX}, \bm{\alpha}_{X}$ and \bm{n}_{XX} are positive definite and $\sigma^2, \sigma^2_{\sigma},$ and σ^2_{σ} are positive and finite and I and IV are ninte. Furthermore, we can write ${\bf v}(\rho_{OLS})={\bf v}(\rho_{GLS})$ as a matrix product which can be shown [for instance by using results in Horn and Johnson (1985, section 7.6)] to be positive definite, in agreement with Gauss-Markov's theorem.

3.3GLS versus OLS: The one regressor case

Consider the relative efficiency of OLS and GLS in the one regressor case, $K = 1$. The results we derive below are not strictly confined to this case, however; they are also valid,

for each regression coefficient, if all regression are orthogonal so that $\Gamma \subset X$ X , $\Gamma \subset X$ X , Γ and \mathbf{R}_{XX} are diagonal matrices. Let, for $K = 1$, $b = \mathbf{B}_{XX}/\mathbf{T}_{XX}$ and $c = \mathbf{C}_{XX}/\mathbf{T}_{XX}$, *i.e.*, the shares of the total variation in the regressor (or a typical regressor, under orthogonality) which are between individual and between period variation, respectively. It then follows from (26) , (33) , and (34) that

$$
\begin{array}{rcl}\ne_{OLS} & = & \frac{\mathsf{V}(\widehat{\boldsymbol{\beta}}_{OLS})}{\mathsf{V}(\widehat{\boldsymbol{\beta}}_{GLS})} & = & \left[\sigma^2(1-b-c) + (\sigma^2 + T\sigma_\alpha^2)b + (\sigma^2 + N\sigma_\gamma^2)c\right] \\
& & \times \left[\frac{1-b-c}{\sigma^2} + \frac{b}{\sigma^2 + T\sigma_\alpha^2} + \frac{c}{\sigma^2 + N\sigma_\gamma^2}\right] \\
& = & (1-b-c)^2 + b^2 + c^2 + bc\left(\frac{\theta_B}{\theta_C} + \frac{\theta_C}{\theta_B}\right) \\
& & \quad + \left[b\left(\theta_B + \frac{1}{\theta_B}\right) + c\left(\theta_C + \frac{1}{\theta_C}\right)\right](1-b-c),\n\end{array}
$$

i.e.,

(35)
$$
e_{OLS} = \left[1 + b\left(\theta_B + \frac{1}{\theta_B} - 1\right) + c\left(\theta_C + \frac{1}{\theta_C} - 1\right)\right](1 - b - c) + b^2 + bc\left(\frac{\theta_B}{\theta_C} + \frac{\theta_C}{\theta_B}\right) + c^2.
$$

Not unexpectedly, this efficiency function is more complicated than (16) for the one-way model. In the particular case where $\sigma_B = \sigma_C = \sigma, \ i.e., \ I \ \sigma_\alpha^- = N \ \sigma_\gamma^-$, we have, however

$$
e_{OLS} = \left[1 + (b + c)\left(\theta + \frac{1}{\theta} - 1\right)\right] (1 - b - c) + (b + c)^2,
$$

which depends on θ and $b + c$ only, and resembles (16).

Since, in general,

$$
\frac{\partial e_{OLS}}{\partial \theta_B} = b(1 - b - c) \left(1 - \frac{1}{\theta_B^2} \right) + \frac{bc}{\theta_C} \left(1 - \frac{\theta_C^2}{\theta_B^2} \right),
$$
\n
$$
\frac{\partial e_{OLS}}{\partial \theta_C} = c(1 - b - c) \left(1 - \frac{1}{\theta_C^2} \right) + \frac{bc}{\theta_B} \left(1 - \frac{\theta_B^2}{\theta_C^2} \right),
$$
\n
$$
\frac{\partial e_{OLS}}{\partial b} = (1 - \theta_B) \left[(1 - 2b) \left(\frac{1}{\theta_B} - 1 \right) - c(1 - \theta_C) \left(\frac{1}{\theta_B} + \frac{1}{\theta_C} \right) \right],
$$
\n
$$
\frac{\partial e_{OLS}}{\partial c} = (1 - \theta_C) \left[(1 - 2c) \left(\frac{1}{\theta_C} - 1 \right) - b(1 - \theta_B) \left(\frac{1}{\theta_B} + \frac{1}{\theta_C} \right) \right],
$$

the efficiency function (35) has the following properties: If $0 < b < 1, 0 < c < 1$, $0 < b+c < 1$, then e_{OLS} attains its minimum, one, for $\sigma_B = \sigma_C = 1$, *t.e.*, $\sigma_{\alpha} = \sigma_{\gamma} = 0$ (which is obvious since OLS and GLS coincide in this case). If $b = 0, c = 1$ or $b = 1, c = 0$,

then $e_{OLS} = 1$ for any θ_B and θ_C . Furthermore,

$$
\theta_C = 1, \theta_B < 1, \quad 0 < b < 1 \quad \implies \quad \frac{\partial e_{OLS}}{\partial \theta_B} < 0,
$$
\n
$$
\theta_B = 1, \theta_C < 1, \quad 0 < c < 1 \quad \implies \quad \frac{\partial e_{OLS}}{\partial \theta_C} < 0,
$$
\n
$$
\theta_B = \theta_C < 1, \quad 0 < b + c < 1 \quad \implies \quad \frac{\partial e_{OLS}}{\partial \theta_B} < 0, \quad \frac{\partial e_{OLS}}{\partial \theta_C} < 0.
$$

It also follows that

$$
\frac{\partial e_{OLS}}{\partial b} > (<) 0 \Longleftrightarrow b < (>) \frac{1}{2} \left[1 - c \frac{1 - \theta_C}{1 - \theta_B} \left(1 + \frac{\theta_B}{\theta_C} \right) \right] \quad (\theta_B < 1),
$$
\n
$$
\frac{\partial e_{OLS}}{\partial c} > (<) 0 \Longleftrightarrow c < (>) \frac{1}{2} \left[1 - b \frac{1 - \theta_B}{1 - \theta_C} \left(1 + \frac{\theta_C}{\theta_B} \right) \right] \quad (\theta_C < 1),
$$
\n
$$
\frac{\partial^2 e_{OLS}}{\partial b^2} = -\frac{2}{\theta_B} (1 - \theta_B)^2,
$$
\n
$$
\frac{\partial^2 e_{OLS}}{\partial c^2} = -\frac{2}{\theta_C} (1 - \theta_C)^2,
$$
\n
$$
\frac{\partial^2 e_{OLS}}{\partial b \partial c} = -\frac{\theta_B + \theta_C}{\theta_B \theta_C} (1 - \theta_B) (1 - \theta_C).
$$

The efficiency of OLS relative to GLS therefore has the following properties: (i) When $0 < \theta_B < 1$ and $0 < \theta_C < 1$, then ϵ_{OLS} is strictly concave in b and c. (ii) When $c(1-\theta_C) = 0$, then ϵ_{OLS} attains its maximum for $\theta = \frac{1}{2}$, and when $0 \leq c(1-\theta_C) \leq 0$ $[{\sigma}{C}(1 - {\sigma}{B})]/({\sigma}{B} + {\sigma}{C})$, then ${\sigma}_{OLS}$ attains its maximum for ${\sigma} \in (0, \frac{1}{2})$. (iii) When $b(1-\theta_B) = 0$, then e_{OLS} attains its maximum for $c = \frac{1}{2}$, and when $0 \leq b(1-\theta_B) \leq 0$ $[vB(1 - vC)]/(vB + vC)$, then ε_{OLS} attains its maximum for $c \in (0, \frac{1}{2})$. In the particular case where $\theta_B = \theta_C < 1$, we have

$$
\frac{\partial e_{OLS}}{\partial (b+c)} > (0) \iff b+c < (>)\frac{1}{2},
$$

so that OLS is maximally inefficient when the sum of the between individual and the between period variation in the regressor is half of the total variation.

3.4The ranking of the within, between, and OLS estimators

We see from (50) = (33) that the ranking or μ_B , μ_C , μ_R , and μ_{OLS} in the general case is determined by $\sigma^-,$ I σ_α^- , Iv σ_γ^- , $\bm B_{XX},$ $\bm C_{XX},$ and $\bm R_{XX}.$ Again, the one regressor (or orthogonal regressors) case is most transparent, and we consider this case specically.

From (26) and (30) – (33) we obtain the following variance ratios when $K = 1$:

$$
\frac{\mathsf{V}(\hat{\beta}_{R})}{\mathsf{V}(\hat{\beta}_{B})} = \frac{b\theta_{B}}{1-b-c},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_{R})}{\mathsf{V}(\hat{\beta}_{C})} = \frac{c\theta_{C}}{1-b-c},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_{B})}{\mathsf{V}(\hat{\beta}_{C})} = \frac{c\theta_{C}}{b\theta_{B}},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_{R})}{\mathsf{V}(\hat{\beta}_{OLS})} = \frac{1}{(1-b-c)\left[1+b\left(\frac{1}{\theta_{B}}-1\right)+c\left(\frac{1}{\theta_{C}}-1\right)\right]},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_{B})}{\mathsf{V}(\hat{\beta}_{OLS})} = \frac{1}{b\theta_{B}\left[1+b\left(\frac{1}{\theta_{B}}-1\right)+c\left(\frac{1}{\theta_{C}}-1\right)\right]},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_{C})}{\mathsf{V}(\hat{\beta}_{OLS})} = \frac{1}{c\theta_{C}\left[1+b\left(\frac{1}{\theta_{B}}-1\right)+c\left(\frac{1}{\theta_{C}}-1\right)\right]}.
$$

Rearranging these expressions, defining $\tau_{\alpha} = \sigma_{\alpha}^2/\sigma$ and $\tau_{\gamma}^2 = \sigma_{\gamma}^2/\sigma$, and using (20), we find

$$
\frac{\mathsf{V}(\hat{\beta}_R)}{\mathsf{V}(\hat{\beta}_B)} > (\langle 1 \rangle \iff \frac{1 - b - c}{1 - c} < (\rangle) \frac{1}{2 + T\tau_{\alpha}^2},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_R)}{\mathsf{V}(\hat{\beta}_C)} > (\langle 1 \rangle \iff \frac{1 - b - c}{1 - b} < (\rangle) \frac{1}{2 + N\tau_{\gamma}^2},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_C)}{\mathsf{V}(\hat{\beta}_B)} > (\langle 1 \rangle \iff \frac{b}{c} > (\langle 1 \rangle) \frac{1 + T\tau_{\alpha}^2}{1 + N\tau_{\gamma}^2},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_R)}{\mathsf{V}(\hat{\beta}_{OLS})} > (\langle 1 \rangle \iff 1 - b - c < (\rangle) \frac{1}{1 + bT\tau_{\alpha}^2 + cN\tau_{\gamma}^2},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_B)}{\mathsf{V}(\hat{\beta}_{OLS})} > (\langle 1 \rangle \iff b < (\rangle) \frac{1 + T\tau_{\alpha}^2}{1 + bT\tau_{\alpha}^2 + cN\tau_{\gamma}^2},
$$
\n
$$
\frac{\mathsf{V}(\hat{\beta}_C)}{\mathsf{V}(\hat{\beta}_{OLS})} > (\langle 1 \rangle \iff c < (\rangle) \frac{1 + N\tau_{\alpha}^2 + cN\tau_{\gamma}^2}{1 + bT\tau_{\alpha}^2 + cN\tau_{\gamma}^2}.
$$

From these expressions, we can rank the five estimators by relative efficiency.

We do not describe the detailed ranking, as there are a substantial number of possible cases and some parameter constellations are more likely to occur than others. In principle, there is a region in the $(v, c, T\tau_{\alpha}^*, N\tau_{\gamma}^-)$ space in which $\bm{\beta}_R$ is superior to $\bm{\beta}_B, \bm{\beta}_B$ is superior to ρ_C, ρ_R is superior to ρ_{OLS} , etc. Genuine panel data from individuals, households, or firms often show substantial individual specific heterogeneity, both in the regressor and in the disturbances, and less pronounced period specific heterogeneity, so that b often by far exceeds $c, \, 1-b-c$ is small (but often larger than c), and σ_{α}^- exceeds σ_{γ}^- . Furthermore,

N is often considerably larger than T. Four realistic cases may then be

$$
V(\hat{\beta}_{C}) > V(\hat{\beta}_{B}) > V(\hat{\beta}_{OLS}) > V(\hat{\beta}_{R}) > V(\hat{\beta}_{GLS}) \text{ if}
$$

\n
$$
1 < \frac{1}{1 - b - c} < 1 + bT\tau_{\alpha}^{2} + cN\tau_{\gamma}^{2} < \frac{1 + T\tau_{\alpha}^{2}}{b} < \frac{1 + N\tau_{\gamma}^{2}}{c},
$$

\n
$$
V(\hat{\beta}_{C}) > V(\hat{\beta}_{B}) > V(\hat{\beta}_{R}) > V(\hat{\beta}_{OLS}) > V(\hat{\beta}_{GLS}) \text{ if}
$$

\n
$$
1 < 1 + bT\tau_{\alpha}^{2} + cN\tau_{\gamma}^{2} < \frac{1}{1 - b - c} < \frac{1 + T\tau_{\alpha}^{2}}{b} < \frac{1 + N\tau_{\gamma}^{2}}{c},
$$

\n
$$
V(\hat{\beta}_{C}) > V(\hat{\beta}_{R}) > V(\hat{\beta}_{B}) > V(\hat{\beta}_{OLS}) > V(\hat{\beta}_{GLS}) \text{ if}
$$

\n
$$
1 < 1 + bT\tau_{\alpha}^{2} + cN\tau_{\gamma}^{2} < \frac{1 + T\tau_{\alpha}^{2}}{b} < \frac{1}{1 - b - c} < \frac{1 + N\tau_{\gamma}^{2}}{c},
$$

\n
$$
V(\hat{\beta}_{R}) > V(\hat{\beta}_{C}) > V(\hat{\beta}_{B}) > V(\hat{\beta}_{OLS}) > V(\hat{\beta}_{GLS}) \text{ if}
$$

\n
$$
1 < 1 + bT\tau_{\alpha}^{2} + cN\tau_{\gamma}^{2} < \frac{1 + T\tau_{\alpha}^{2}}{b} < \frac{1 + N\tau_{\gamma}^{2}}{c} < \frac{1}{1 - b - c}.
$$

Let us consider an *empirical illustration* taken from Biørn (1994, Table 8), in which marginal budget shares for 28 disaggregate consumption commodities (exhausting the complete budget) and their variances are estimated from Norwegian household panel data with $K = 1, N = 418$ and $T = 2$. The only regressor variable is total expenditure which corresponds to x . Its shares of the total variation which is between individual. between period, and residual variation in this data set are $b = 80.5\%$, $c = 4.2\%$, and $1 - b - c = 15.3 %$, respectively. (Values of b and c of similar magnitudes are often found for logarithms of outputs and inputs in firm data.) The variance components o , o_{α}^- and σ_γ^- are commodity specific and are estimated consistently from residuals, as explained in Bigrn $(1994, p. 142)$. We find from these estimates of the variance components that the first of the above four sets of inequalities is satisfied for one commodity (tobacco), the second for one commodity (fuel and power), and the third for the remaining 26 commodities (including foods, services, housing, durables etc.). It should come as no surprise that the first inequality is the one to hold for tobacco, since this is a strongly addictive commodity whose value of τ_{α}^- is substantially larger than for other commodities. For none of the 28 commodities the fourth set of inequalities is satisfied according to the estimated variance components.

For this two-way, unlike the one-way, model it is possible for the between individual estimator to be more emclent than OLS. This will happen when $cN \tau_\gamma^-$ is so large that $b > (1 + T i_{\alpha}^2) / (1 + 0T i_{\alpha}^2 + C N i_{\gamma}^2)$ [if $i_{\alpha}^2 = 0$ or $c = 0$, and $b < 1$, we always have $V(B_B) > 0$ $V(\rho_{OLS})$. Likewise, it is possible for the between period estimator to be more enficient than OLS. This will happen when bI τ_{α}^- is so large that $c > (1 + N \tau_{\alpha}^-) / (1 + bI \tau_{\alpha}^- + cN \tau_{\alpha}^-)$ μ τ_{α} = 0 or θ = 0, and c < 1, we always have ${\bf v}(\rho_C) > {\bf v}(\rho_{OLS})$. Netther of these

situations occur in our marginal budget shares example, however. In fact, the between period estimator has by far the lowest estimated eciency for all the 28 commodities. The OLS estimator is ranked second for all commodities except one (tobacco), for which it is ranked third and the residual estimator is ranked second.

4 Concluding remarks

Although it is well established that the GLS is the optimal estimator of the coefficient vector in random effects panel data regression models, when the model is correctly specied, we can conclude from the results in this paper that both OLS and various between and within estimators may be of interest for practical purposes. First, in many realistic situations, the estimation efficiency may not be much improved by using GLS instead of one of its competitors. Second, GLS depends on variance components which are rarely available and may be estimable only with substantial margins of errors, so that the Feasible GLS may depart substantially from the strict GLS. Other panel data estimators do not require this kind of information, although it is needed in order to estimate their variances correctly. Third, the consistency of GLS, like OLS, may be vulnerable to model specification errors. For instance, the consistency of within estimators is robust to correlation between the random latent effects and the covariate vector, and the consistency of the between individual (between period) estimator is robust to errors of measurement in the regressors when the number of periods (individuals) goes to infinity.

In the paper, we have, for both the one-way and the two-way random effects models, reconsidered on the one hand the efficiency of the GLS over its competitors, on the other hand the mutual efficiency of the OLS and the within and between estimators. A detailed investigation has been done for the one regressor (or the orthogonal regressors) case, in which the efficiency can be characterized by variance ratios. A further examination of the multiple (non-orthogonal) regressor case is left for future research. Of course, the precise ranking of the estimators is indeterminate unless the disturbance variance components are known. Still, our results may give guidelines about which estimator of the coefficient vector to choose when the relative composition of the (co)variation of the regressors into within and between (co)variation is known and we have estimates of disturbance variance component ratios, for instance obtained from OLS residuals, even if they are inaccurate.

Acknowledgements. The results presented in this note had its origin in lecture notes which I prepared for courses in panel data econometrics at the University of Oslo. I thank Terje Skjerpen for valuable comments.

References

- Baltagi, B.H. (1981): Pooling: An Experimental Study of Alternative Testing and Estimation Procedures in a Two-Way Error Component Model. Journal of Econometrics, 17 (1981), $21 - 49$.
- Birn, E. (1994): Moment Estimators and the Estimation of Marginal Budget Shares from Household Panel Data. Structural Change and Economic Dynamics, 5 (1994), $133 - 154.$
- Birn, E. (1996): Panel Data with Measurement Errors. Chapter 10 in: Matyas, L., and Sevestre, P. (eds.): The Econometrics of Panel Data. A Handbook of the Theory with Applications. Dordrecht: Kluwer, 1996.
- Fuller, W.A., and Battese, G.E. (1974): Estimation of Linear Models with Crossed-Error Structure. Journal of Econometrics, 2 (1974), $67 - 78$.
- Greene, W.H. (2000): *Econometric Analysis*, Fourth Edition. London: Prentice Hall, 2000.
- Horn, R.A., and Johnson, C.R. (1985): *Matrix Analysis*. Cambridge: Cambridge University Press, 1985.
- Hsiao, C. (1986): Analysis of Panel Data. Cambridge: Cambridge University Press, 1986.
- Maddala, G.S., and Mount, T.D. (1973): A Comparative Study of Alternative Estimators for Variance Components Models Used in Econometric Applications. Journal of the American Statistical Association, 68 (1973), $324 - 328$.
- Magnus, J.R., and Neudecker, H. (1988): *Matrix Differential Calculus with Applications* in Statistics and Econometrics. Chichester: Wiley, 1988.
- Malinvaud, E. (1978): Methodes Statistiques de l'Ec onometrie, 3eme Edition. Paris: Dunod, 1978.
- Matyas, L. (1996): Error Components Models. Chapter 4 in: Matyas, L., and Sevestre, P. (eds.): The Econometrics of Panel Data. A Handbook of the Theory with Applications. Dordrecht: Kluwer, 1996.
- Swamy, P.A.V.B., and Arora, S.S. (1972): The Exact Finite Sample Properties of the Estimators of Coefficients in the Error Components Regression Models. *Econometrica*, 40 (1972), $261 - 275$.
- Taylor, W.E. (1980): Small Sample Considerations in Estimation from Panel Data. Journal of Econometrics, 13 (1980), 203 -223 .